

hep-ph/0212305

Cosmological Baryon Asymmetry and Neutrinos: Baryogenesis via Leptogenesis in Supersymmetric Theories

Koichi Hamaguchi

Deutsches Elektronen-Synchrotron DESY, D-22603, Hamburg, Germany

*Dr.thesis submitted to
Department of Physics, University of Tokyo
January 2002*

Ph.D thesis

**Cosmological Baryon Asymmetry and Neutrinos:
Baryogenesis via Leptogenesis in Supersymmetric Theories**

Koichi Hamaguchi

Department of Physics, University of Tokyo, Tokyo 113-0033, Japan

Submitted to Department of Physics, University of Tokyo: *January 2002*

Accepted: *February 2002*

Revised to submit to the eprint arXiv: *December 2002*

Contents

1	Introduction	1
1.1	Overview	1
1.2	Outline of this thesis	5
1.3	Baryon asymmetry	6
1.4	Sphaleron	7
1.5	Cosmological gravitino problems	14
2	Leptogenesis by the decay of right-handed neutrino	18
2.1	Asymmetric decay of the right-handed neutrino	19
2.2	Leptogenesis by thermally produced right-handed neutrino	23
2.3	Leptogenesis in inflaton decay	30
2.3.1	Lepton asymmetry	30
2.3.2	Froggatt-Nielsen model for neutrino	32
2.3.3	Hybrid inflation	35
2.3.4	New inflation	45
2.3.5	Topological inflation	54
2.3.6	Production of right-handed neutrinos at preheating	57
2.4	Leptogenesis from $\widetilde{N}$ -dominant universe	60
2.4.1	Lepton asymmetry from $\widetilde{N}$ -dominant universe	60
2.4.2	Necessary conditions	62
2.4.3	Dilution of the gravitinos	64
2.4.4	Remarks	67

2.A	N -dominant universe from inflaton decay	74
3	Leptogenesis via LH_u flat direction	75
3.1	Overview	76
3.2	Lepton asymmetry from LH_u flat direction	78
3.2.1	The LH_u flat direction	78
3.2.2	Initial amplitude	79
3.2.3	Evolution of the flat direction	80
3.2.4	Lepton asymmetry	87
3.3	Some models	95
3.3.1	Model for the ultralight neutrino with a broken discrete symmetry .	95
3.3.2	Large effective masses for right-handed neutrinos	98
3.4	Effects of the gauged $U(1)_{B-L}$ symmetry	102
3.4.1	The model	103
3.4.2	Initial amplitude and initial phase	105
3.4.3	Lepton asymmetry	110
3.4.4	Conditions to avoid an early oscillation	115
3.5	The neutrino-less double-beta decay	117
3.A	D -flat condition for the LH_u flat direction	121
4	Conclusions and discussion	122
A	Notations	128
B	Some notes of standard cosmology	129
B.1	Friedmann-Robertson-Walker universe	129
B.2	Thermodynamics	131

Chapter 1

Introduction

1.1 Overview

The origin of baryon asymmetry (matter-antimatter asymmetry) in the present universe is one of the fundamental puzzles in particle physics as well as in cosmology. Assuming an inflationary phase in the early universe, any initial baryon asymmetry is diluted and becomes essentially zero during the inflation. Therefore, the observed baryon asymmetry should be generated dynamically after the inflation. Such a dynamical generation of baryon asymmetry (baryogenesis) is possible if the following three conditions are satisfied; (i) baryon-number violation, (ii) C - and CP -violations, and (iii) departure from thermal equilibrium, which are known as the Sakharov's three conditions [1].

The original idea of the baryogenesis in the grand unified theory (GUT) [2] satisfies all of these conditions [3], and provides an elegant interplay between particle physics and cosmology. Actually, the GUT [4] unifies baryons and leptons in the same gauge multiplets and predicts the existence of baryon number violation, the delayed decay of superheavy GUT particle satisfies the out-of-equilibrium condition, and its CP -violating asymmetric decay into baryons and antibaryons can offer a net baryon asymmetry.

On the other hand, the electroweak gauge theory itself violates the baryon asymmetry by a quantum anomaly [5]. This effect is highly suppressed by a large exponential factor at zero temperature, and hence the stability of the proton is practically guaranteed. However, it was found that in the early universe at high temperatures above the electroweak scale, the baryon-number violating interaction is not suppressed and even be rapid enough to be in thermal equilibrium [6]. An important point here is that this baryon number violating (“sphaleron”) process violates lepton (L) number as well as baryon (B) number, and it conserves a linear combination of them, $B - L$. Therefore, if the baryon asymmetry is produced in $(B - L)$ -conserving processes, like those in the $SU(5)$ GUT, it would be

washed out by the sphaleron effect before the electroweak phase transition, and hence no baryon asymmetry can remain until the stage of big-bang nucleosynthesis to generate the light nuclei.

The existence of the sphaleron effect opens a new possibility of baryogenesis, the generation of baryon asymmetry at electroweak phase transition [6, 7]. This mechanism does not use any $(B - L)$ -violation, and the baryon asymmetry is generated by the sphaleron-induced $(B + L)$ -violating process. (Hence, same amount of baryons and leptons are produced.) As other baryogenesis scenarios have, this mechanism also has a double-edged behavior, i.e., the produced baryon (and lepton) asymmetry tends to be washed out afterwards by the sphaleron process itself. In order to avoid the erasure of the produced baryon asymmetry, the electroweak phase transition must be strongly first-order. However, in the case of the minimal standard model with one Higgs doublet, it turns out that, for Higgs masses allowed by current experimental bounds, the electroweak transition is too weakly first order or just a smooth transition, which excludes the electroweak baryogenesis in the standard model. Although the extension of the standard model to the supersymmetric version may cure this difficulty, there remains only a small parameter range [8].

Therefore, it seems natural to consider that the baryon asymmetry was generated before the electroweak phase transition. As long as we consider a baryogenesis above the electroweak scale, some $(B - L)$ -violating interaction is mandatory, since the sphaleron process washes out any baryon asymmetry in the universe unless there exists nonzero $B - L$ asymmetry. (Namely, the first one of the Sakharov's conditions, (i) B -violation, should be replaced with (i)' $(B - L)$ -violation.) This means, on the other hand, as first suggested by Fukugita and Yanagida [9], that the *lepton*-number violation is enough for baryogenesis and explicit baryon-number violation is not necessarily required. Actually, there now exists an implication of lepton-number (L) violation, whereas there has been discovered no evidence for baryon-number (B) violations: that is, the neutrino oscillation.

Neutrino oscillation [10, 11],¹ especially the strong evidence for the atmospheric neutrino oscillation reported in 1998 by the Super-Kamiokande Collaboration [10], is one of the greatest discoveries in particle physics after the success of the standard model. The experimental data strongly suggest that the neutrinos have small but finite masses, impelling the standard model of particle physics to be modified.

Such tiny masses of light neutrinos are naturally understood if we introduce heavy right-handed Majorana neutrinos to the standard model, in terms of so-called seesaw mechanism [13]. Because the right-handed neutrinos N are singlets under the standard

¹Very recently the KamLAND experiment has announced the first results, which exclude all oscillation solutions but the 'Large Mixing Angle' solution to the solar neutrino problem [12].

gauge symmetries, they can have Majorana masses as well as Yukawa couplings to the left-handed leptons (l) and Higgs (φ) doublets:²

$$\mathcal{L} = -\frac{1}{2}M N^T N + \left(-h N^T l \varphi + \text{H.c.}\right), \quad (1.1)$$

where we have taken the N to be the Majorana mass eigenstates. Then by integrating out the heavy right-handed neutrinos, we obtain the small Majorana masses for light neutrinos, $m_\nu = h^2 \langle \varphi \rangle^2 / M$.

A crucial observation in the lagrangian Eq. (1.1) is that the Majorana mass term MN^TN represents nothing but a lepton number (L) violation, actually a $(B-L)$ -violation. In fact, the heavy right-handed neutrinos have two distinct decay channels into leptons and anti-leptons, and produce $B-L$ asymmetry if the Yukawa couplings h violate the CP and if the decay is out of thermal equilibrium. The produced lepton asymmetry is partially converted into baryon asymmetry via the aforementioned sphaleron effect, which explains the baryon asymmetry in the present universe. This is the original idea of the leptogenesis [9].

We should also note that if we assume a gauged $U(1)_{B-L}$ symmetry, the existence of three generations of the right-handed neutrinos are automatically required in order to cancel the gauge anomaly. As well known, the $U(1)_{B-L}$ symmetry is the unique extra $U(1)$ symmetry which can be gauged consistently with the standard model. Furthermore, the breaking of this $U(1)_{B-L}$ symmetry naturally provides large Majorana masses to the right-handed neutrinos, which leads to the tiny neutrino masses via the seesaw mechanism. It should be noticed that the $U(1)_{B-L}$ symmetry is consistent with the $SU(5)$ GUT, and is embedded in larger GUT groups such as $SO(10)$.

Meantime, the supersymmetry (SUSY) [14] has been attracting wide interests in particle physics as one of the best candidates for new physics beyond the standard model. It protects the huge hierarchy between the electroweak scale and unification (or Planck) scale against the quadratically divergent radiative corrections, and the particle contents of the minimal SUSY standard model (MSSM) lead to a beautiful unification of the three gauge couplings of the standard model at the scale $\simeq 2 \times 10^{16}$ GeV [15], which strongly suggests the SUSY GUT [16]. The MSSM also gives a natural framework to break the electroweak symmetry radiatively [17]. On the other hand, from the viewpoint of cosmology, SUSY provides an ideal dark matter candidate, the lightest SUSY particle [18]. It also protects the flatness of the inflaton potential against the radiative corrections, which is inevitable for successful inflation.

²Here, we have omitted the family- and $SU(2)_L$ -indices for simplicity.


However, SUSY also causes a cosmological difficulty if the reheating temperature of the inflation is too high, that is, the cosmological gravitino problems [19, 20]. After the end of the inflation, the gravitinos are produced by scattering processes of particles from the thermal bath, and its abundance is proportional to the reheating temperature. Because the gravitino's interaction is suppressed by the gravitational scale, it has a very long lifetime unless it is completely stable, and its decay during or after the big-bang nucleosynthesis (BBN) epoch ($t \sim 1\text{--}100$ sec) might spoil the success of the BBN [19, 21]. On the other hand, if the gravitino is completely stable, its present mass density must be below the critical density of the present universe [20]. In both cases, the primordial gravitino abundance should be low enough, and hence the reheating temperature of the inflation is severely constrained from above, depending on the gravitino mass.

In this thesis, we study in detail several leptogenesis scenarios in the framework of the SUSY. There have been proposed, in fact, various leptogenesis scenarios depending on the production mechanisms of the right-handed neutrinos:

- In the simplest and the most conventional leptogenesis mechanism, the right-handed neutrino is produced by thermal scatterings [9, 22]. The delayed decay of the right-handed neutrino satisfies the out-of-equilibrium condition à la the original GUT baryogenesis. This mechanism requires a relatively high reheating temperature to produce the heavy right-handed neutrino, and the aforementioned gravitino problem makes it somewhat difficult, depending on the gravitino mass.
- Another mechanism is given when the right-handed neutrinos are produced non-thermally in inflaton decay [23]. In this scenario the reheating temperature can be lower than the case of thermal production, and the gravitino problem is avoided in a wider range of gravitino mass.
- The third mechanism is inherent in the SUSY. The lepton asymmetry is produced by the decay of coherent oscillation of the right-handed “s” neutrino, which is the supersymmetric scalar partner of the right-handed neutrino [24, 25]. If the right-handed sneutrino's oscillation dominates the energy density of the universe, the gravitino problem is drastically ameliorated [26].

In the framework of SUSY, there is yet another, completely different mechanism. The lepton asymmetry is produced not by the right-handed neutrino decay, but by a coherent oscillation (actually, a rotation) of a flat direction field including the lepton doublet L :

- The leptogenesis via LH_u flat direction [25], which is based on the Affleck-Dine mechanism [27].



1.2 Outline of this thesis

The outline of this thesis is as follows. The rest of this chapter is devoted to some reviews. We briefly mention the observed baryon asymmetry in Sec. 1.3. A review of the sphaleron effect, which is a crucial ingredient of the leptogenesis, is given in Sec. 1.4. We also briefly review the results of the cosmological gravitino problems in Sec. 1.5.

In Chapter 2, we discuss leptogenesis scenarios by the decay of the right-handed (s)neutrino. First, we study the asymmetric decay of the right-handed neutrino into leptons and anti-leptons in Sec. 2.1. The conventional leptogenesis mechanism where the right-handed neutrinos are thermally produced is briefly discussed in Sec. 2.2. Then we perform a comprehensive study of the leptogenesis mechanism in inflaton decay in Sec. 2.3, adopting various SUSY inflation models. In Sec. 2.4, we investigate the leptogenesis from coherent right-handed sneutrino.

In the latter half of this thesis, in Chapter 3, we perform a detailed analysis on the leptogenesis via LH_u flat direction. This scenario may require another overview, which will be given in Sec. 3.1. Here we mention one point, that the most important parameter in this scenario which determines the baryon asymmetry is the mass of the *lightest* neutrino, $m_{\nu 1}$. (Notice that the data from neutrino-oscillation experiments [10, 11, 12] suggest the difference of the neutrino mass squared, indicating the masses of the heavier neutrinos, $m_{\nu 2}$ and $m_{\nu 3}$.) It is amazing that the observed baryon asymmetry, which was generated in the very early universe, is directly related to such a low-energy physics, the neutrino mass.

Chapter 4 is devoted to conclusions and discussion. Some of our notations are summarized in Appendix A. Some ingredients of the standard cosmology are reviewed very briefly in Appendix. B.

1.3 Baryon asymmetry

We will use the following value for the baryon asymmetry in the present universe [28, 29]:

$$\frac{n_B}{s} \equiv \frac{n_b - n_{\bar{b}}}{s} = (0.4 - 1) \times 10^{-10}, \quad (1.2)$$

where n_b and $n_{\bar{b}}$ are baryon and anti-baryon number density, respectively, and s is the entropy density of the universe. This ratio takes a constant value, as long as the baryon number is conserved and no entropy production takes place. Notice that $n_{\bar{b}} \ll n_b$ and hence $n_B \simeq n_b$ in the present universe. (We sometimes call the n_B just “baryon number density,” for simplicity.) This value is determined from the big-bang nucleosynthesis (BBN). The BBN occurs at the temperature of $T \simeq (1-0.1)$ MeV, or equivalently at the cosmic time $t \sim (1-100)$ sec, and generates light nuclei, D, ^3He , ^4He , and ^7Li . (For reviews, see, for example, Refs. [28, 29].) The agreement between the predictions of the BBN theory for the abundances of these light nuclei and the primordial abundances of them (which are inferred from observational data) is one of the most important successes of the standard big-bang cosmology. Actually, the theory of the BBN has basically only one free parameter,³ i.e., the baryon asymmetry, and all the above light-element abundances are well explained with n_B/s in the range of Eq. (1.2).

To demonstrate the smallness of the baryon asymmetry, let us calculate the baryon-number and anti-baryon number density at high temperature. For example, consider a temperature between the QCD phase transition and electroweak phase transition, say, $T \sim 10$ GeV. At this temperature, all the baryons and anti-baryons are expected to exist as quarks and anti-quarks, and they are well in thermal equilibrium. The ratio of the baryon-number and anti-baryon-number densities to the entropy density are then given by

$$\frac{n_b}{s} \simeq \frac{n_{\bar{b}}}{s} \simeq 0.02, \quad (1.3)$$

where we have used Eq. (B.19). Compared with Eq. (1.2), this means, $(n_b - n_{\bar{b}})/(n_b + n_{\bar{b}}) \sim \mathcal{O}(10^{-9})$, i.e., there were (1,000,000,000+1) baryons per 1,000,000,000 anti-baryons at this epoch.

³We assume the number of neutrino generations to be three. As well known, this fact itself is one of the most important implications of the BBN.

1.4 Sphaleron

In this section, we give a brief review of the “sphaleron” effect, which is a crucial ingredient of the leptogenesis. If the lepton asymmetry is successfully generated, it is partially converted into the baryon asymmetry at equilibrium thanks to the sphaleron-induced baryon and lepton number violating process, which then explains the observed baryon asymmetry in the present universe.

At the classical level, the Lagrangian of the standard $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge theory clearly conserves the baryon (B) and lepton (L) numbers. However, as was first pointed out by 't Hooft [5], both B and L are violated by quantum effects:

$$\partial_\mu J_B^\mu = \partial_\mu J_L^\mu = N_f \frac{g_2^2}{32\pi^2} \epsilon_{\mu\nu\rho\sigma} \text{Tr} F^{\mu\nu} F^{\rho\sigma}, \quad (1.4)$$

where N_f is the number of fermion generations and g_2 and $F_{\mu\nu}$ are the coupling constant and the field strength of the $SU(2)_L$ gauge group, respectively. It was shown later that the $SU(2)_L \times U(1)_Y$ gauge theory has a non-contractible loop (path) in the field configuration space, which connects topologically distinct vacua with different baryon and lepton numbers [30], and it was found that the highest energy configuration along this “path” corresponds to a spatially localized and static, but unstable solution [31, 32]. This solution was named “sphaleron”. Since this sphaleron solution is a saddle point of the field potential energy, its energy represents the height of the barrier between the vacua with different baryon and lepton numbers (see Fig. 1.1).

At zero temperature, the rate of baryon (and lepton) number violating process via a tunneling between the topologically distinct vacua (for example, proton decay) is extremely small [5], since it is suppressed by the factor of $\exp(-16\pi^2/g_2^2) \sim 10^{-170}$. However, as suggested by Kuzmin, Rubakov, and Shaposhnikov [6, 33], such processes are not suppressed and can even be efficient at temperatures close to (and above) the electroweak phase transition. The transition of the fermion number occurs simultaneously for each fermion doublets, so that the total changes in the numbers of baryons and leptons are $\Delta B = \Delta L = N_f$. Therefore, $B - L$ is conserved. (This is clear from the absence of the anomaly in $B - L$ in Eq. (1.4).)

Let us roughly estimate the temperature T_* above which this $(B+L)$ -violating process becomes in thermal equilibrium, according to Ref. [6]. We assume that the electroweak phase transition is the second order, and consider the temperature $T < T_C$, where T_C is the critical temperature above which the vacuum expectation value $v(T)$ of the Higgs field vanishes. In such a situation, the transition from one vacuum (say, $B = b_0$ and $L = l_0$)




Figure 1.1: A Schematic behavior of the energy dependence on the configuration of the gauge and Higgs fields $[\mathbf{A}(x), \varphi(x)]$ [6]. The minima correspond to topologically distinct vacua with different baryon (B) and lepton (L) numbers. The configuration $[\mathbf{A}_{\text{sph}}(x), \varphi_{\text{sph}}(x)]$ represents the saddle point of the energy functional, the sphaleron solution.

to the next vacuum ($B = b_0 \pm N_f$ and $L = l_0 \pm N_f$) occurs at the rate [6]

$$\Gamma = C(T) T \exp \left(- \frac{M_{\text{sph}}(T)}{T} \right), \quad (1.5)$$

where dimensionless factor $C(T)$ depends on the ratio $v(T)/T$ and the coupling constants.⁴ $M_{\text{sph}}(T)$ represents the free energy of the sphaleron configuration (at temperature T), which is given by [31]

$$M_{\text{sph}}(T) = 4\pi B(T) \frac{v(T)}{g_2(T)}, \quad (1.6)$$

where $B(T)$ depends on the gauge coupling $g_2(T)$ and the 4-point coupling constant of the Higgs potential $\lambda(T)$ as $B = B(\lambda/g_2^2)$, varying from 1.5 ($\lambda/g_2^2 \rightarrow 0$) to 2.7 ($\lambda/g_2^2 \rightarrow \infty$) [31]. The rate in Eq. (1.5) should be compared with the Hubble expansion rate $H = (\pi^2 g_*/90)^{1/2} \times T^2/M_G$. ($M_G = 2.4 \times 10^{18}$ GeV is the reduced Planck scale and g_* is defined in Appendix B.2.) Then, it is found that the sphaleron rate in Eq. (1.5) indeed exceeds the Hubble expansion rate for $T > T_*$, where T_* is given by

$$T_* \simeq 4\pi B(T_*) \frac{v(T_*)}{g_2(T_*)} \times \left[\ln \left(\frac{M_G}{T_*} \right) \right]^{-1}. \quad (1.7)$$

⁴See comments below.

Here, we have neglected the $\ln C(T)$ and $\ln(\pi^2 g_*/90)$ compared with the large number $\ln(M_G/T_*)$. Recalling that $v(T)$ approaches to zero for $T \rightarrow T_C$, this occurs below the critical temperature, i.e., $T_* < T_C$. Therefore, the sphaleron-induced $(B+L)$ -violating process becomes in equilibrium for $T_* < T (< T_C)$. Precisely speaking, a more careful treatment including the prefactor $C(T)$, which turns out to be proportional to $(v(T)/T)^7$ [34], is required. A numerical estimation shows that T_* is just below the critical temperature, $(T_C - T_*) \lesssim 10$ GeV, for $50 \text{ GeV} \lesssim T_C \lesssim 200 \text{ GeV}$ [35].

The evaluation of the “sphaleron” rate for $T > T_C$ is a complicated problem, since the sphaleron configuration no longer exists in the symmetric phase.⁵ Naively thinking, there seems no reason for the $(B+L)$ -violating processes to be suppressed, since the potential barrier vanishes. Actually, theoretical arguments as well as numerical calculations suggest [36] that the sphaleron rate per unit time per unit volume is $\Gamma/V \sim \alpha_2^5 T^4$ for $T > T_C$, where $\alpha_2 \equiv g_2^2/(4\pi)$. By using their result, it is found that the sphaleron rate exceeds the Hubble expansion rate for $T \lesssim 10^{12}$ GeV.

Therefore, the sphaleron-induced $(B+L)$ -violating process is in thermal equilibrium in the range of

$$100 \text{ GeV} \sim T_* < T \lesssim 10^{12} \text{ GeV}. \quad (1.8)$$

• relation between baryon and lepton asymmetry

Let us calculate the relation between the baryon and lepton number in the presence of the sphaleron process, by means of the analysis of the chemical potentials [37, 38]. At first sight, it seems that the relation is just given by $B+L=0$ since the sphaleron effect violates $B+L$, preserving $B-L$. However, as we will see, a nontrivial relation is derived if there exists a non-vanishing $B-L$ asymmetry.

We first consider the standard model without supersymmetric particles, but with N_φ Higgs doublets, φ_i ($i = 1 \cdots N_\varphi$), and N_f generations of fermions, i.e., left-handed quark doublets q_j , right-handed up-type and down-type quarks $\bar{u}_j$ and $\bar{d}_j$, left-handed lepton doublets l_j and right-handed charged leptons $\bar{e}_j$ ($j = 1 \cdots N_f$). Hereafter, we consider the symmetric phase $T > T_C$. Hence, the $\text{SU}(2)_L \times \text{U}(1)_Y$ symmetry recovers. We denote the asymmetry of the number density of particle i by Δn_i , which is related to the chemical

⁵Although the saddle-point sphaleron solution does not exist in the symmetric phase, we will call this anomalous $(B+L)$ -violating process “sphaleron” also for $T > T_C$.

potential of that particle as follows [see Eq. (B.20)]:

$$\Delta n_i \equiv n_i^{(+)} - n_i^{(-)} = \begin{cases} \frac{1}{6} g_i T^3 \left(\frac{\mu_i}{T} \right) & \text{for fermion,} \\ \frac{1}{3} g_i T^3 \left(\frac{\mu_i}{T} \right) & \text{for boson.} \end{cases} \quad (1.9)$$

Therefore, the baryon and lepton number asymmetries are given by

$$\begin{aligned} n_B &\equiv \sum_j \left(\frac{1}{3} \Delta n_{q_j} - \frac{1}{3} \Delta n_{\bar{u}_j} - \frac{1}{3} \Delta n_{\bar{d}_j} \right) \\ &= \frac{1}{6} T^2 \times \sum_j \left(2\mu_{q_j} - \mu_{\bar{u}_j} - \mu_{\bar{d}_j} \right), \\ n_{L_j} &\equiv \Delta n_{l_j} - \Delta n_{\bar{e}_j} \\ &= \frac{1}{6} T^2 \times \left(2\mu_{l_j} - \mu_{\bar{e}_j} \right), \end{aligned} \quad (1.10)$$

where we have used the fact that the chemical potentials of particles in the same gauge multiplet are the same, which is ensured by the gauge interactions. (Note that the $SU(2)_L$ is also recovered, since we consider $T > T_C$.)

In the following, we derive the relations between the chemical potentials μ_i . First of all, we assume that all the Higgs doublets have the same chemical potentials due to the mixings between themselves, i.e., $\mu_{\varphi_i} \equiv \mu_{\varphi}$. (If there is a Higgs doublet with a conjugate quantum number, like in the MSSM, we redefine the chemical potential of that Higgs doublet with an additional minus sign.) Then, the interactions via Yukawa couplings

$$y_{e,j} \varphi^* l_j \bar{e}_j, \quad y_{u,jk} \varphi q_j \bar{u}_k, \quad y_{d,jk} \varphi^* q_j \bar{d}_k, \quad (1.11)$$

lead to

$$\begin{aligned} -\mu_{\varphi} + \mu_{l_j} + \mu_{\bar{e}_j} &= 0 & j = 1 \cdots N_f, \\ \mu_{\varphi} + \mu_{q_j} + \mu_{\bar{u}_k} &= 0 & j, k = 1 \cdots N_f, \\ -\mu_{\varphi} + \mu_{q_j} + \mu_{\bar{d}_k} &= 0 & j, k = 1 \cdots N_f. \end{aligned} \quad (1.12)$$

Here, we have taken a basis of gauge eigenstates for the quarks. Thus, because of the mixing in the Yukawa couplings, the chemical potentials of the quarks become generation independent: $\mu_{q_j} \equiv \mu_q$, $\mu_{\bar{u}_j} \equiv \mu_{\bar{u}}$ and $\mu_{d_j} \equiv \mu_{\bar{d}}$ ($j = 1 \cdots N_f$). Notice that the relations in Eq. (1.12) hold as long as these interactions are in thermal equilibrium, and even the electron Yukawa coupling, which is the smallest one, is in equilibrium for $T \lesssim 10^4$ GeV. Next, the charge neutrality of the universe requires vanishing total $U(1)_Y$ charge:

$$\sum_i (\Delta n_i Y_i) = 0, \quad (1.13)$$

where Y_i denotes the $U(1)_Y$ charge of particle i . This leads to

$$N_f (\mu_q - 2\mu_{\bar{u}} + \mu_{\bar{d}}) + \sum_j (-\mu_{l_j} + \mu_{\bar{e}_j}) + 2N_\varphi \mu_\varphi = 0. \quad (1.14)$$

Finally, the sphaleron interaction can be understood as

$$\sum_{\substack{i = \text{all SU}(2)_L \\ \text{doublet fermions}}} \psi_i \longleftrightarrow \text{vacuum}, \quad (1.15)$$

which leads to

$$3N_f \mu_q + \sum_j \mu_{l_j} = 0. \quad (1.16)$$

From Eqs. (1.12), (1.14) and (1.16), all the chemical potentials can be written in terms of N_f independent ones, say, $\mu_{\bar{e}_j}$. Then, from Eq. (1.10), we obtain

$$\begin{aligned} n_B &= \frac{1}{6} T^2 \times \frac{8N_f + 4N_\varphi}{2N_f + 3N_\varphi} \sum_j \mu_{\bar{e}_j}, \\ n_{L_j} &= \frac{1}{6} T^2 \times \left[\left(\frac{-8}{2N_f + 3N_\varphi} \sum_k \mu_{\bar{e}_k} \right) - 3\mu_{\bar{e}_j} \right], \\ n_L &\equiv \sum_j n_{L_j} \\ &= \frac{1}{6} T^2 \times \frac{-14N_f - 9N_\varphi}{2N_f + 3N_\varphi} \sum_j \mu_{\bar{e}_j}. \end{aligned} \quad (1.17)$$

Because the conserved quantity is the $B - L$ asymmetry in the present analysis,⁶ it is suitable to rewrite the above relations in terms of $n_B - n_L$:

$$n_B = C (n_B - n_L), \quad (1.18)$$

where C is given by [37]

$$C = \frac{8N_f + 4N_\varphi}{22N_f + 13N_\varphi}. \quad (1.19)$$

In the case of the standard model, $N_f = 3$ and $N_\varphi = 1$ yield $C = 28/79$.

We can see from Eq. (1.18) that, if $B - L$ asymmetry is absent, any baryon asymmetry vanishes at equilibrium in the presence of the sphaleron effect. (See the arrow (i) in Fig. 1.2.) On the other hand, if there occurs a successful leptogenesis, i.e., if nonzero

⁶Actually, not only the total $B - L$ but each $(1/N_f)B - L_j$ ($j = 1 \cdots N_f$) is conserved in the present situation.




Figure 1.2: The relation between baryon (B) and lepton (L) number. Thin dotted lines correspond to constant $B - L$, along which B and L can move via the sphaleron process. At equilibrium, B and L reach the thick dashed line, which represents $B = C(B - L)$.

lepton asymmetry (and hence $B - L$ asymmetry) is generated in an out-of-equilibrium way, it is partially converted into baryon asymmetry [9]. (See the arrow (ii) in Fig. 1.2.) The amount of the baryon asymmetry at equilibrium is obtained from Eq. (1.18) as

$$\left. \frac{n_B}{s} \right|_{\text{eq}} = C \left. \frac{n_B - n_L}{s} \right|_{\text{eq}} = -C \left. \frac{n_L}{s} \right|_{\text{initial}}, \quad (1.20)$$

where we have normalized the number densities by the entropy density, so that they become constant against the expansion of the universe.

In the case of the MSSM, we have an additional Higgs doublet as well as supersymmetric partners. Let us calculate the coefficient C in the presence of those particles. First, the (gaugino)-(fermion)-(sfermion)* interactions ensure that the chemical potential of each sfermion⁷ is the same as that of the corresponding fermion. (Note that the chemical potentials of the gauginos vanish since they are Majorana particles.) Next, the supersymmetric mass term $W = \mu H_u H_d$ makes the chemical potentials of the up-type and

⁷”sfermion” denotes a scalar partner of the fermion.

down-type Higgsinos have opposite signs, and hence the relation induced by the sphaleron effect, Eq. (1.16), does not change. However, the condition of the vanishing $U(1)_Y$ charge [Eq. (1.14)] changes. (Note that the contributions from bosons and fermions are different. See Eq. (1.9).) Consequently, the coefficient C becomes again $C = 28/79$, in spite of the presence of two Higgs doublets. In the actual thermal history, however, the Higgsino and sfermions likely become massive and are essentially absent at the time of electroweak phase transition. Hence, it is more appropriate to calculate C without them, but with two Higgs doublets. This ($N_f = 3$, $N_\varphi = 2$) leads to $C = 8/23$.

So far, we have considered the symmetric phase $T > T_C$. The actual ratio of the baryon to lepton asymmetry in the present universe depends on how the electroweak phase transition occurs [38]. More precisely, it depends on the value of $v(T)/T$ just before the sphaleron decoupling $T \simeq T_*$. Fortunately, however, numerically it does not change much from Eq. (1.19). (The coefficient C changes from $8/23$ ($v \ll T$) to $10/31$ ($v \gg T$) for $N_f = 3$ and $N_\varphi = 2$ [38], so that the difference is at most a few percent.) Thus, we will take $C = 8/23 \simeq 0.35$ throughout this thesis, for simplicity.

Finally, we should also note the sign of the baryon asymmetry. We know that the sign of the present baryon asymmetry is positive, i.e., $n_B/s > 0$.⁸ Thus, from Eq. (1.20), it is found that the leptogenesis must generate a lepton asymmetry with a negative sign, $n_L/s < 0$ (i.e., more anti-leptons than leptons should be produced). In principle, the sign of the generated lepton asymmetry could be determined if we know the sign of the effective CP -violating phase in each leptogenesis mechanism. In the case of leptogenesis by the decay of right-handed neutrino (discussed in Chapter 2), it depends on the phases of the Yukawa couplings of the right-handed neutrino [see Eq. (2.14)]. On the other hand, in the case of leptogenesis via LH_u flat direction (discussed in Chapter 3), the effective CP -violating phase is determined by the relative phase between the phase of the SUSY-breaking term and the initial phase of the flat direction field ϕ , which depends on the coupling of ϕ to the inflaton field.

In both of those cases, however, it is highly difficult to determine the sign of the phases. Thus, we will simply assume that the leptogenesis mechanisms we will discuss produce the lepton asymmetry with a correct sign. Keeping in mind the discussion above, we will omit the relative sign in Eq. (1.20) for simplicity, and use the following relation throughout this thesis:

$$\frac{n_B}{s} = 0.35 \times \frac{n_L}{s} \Big|_{\text{initial}}. \quad (1.21)$$

⁸This is not a matter of definition, since we can distinguish the matter from antimatter by CP -violating processes in the laboratory experiments (e.g., the asymmetry in the decay $K_L^0 \rightarrow \pi^\pm e^\mp \nu$), which is independent of the definition of “matter” we would name from the cosmological baryon asymmetry.

1.5 Cosmological gravitino problems

In this section, we briefly review the results of the cosmological gravitino problems obtained in the literature, since they give a very important and severe constraint on the baryogenesis, i.e., the upper bound on the reheating temperature. For a review, see Ref. [39].

• unstable gravitino

There are two cases; unstable and stable gravitino. (See Fig. 1.3.) Let us first consider the unstable gravitino. Since the couplings of the gravitino with ordinary matter are strongly suppressed by the gravitational scale, it has a very long lifetime:

$$\tau_{3/2} \simeq 4 \times 10^5 \times \left(\frac{m_{3/2}}{1 \text{ TeV}} \right)^{-3} \text{ sec}, \quad (1.22)$$

where $m_{3/2}$ denotes the gravitino mass, and we have assumed that the gravitino dominantly decays into a photon and a photino⁹ and omitted the phase space suppression of the decay rate, for simplicity. Therefore, it decays after the big-bang nucleosynthesis (BBN) epoch ($t \sim 1\text{--}100$ sec), unless gravitino is heavier than ~ 10 TeV [40]. Then the energetic photon (or some charged particle) produced in the gravitino decay induce electromagnetic cascade process. This cascade might destroy the light elements and change their abundances, and spoil the success of the BBN [19]. Since the abundance of the gravitinos produced from thermal environment at reheating epoch is roughly proportional to the reheating temperature T_R , usually there are upper bounds on the T_R depending on the gravitino mass. Recently, a detailed calculation of the gravitino production rate in SUSY QCD at high temperature has been done [41].¹⁰ A recent analysis of the effect of the radiative decays of massive particles on the BBN is found in Ref. [42]. By using their results, the upper bounds are given by $T_R \lesssim 10^6$, 10^9 , and 10^{12} GeV for $m_{3/2} = 100$ GeV, 1 TeV and 3 TeV, respectively.¹¹

If the gravitino is even heavier, as in the anomaly mediated SUSY breaking models [43], it decays during or near the BBN epoch ($t \sim 10^{-2}\text{--}10^2$ sec). In this case, there is another constraint which comes from hadronic decay [21]. If energetic hadrons are emitted at this epoch, they interconvert the neutrons (n) and protons (p) in the background even

⁹If the gravitino mainly decays into a neutrino and a sneutrino, the upper bound on the reheating temperature becomes higher. See discussion in remarks below.

¹⁰Their calculation [41] shows a slightly smaller abundance of the produced gravitinos than in earlier works.

¹¹Here, we have taken the gluino mass to be $m_{\tilde{g}} = 1$ TeV. Precisely speaking, the abundance of the gravitinos produced by thermal scatterings depends on the gluino mass as $\propto \{1 + m_{\tilde{g}}^2(T)/(3m_{3/2}^2)\}$ [41].




Figure 1.3: A schematic figure of the thermal history of the universe relevant to the cosmological gravitino problems.

after the freeze-out time of the n/p ratio ($t \sim 1 \text{ sec}$), which results in the change of the abundances of the light elements. Thus, there are upper bounds on the reheating temperature T_R times the branching ratio of the gravitino decay into hadrons B_h . By using the results of Ref. [41] and a recent analysis of the effects of the hadronic decay on the BBN in Ref. [44], the upper bounds on the reheating temperatures are given by $(B_h/0.1) \times T_R \lesssim 10^9 - 10^{11} \text{ GeV}$ for $m_{3/2} \simeq (\text{a few} - 100) \text{ TeV}$.¹²

• stable gravitino

Now let us turn to discuss the case of stable gravitino, which is the case if the gravitino is the lightest SUSY particle (LSP). (We assume the R -parity and hence the LSP is stable. A gravitino much lighter than the weak scale is realized in a low-energy SUSY breaking scenario such as gauge-mediated SUSY breaking models [45].¹³) In this case,

¹²We have taken the gluino mass to be $m_{\tilde{g}} = 1 \text{ TeV}$. See footnote 11.

¹³This is not the case if the SUSY breaking is mediated by a bulk gauge field in higher dimension spacetime [46].

the upper bound on the reheating temperature T_R comes from the requirement that the energy density of the gravitino not overclose¹⁴ the present universe [20]. From the result in Ref. [41], the relic abundance of the gravitinos which are produced by scattering processes of particles from the thermal bath after the inflation is given by¹⁵

$$\Omega_{3/2} h^2 \simeq 0.3 \times \left(\frac{m_{\tilde{g}}}{1 \text{ TeV}} \right)^2 \left(\frac{m_{3/2}}{10 \text{ MeV}} \right)^{-1} \left(\frac{T_R}{10^6 \text{ GeV}} \right). \quad (1.23)$$

Here, $m_{\tilde{g}}$ is the gluino mass, h is the present Hubble parameter in units of $100 \text{ km sec}^{-1} \text{ Mpc}^{-1}$ and $\Omega_{3/2} = \rho_{3/2}/\rho_c$. ($\rho_{3/2}$ and ρ_c are the present energy density of the gravitino and the critical energy density of the present universe, respectively.) It is found from Eq. (1.23) that the overclosure limit $\Omega_{3/2} < 1$ puts a severe upper bound on the reheating temperature T_R , depending on the gravitino mass $m_{3/2}$. Here, we have omitted the contribution from the decays of squarks and sleptons into gravitinos. This effect makes the upper bound on the reheating temperature slightly severer for a smaller gravitino mass region, $m_{3/2} \sim (1\text{--}100) \text{ keV}$ [20]. For a even lighter gravitino $m_{3/2} \lesssim 1 \text{ keV}$, there is no cosmological gravitino problem [47], since in this case the gravitino does not overclose the energy density of the universe even if it is thermalized.

Meanwhile, we should also take care of the decays of the next-to-lightest SUSY particle (NLSP) [20]. Since the NLSP can decay only to the LSP gravitino through the suppressed interaction, it has a long lifetime. Then its decay during or after the BBN might destroy the success of the BBN, just like the case of unstable gravitino. Thus, there are constraints on the abundance of the NLSP at the decay time depending on the lifetime of the NLSP. (The lifetime of the NLSP depends on the NLSP mass as well as the gravitino mass, and its abundance at the time of the decay is determined by its annihilation cross section.) This constraint leads to an upper bound on the gravitino mass $m_{3/2}$ [20], depending on the mass and couplings of the NLSP. Then combined with the constraint from the overclosure limit of the gravitino explained above, we obtain an upper bound on the reheating temperature T_R . Detailed analyses show that reheating temperature can be as high as $T_R \sim 10^9\text{--}10^{10} \text{ GeV}$ for $m_{3/2} \sim 10\text{--}100 \text{ GeV}$ [48].

¹⁴The word “overclose” might not be appropriate, since the open or flat universe ($\Omega \leq 1$) cannot change to a closed ($\Omega > 1$) universe. The actual problem is that the Hubble expansion would be too high compared with the observation at a temperature of $T = T_0 = 2.73K$, if the calculated value of Ω exceeds unity [28].

¹⁵Besides the factor in Eq. (1.23), the abundance of the gravitino depends on the reheating temperature also through the $SU(3)_C$ running coupling, $\alpha_3(T_R)$. Precisely speaking, the final abundance is proportional to $\alpha_3(T_R)^3 \times \{\text{a logarithmic correction (known)}\}$ for $m_{\tilde{g}} \gg m_{3/2}$ [41].

- **remarks**

Several comments are in order. First, all of the above arguments have assumed that no extra entropy production takes place after gravitinos are produced. If there is a dilution of the gravitino by a late-time entropy production, the bounds discussed in this section are relaxed. Care has to be taken in this case, however, since baryon asymmetry is also diluted if the baryogenesis occurs before that entropy production.

Next, in the case of unstable gravitino, if the gravitino decays mainly into a neutrino and a sneutrino, the upper bound on the reheating temperature becomes weaker since the neutrino have only weak interactions. In this case, the high energy neutrinos emitted from the gravitino decay scatter off the background neutrinos and produce charged leptons, which cause electroweak cascade and produce many photons. The requirement that those photons do not alter the abundances of light elements gives an upper bound on the reheating temperature, $T_R \lesssim 10^{10}\text{--}10^{12}$ GeV for $m_{3/2} \simeq 100$ GeV–1 TeV [49].

In some cases, considerably higher reheating temperatures are allowed. For example, if the LSP is the axino (which is a fermionic superpartner of the axion) and the gravitino is the NLSP, the reheating temperature can be as high as 10^{15} GeV for $m_{3/2} \simeq 100$ GeV [50]. Another interesting case is given when the leptogenesis takes place from the universe dominated by the coherent oscillation of the right-handed sneutrino [26] (see Sec. 2.4). See also a model in Sec. 3.3.2. In the context of gauge-mediated SUSY breaking models, an attractive scenario to solve the gravitino problem has been proposed [51], recently.

Finally, we mention the nonthermal production of the gravitino during the preheating epoch after the inflation [52, 53]. At the preheating epoch, gravitinos can be produced either through the scattering of particles which are created by the parametric resonance of the oscillating inflaton [52] or directly from the oscillating inflaton [53]. These nonthermal productions may increase the primordial abundance of the gravitino and hence might make the upper bound on the reheating temperature T_R severer than discussed in this section. However, in both cases the produced gravitino abundance depends on the amplitude and coupling of the oscillating inflaton. In particular, as for the second mechanism, it was shown that the gravitino abundance produced directly from oscillating inflaton is sufficiently small as long as the two sectors, the one responsible for supersymmetry breaking at true vacuum and the one for the inflation, are distinct and coupled only gravitationally [54]. (Notice that this is the case for the SUSY inflation models which will be discussed in Sec. 2.3.) We will use, therefore, the constraint from the thermally produced gravitinos as a conservative bound.

Chapter 2

Leptogenesis by the decay of right-handed neutrino

In this chapter, we discuss leptogenesis scenarios by the decays of right-handed neutrino. The lepton asymmetry is produced by the CP -violating decay of the right-handed neutrino N into leptons L and anti-leptons $\bar{L}$. As discussed in Sec. 1.4, the produced lepton asymmetry is partially converted to the baryon asymmetry [9] by the sphaleron process [6].

In Sec. 2.1 we discuss the amount of the lepton asymmetry produced in the decays of right-handed (s)neutrino. Then we turn to discuss each scenario depending on the production mechanism of the right-handed (s)neutrinos. The original, and the most extensively studied mechanism is the thermal production. We briefly discuss it in Sec. 2.2. Next, we investigate the production of right-handed (s)neutrinos in inflaton decay in Sec. 2.3, adopting several SUSY inflation models. Here, we also study in detail the inflation dynamics in each inflation model. In Sec. 2.4, we discuss the leptogenesis from coherent oscillation of the right-handed sneutrino. In particular, we mainly discuss the most interesting case, the leptogenesis from the universe dominated by the coherent oscillation of the right-handed sneutrino.

2.1 Asymmetric decay of the right-handed neutrino

Let us start by introducing three generations of heavy right-handed neutrinos to the minimal supersymmetric standard model (MSSM), which have a superpotential;

$$W = \frac{1}{2} M_i N_i N_i + h_{i\alpha} N_i L_\alpha H_u, \quad (2.1)$$

where N_i ($i = 1, 2, 3$), L_α ($\alpha = e, \mu, \tau$) and H_u denote the supermultiplets of the heavy right-handed neutrinos, lepton doublets and the Higgs doublet which couples to up-type quarks, respectively. (Here and hereafter, we omit the $SU(2)_L$ indices for simplicity.) M_i are the masses of the right-handed neutrinos. Here, we have taken a basis where the mass matrix for N_i is diagonal and real.

As can be seen in Eq. (2.1), the masses of the right-handed neutrinos violate the lepton number. This gives rise to the following two important consequences. First, the tiny neutrino masses, which are now strongly suggested by the neutrino-oscillation experiments, are explained via the seesaw mechanism [13]. From Eq. (2.1) we obtain the mass matrix for the light neutrinos by integrating out the heavy right-handed neutrinos:

$$(m_\nu)_{\alpha\beta} = - \sum_i h_{i\alpha} h_{i\beta} \frac{\langle H_u \rangle^2}{M_i}. \quad (2.2)$$

The second one is the production of the lepton asymmetry by the right-handed neutrino decay [9] which we discuss now. Because of their Majorana masses, the right-handed neutrinos have two distinct decay channels into leptons and anti-leptons. At tree level, these two kinds of decay channels have the same decay widths:

$$\begin{aligned} \frac{1}{4} \Gamma_{N_i} &= \Gamma(N_i \rightarrow \tilde{L} + \widetilde{h_u}) = \Gamma(N_i \rightarrow \tilde{L}^* + \overline{\widetilde{h_u}}) \\ &= \Gamma(N_i \rightarrow l + H_u) = \Gamma(N_i \rightarrow \bar{l} + H_u^*) = \frac{1}{16\pi} \sum_\alpha |h_{i\alpha}|^2 M_i, \\ \frac{1}{2} \Gamma_{\tilde{N}_i} &= \Gamma(\tilde{N}_i \rightarrow \tilde{L} + H_u) = \Gamma(\tilde{N}_i \rightarrow \bar{l} + \overline{\widetilde{h_u}}) = \frac{1}{8\pi} \sum_\alpha |h_{i\alpha}|^2 M_i, \\ \frac{1}{2} \Gamma_{\tilde{N}_i^*} &= \Gamma(\tilde{N}_i^* \rightarrow l + \widetilde{h_u}) = \Gamma(\tilde{N}_i^* \rightarrow \tilde{L}^* + H_u^*) = \frac{1}{8\pi} \sum_\alpha |h_{i\alpha}|^2 M_i, \end{aligned} \quad (2.3)$$

where $N_i = \nu_{Ri}^c + \overline{\nu_{Ri}^c}^T$, $\{\nu_{Ri}^c, l, \widetilde{h_u}\}$ and $\{\tilde{N}, \tilde{L}, H_u\}$ denote fermionic and scalar components of corresponding supermultiplets $\{N, L, H_u\}$, and $\bar{y}$ and Y^* represent antiparticles of fermion y and scalar Y , respectively. Here, we have summed the final states over flavor ($\alpha = e, \mu, \tau$) and $SU(2)_L$ indices. We can symbolically write the above widths as

$$\frac{1}{2} \Gamma_{N_i} = \Gamma(N_i \rightarrow L + H_u) = \Gamma(N_i \rightarrow \bar{L} + \overline{H_u}) = \frac{1}{8\pi} \sum_\alpha |h_{i\alpha}|^2 M_i, \quad (2.4)$$

where N , L and H_u ($\bar{L}$ and $\bar{H}_u$) denote fermionic or scalar components of corresponding supermultiplets (and their anti-particles).

If CP is violated in the Yukawa matrix $h_{i\alpha}$, the interference between decay amplitudes of tree and one-loop diagrams results in lepton-number violation [9]. Hereafter, we concentrate on the decay of lightest right-handed (s)neutrino N_1 , since we will consider only the N_1 decay in the following sections.¹ The lepton asymmetry produced in the N_1 decay is represented by the following parameter ϵ_1 :

$$\epsilon_1 \equiv \frac{\Gamma(N_1 \rightarrow L + H_u) - \Gamma(N_1 \rightarrow \bar{L} + \bar{H}_u)}{\Gamma_{N_1}}, \quad (2.5)$$

which means the lepton number asymmetry produced per one right-handed neutrino decay. Summing up the one-loop vertex and self-energy corrections [55], the ϵ_1 has the following form:

$$\epsilon_1 = -\frac{1}{8\pi} \frac{1}{(hh^\dagger)_{11}} \sum_{i=2,3} \text{Im} [\{(hh^\dagger)_{1i}\}^2] \left[f^V \left(\frac{M_i^2}{M_1^2} \right) + f^S \left(\frac{M_i^2}{M_1^2} \right) \right], \quad (2.6)$$

where $f^V(x)$ and $f^S(x)$ represent the contributions from vertex and self-energy corrections, respectively. In the case of the non-supersymmetric standard model with right-handed neutrinos, they are given by [55]

$$f_{\text{non-SUSY}}^V(x) = \sqrt{x} \left[-1 + (x+1) \ln \left(1 + \frac{1}{x} \right) \right], \quad f_{\text{non-SUSY}}^S(x) = \frac{\sqrt{x}}{x-1}, \quad (2.7)$$

while in the case of MSSM plus right-handed (s)neutrinos, they are given by [56],²

$$f_{\text{SUSY}}^V(x) = \sqrt{x} \ln \left(1 + \frac{1}{x} \right), \quad f_{\text{SUSY}}^S(x) = \frac{2\sqrt{x}}{x-1}. \quad (2.8)$$

Hereafter, we use Eq. (2.8) since we assume SUSY. Notice that all of the N_1 , $\tilde{N}_1$ and $\tilde{N}_1^*$ in Eq. (2.3) produce the same amount of lepton asymmetry with the same sign [57, 56, 58],³ given by the above ϵ_1 . Assuming a mass hierarchy $M_1 \ll M_2, M_3$ in the right-handed neutrino sector (i.e., $x \gg 1$), the above formula is simplified to the following one:

$$\epsilon_1 \simeq -\frac{3}{8\pi} \frac{1}{(hh^\dagger)_{11}} \sum_{i=2,3} \text{Im} [\{(hh^\dagger)_{1i}\}^2] \frac{M_1}{M_i}. \quad (2.9)$$

¹We will assume $M_1 \ll M_2, M_3$. In this case, even if the heavier right-handed neutrinos $N_{2,3}$ produce lepton asymmetry, it is usually erased before the decays of N_1 .

²See also Refs. [57, 24].

³Here, we have neglected the three body decay of the right-handed sneutrino, e.g., $\tilde{N}_1 \rightarrow \tilde{L}^* + \tilde{Q} + \tilde{u}$, which gives only tiny corrections [58]. We also neglect the effects of the soft-SUSY breaking terms, since the mass scale of soft terms ($\sim 100 \text{ GeV} - 1 \text{ TeV}$) are much smaller than right-handed neutrino mass.

Now let us rewrite this ϵ_1 parameter in terms of the light neutrino mass m_ν , so that we can relate the lepton asymmetry (and hence the baryon asymmetry in the present universe) to the neutrino mass, which is observed by the neutrino-oscillation experiments. First of all, by using $\text{Im} \left[\left(hh^\dagger \right)_{11} \right]^2 = 0$, we can write the ϵ_1 as follows:

$$\epsilon_1 = -\frac{3}{8\pi} \frac{M_1}{(hh^\dagger)_{11}} \text{Im} \left[\left(hh^\dagger \frac{1}{M} h^* h^T \right)_{11} \right], \quad (2.10)$$

where a matrix notation $M = \text{diag}(M_1, M_2, M_3)$ is adopted. Next, by using the seesaw formula in Eq. (2.2), it is reduced to

$$\epsilon_1 = \frac{3}{8\pi} \frac{M_1}{\langle H_u \rangle^2} \frac{\text{Im} \left[h(m_\nu^*) h^T \right]_{11}}{(hh^\dagger)_{11}}. \quad (2.11)$$

The neutrino mass matrix $(m_\nu)_{\alpha\beta}$ can be diagonalized by an unitary matrix $U_{\alpha i}$ (Maki-Nakagawa-Sakata matrix [59]) as $(m_\nu)_{\alpha\beta} = \sum_i U_{\alpha i} U_{\beta i} (\widehat{m}_\nu)_i$, where $\widehat{m}_\nu$ is a diagonal mass matrix $\widehat{m}_\nu = \text{diag}(m_{\nu 1}, m_{\nu 2}, m_{\nu 3})$ and $m_{\nu 1} < m_{\nu 2} < m_{\nu 3}$. Then, with rotated Yukawa couplings

$$\widehat{h}_{ik} \equiv \sum_\alpha h_{i\alpha} U_{\alpha k}^* \quad (2.12)$$

we obtain

$$\begin{aligned} \epsilon_1 &= \frac{3}{8\pi} \frac{M_1}{\langle H_u \rangle^2} \frac{\text{Im} \left[\widehat{h}(\widehat{m}_\nu) \widehat{h}^T \right]_{11}}{(\widehat{h} \widehat{h}^\dagger)_{11}} \\ &= \frac{3}{8\pi} \frac{M_1}{\langle H_u \rangle^2} m_{\nu 3} \delta_{\text{eff}}, \end{aligned} \quad (2.13)$$

where the effective CP -violating phase δ_{eff} is defined by

$$\delta_{\text{eff}} \equiv \frac{\text{Im} \left[(\widehat{h}_{13})^2 + \frac{m_{\nu 2}}{m_{\nu 3}} (\widehat{h}_{12})^2 + \frac{m_{\nu 1}}{m_{\nu 3}} (\widehat{h}_{11})^2 \right]}{|\widehat{h}_{13}|^2 + |\widehat{h}_{12}|^2 + |\widehat{h}_{11}|^2}. \quad (2.14)$$

As clearly seen from the above explicit expression, δ_{eff} is always less than one [26], but it is in general order one unless the phase of the Yukawa coupling $\widehat{h}_{13}$ is accidentally suppressed or the couplings have a inverted hierarchy, i.e., $|\widehat{h}_{12}|^2 \gg (m_{\nu 2}/m_{\nu 1})|\widehat{h}_{13}|^2$ or $|\widehat{h}_{11}|^2 \gg (m_{\nu 3}/m_{\nu 1})|\widehat{h}_{13}|^2$. From Eq. (2.13), the ϵ_1 parameter is given by

$$\epsilon_1 \simeq 2.0 \times 10^{-10} \left(\frac{M_1}{10^6 \text{ GeV}} \right) \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}. \quad (2.15)$$

This relation is consistent with the one obtained in Ref. [60]. Here, we have used $\langle H_u \rangle = 174 \text{ GeV} \times \sin \beta$, where $\tan \beta \equiv \langle H_u \rangle / \langle H_d \rangle$. (H_d is the Higgs field which couples to

down-type quarks.) Here and hereafter, we will take $\sin\beta \simeq 1$ for simplicity.⁴ As for the heaviest neutrino mass, we take $m_{\nu 3} \simeq 0.05$ eV as a typical value throughout this chapter, suggested from the atmospheric neutrino oscillation observed in the Super-Kamiokande experiments [10].

We should stress here that the ϵ_1 parameter has an explicit formula given in Eq. (2.15) with δ_{eff} in Eq. (2.14) (as long as $M_1 \ll M_2, M_3$), although in the literature it is sometimes treated as a free parameter. In particular, it is proportional to the right-handed neutrino mass M_1 for fixed values of $m_{\nu 3}$ and $\delta_{\text{eff}} (\leq 1)$.

Let us mention one last point, the possibility of an enhancement of the asymmetry parameter ϵ_1 . For the self-energy contribution $f^S(x)$ in Eq. (2.8) (and Eq. (2.7)), we have assumed the masses of the right-handed neutrinos are not so degenerate, i.e., $|M_i - M_1| \gg \Gamma_{N_i}$. However, if the mass difference becomes as small as the decay width, $|M_i - M_1| \sim \Gamma_{N_i}$, one expects an enhancement of the self-energy contribution $f^S(x)$ [61]. Actually, it was shown that the asymmetry parameter ϵ_1 can reach its maximum value of $\epsilon_1 \sim \mathcal{O}(1)$ for $|M_i - M_1| \simeq \Gamma_{N_i}/2$ [61]. (ϵ_1 cannot be arbitrarily large. Notice that the lepton asymmetry vanishes in the limit where the right-handed neutrinos become exactly mass degenerate, since in this case CP -violating phases of the Yukawa couplings can be eliminated by a change of basis.) Nevertheless, it requires an extreme degeneracy of right-handed neutrino masses $|M_i - M_1|/M_1 \simeq h^2/(8\pi)$, and hence we do not consider this possibility in the following discussion.

⁴This is the case as long as $\tan\beta > 1$. Even for $\tan\beta \simeq 1$, the final lepton asymmetry changes (increases) by a factor of 2.

2.2 Leptogenesis by thermally produced right-handed neutrino

In this section, we briefly discuss the leptogenesis by thermally produced right-handed neutrinos. After first suggested by Fukugita and Yanagida [9], this scenario has been extensively studied. (See, for example, Refs. [62, 57, 55, 61, 63]. For reviews and references, see Refs. [22, 63].) From the viewpoint of the production mechanism of the right-handed neutrino, compared with other production mechanisms, this scenario has a very attractive point that it requires no extra assumption to create the right-handed neutrinos, besides high enough temperature.

Suppose that the right-handed (s)neutrinos N_1 are produced thermally and become as abundant as in thermal equilibrium at temperature $T > M_1$. (Hereafter, we consider the lightest right-handed neutrino N_1 , since lepton asymmetry produced by the decays of the heavier two right-handed neutrinos $N_{2(3)}$ is likely to be erased before the N_1 's decay.) Then the ratio of the total number density of the right-handed (s)neutrinos to the entropy density is simply given by the following number [see Eq. (B.19)]:⁵

$$\frac{n_{N_1}}{s} \simeq \frac{n_{N_1}^{\text{eq}}}{s} = \frac{45\zeta(3)}{2\pi^4} \frac{1}{g_*} \times \left(\frac{3}{4}g_{N_1} + g_{\widetilde{N}_1} \right) \simeq \frac{1}{240}, \quad (2.16)$$

where $g_{N_1} = g_{\widetilde{N}_1} = 2$ are the numbers of degrees of freedom for right-handed (s)neutrinos and $g_* = 232.5$ for the MSSM plus right-handed neutrino multiplet [see Eq. (B.15)].

Boltzmann equations (a toy model) The evolution of the number density n_{N_1} of the right-handed neutrino and that of the lepton number density n_L are described by coupled Boltzmann equations. In order to understand qualitative behaviors of the n_{N_1} and n_L , in particular to understand the “out-of-equilibrium condition,” let us consider the following simple set of Boltzmann equations:

$$\begin{aligned} \frac{d}{dt}n_{N_1} + 3Hn_{N_1} &= -\gamma(N_1 \rightarrow l\varphi) - \gamma(N_1 \rightarrow \bar{l}\varphi^*) \\ &\quad + \gamma(l\varphi \rightarrow N_1) + \gamma(\bar{l}\varphi^* \rightarrow N_1), \end{aligned} \quad (2.17)$$

$$\begin{aligned} \frac{d}{dt}n_l + 3Hn_l &= \gamma(N_1 \rightarrow l\varphi) - \gamma(l\varphi \rightarrow N_1) \\ &\quad - \gamma'(l\varphi \rightarrow \bar{l}\varphi^*) + \gamma'(\bar{l}\varphi^* \rightarrow l\varphi), \end{aligned} \quad (2.18)$$

$$\begin{aligned} \frac{d}{dt}n_{\bar{l}} + 3Hn_{\bar{l}} &= \gamma(N_1 \rightarrow \bar{l}\varphi^*) - \gamma(\bar{l}\varphi^* \rightarrow N_1) \\ &\quad + \gamma'(l\varphi \rightarrow \bar{l}\varphi^*) - \gamma'(\bar{l}\varphi^* \rightarrow l\varphi), \end{aligned} \quad (2.19)$$

⁵Numerically the value in Eq. (2.16) is almost the same as $1/g_*$.

which describe the evolutions of the number densities of right-handed neutrino N_1 , lepton l , and anti-lepton $\bar{l}$. For a while, we will consider a non-SUSY case, for simplicity. The terms proportional to the Hubble parameter H describe the effect of the expansion of the universe. $\gamma(A \rightarrow B)$ denotes the reaction density, which means the rate of the process $A \rightarrow B$ per unit time per unit volume. Thus, the γ -terms in the right-hand sides of the equations describe the change of the number densities due to the corresponding interactions. We have included only the three kinds of processes ($N_1 \leftrightarrow l\varphi$, $N_1 \leftrightarrow \bar{l}\varphi^*$ and $l\varphi \leftrightarrow \bar{l}\varphi^*$) which are necessary in the following discussion. (In actual calculation, one must include many other interactions. For other processes, which include the superpartners $\widetilde{N}_1$, $\widetilde{L}$ and $\widetilde{h}_u$, see Ref. [58].) As for the primes in $\gamma'(l\varphi \leftrightarrow \bar{l}\varphi^*)$, we will give an explanation below.

The reaction densities of the decay and inverse decay are given by

$$\begin{aligned}\gamma(N_1 \rightarrow l\varphi) &= \frac{1}{2}(1 + \epsilon_1) \langle \Gamma_{N_1} \rangle n_{N_1}, \\ \gamma(N_1 \rightarrow \bar{l}\varphi^*) &= \frac{1}{2}(1 - \epsilon_1) \langle \Gamma_{N_1} \rangle n_{N_1}, \\ \gamma(l\varphi \rightarrow N_1) &= \frac{1}{2}(1 - \epsilon_1) \langle \Gamma_{N_1} \rangle n_{N_1}^{\text{eq}} \times \frac{n_l}{n_l^{\text{eq}}}, \\ \gamma(\bar{l}\varphi^* \rightarrow N_1) &= \frac{1}{2}(1 + \epsilon_1) \langle \Gamma_{N_1} \rangle n_{N_1}^{\text{eq}} \times \frac{n_{\bar{l}}}{n_{\bar{l}}^{\text{eq}}}.\end{aligned}\tag{2.20}$$

Here, $\langle \Gamma_{N_1} \rangle$ denotes thermally averaged total decay rate of N_1 . For low temperature $T \ll M_1$, it is just given by the decay rate at rest frame: $\langle \Gamma_{N_1} \rangle \simeq \Gamma_{N_1}$, while for high temperature $T \gg M_1$, it is given by $\langle \Gamma_{N_1} \rangle \simeq (M_1/2T) \Gamma_{N_1}$ due to the time-dilatation effect.

From Eqs. (2.17)–(2.20), the Boltzmann equations of $Y_{N_1} \equiv n_{N_1}/s$ and $Y_L \equiv n_L/s = (n_l - n_{\bar{l}})/s$ are given by

$$\frac{d}{dt}Y_{N_1} = -\langle \Gamma_{N_1} \rangle (Y_{N_1} - Y_{N_1}^{\text{eq}}) - \frac{1}{2}\epsilon_1 \langle \Gamma_{N_1} \rangle Y_{N_1}^{\text{eq}} \times \frac{Y_L}{Y_l^{\text{eq}}},\tag{2.21}$$

$$\begin{aligned}\frac{d}{dt}Y_L &= \epsilon_1 \langle \Gamma_{N_1} \rangle (Y_{N_1} + Y_{N_1}^{\text{eq}}) - \frac{1}{2} \langle \Gamma_{N_1} \rangle Y_{N_1}^{\text{eq}} \times \frac{Y_L}{Y_l^{\text{eq}}} \\ &\quad - 2\gamma'(l\varphi \rightarrow \bar{l}\varphi^*) + 2\gamma'(\bar{l}\varphi^* \rightarrow l\varphi).\end{aligned}\tag{2.22}$$

Here, there is a subtle point one should take care [64, 65]. As can be seen from Eq. (2.20), the inverse decay processes ($l\varphi \rightarrow N_1$ and $\bar{l}\varphi^* \rightarrow N_1$) produce a net lepton asymmetry with the *same* sign as the decay processes ($N_1 \rightarrow l\varphi$ and $N_1 \rightarrow \bar{l}\varphi^*$) themselves. This can be seen by applying the CPT invariance to the matrix elements of those processes: $\mathcal{M}(N_1 \rightarrow l\varphi) = \mathcal{M}(\bar{l}\varphi^* \rightarrow N_1)$ and $\mathcal{M}(N_1 \rightarrow \bar{l}\varphi^*) = \mathcal{M}(l\varphi \rightarrow N_1)$. Therefore, if there would exist only those decay and inverse decay processes, lepton asymmetry would not

vanish even in thermal equilibrium. In other words, if we ignore the $\gamma'(l\varphi \leftrightarrow \bar{l}\varphi^*)$ terms, the right-hand side of the Eq. (2.22) would not vanish even for $Y_L = 0$ and $Y_{N_1} = Y_{N_1}^{\text{eq}}$.

On the other hand, there are lepton number violating two-body scattering processes; $l\varphi \rightarrow \bar{l}\varphi^*$ and $\bar{l}\varphi^* \rightarrow l\varphi$. By using unitarity, one can show that these processes do not produce a net lepton asymmetry if we consider the N_1 only as a virtual particle [65, 66]. However, the Boltzmann equations already include N_1 as a real particle. Thus one should subtract from the above two body scatterings the resonant s -channel contribution mediated by N_1 (which is understood as an on-shell real particle) to avoid a double counting of reactions. After subtracting the resonant contribution, the reaction densities of the two-body scatterings in Eq. (2.22) leads to [65, 66]

$$\begin{aligned} & -2\gamma'(l\varphi \rightarrow \bar{l}\varphi^*) + 2\gamma'(\bar{l}\varphi^* \rightarrow l\varphi) \\ = & -2\epsilon_1 \langle \Gamma_{N_1} \rangle n_{N_1}^{\text{eq}} - n_L n_\varphi^{\text{eq}} \left[\langle v\sigma'(l\varphi \rightarrow \bar{l}\varphi^*) \rangle + \langle v\sigma'(\bar{l}\varphi^* \rightarrow l\varphi) \rangle \right], \end{aligned} \quad (2.23)$$

where the primes mean that the contribution from resonant s -channel N_1 exchange has been subtracted. (σ is the cross section, v is the relative velocity between the initial particles and the bracket denotes the thermal average.) Notice that there appears a term $-2\epsilon_1 \langle \Gamma_{N_1} \rangle n_{N_1}^{\text{eq}}$, which originates in the subtraction of the resonant contribution [65]. After all, the Boltzmann equations are reduced to the following forms.

$$\frac{d}{dt}Y_{N_1} = -\langle \Gamma_{N_1} \rangle (Y_{N_1} - Y_{N_1}^{\text{eq}}) - \frac{1}{2} \left(\epsilon_1 \frac{Y_L}{Y_l^{\text{eq}}} \right) \langle \Gamma_{N_1} \rangle Y_{N_1}^{\text{eq}}, \quad (2.24)$$

$$\begin{aligned} \frac{d}{dt}Y_L = & \epsilon_1 \langle \Gamma_{N_1} \rangle (Y_{N_1} - Y_{N_1}^{\text{eq}}) \\ & - \left(\frac{1}{2} \langle \Gamma_{N_1} \rangle \frac{Y_{N_1}^{\text{eq}}}{Y_l^{\text{eq}}} + n_\varphi^{\text{eq}} \left[\langle v\sigma'(l\varphi \rightarrow \bar{l}\varphi^*) \rangle + \langle v\sigma'(\bar{l}\varphi^* \rightarrow l\varphi) \rangle \right] \right) Y_L. \end{aligned} \quad (2.25)$$

We see that, for $Y_{N_1} = Y_{N_1}^{\text{eq}}$ (and $Y_L = 0$), the lepton asymmetry produced by [decay and inverse decay] processes is canceled out by the lepton asymmetry from [two body scatterings minus resonance] processes, which ensures vanishing lepton asymmetry at equilibrium.

Now let us discuss the evolutions of Y_{N_1} and Y_L . As can be seen from Eq. (2.25), a deviation of the number density of the N_1 from its equilibrium value ($Y_{N_1} \neq Y_{N_1}^{\text{eq}}$) is mandatory in order to produce a net lepton asymmetry $Y_L \neq 0$ from $Y_L = 0$. This can be realized when the temperature of the universe T cools down and becomes below the mass of the right-handed neutrino M_1 . To see the out-of-equilibrium condition for N_1 , let us adopt a dimensionless variable $z \equiv M_1/T$ instead of the cosmic time t . Then Eq. (2.24)

becomes

$$\frac{dY_{N_1}}{dz} \simeq -\frac{\langle \Gamma_{N_1} \rangle}{zH} (Y_{N_1} - Y_{N_1}^{\text{eq}}). \quad (2.26)$$

Here, we have omitted the higher order term proportional to $\epsilon_1(Y_L/Y_l^{\text{eq}})$, for simplicity. For $T < M_1$ ($z > 1$), the equilibrium value of the N_1 's abundance $Y_{N_1}^{\text{eq}}$ decreases due to the Boltzmann suppression $\propto \exp(-M_1/T)$. At this stage, if the dimensionless prefactor $\langle \Gamma_{N_1} \rangle / zH$ is small enough, Y_{N_1} can no longer catches up the decreasing equilibrium value $Y_{N_1}^{\text{eq}}$. Namely, Y_{N_1} deviates from $Y_{N_1}^{\text{eq}}$, if $\langle \Gamma_{N_1} \rangle / zH \lesssim 1$ for $z \gtrsim 1$. Thus, the out-of-equilibrium condition is roughly given by

$$\frac{\Gamma_{N_1}}{H|_{T=M_1}} \lesssim 1. \quad (2.27)$$

If this condition is satisfied, $Y_{N_1} > Y_{N_1}^{\text{eq}}$ is realized, and a net lepton asymmetry is produced [see Eq. (2.25)].

The ratio $\Gamma_{N_1}/H(T = M_1)$ is related to a mass parameter $\widetilde{m}_1$, which is defined as [22]:

$$\begin{aligned} \widetilde{m}_1 &\equiv \sum_{\alpha} |h_{1\alpha}|^2 \frac{\langle H_u \rangle^2}{M_1} \\ &\simeq 0.8 \times 10^{-3} \text{ eV} \times \left(\frac{\Gamma_{N_1}}{H|_{T=M_1}} \right). \end{aligned} \quad (2.28)$$

Notice that the above formula looks similar to the neutrino mass given in Eq. (2.2) but different from that. In terms of $\widetilde{m}_1$, the out-of-equilibrium condition in Eq. (2.27) is roughly equivalent to $\widetilde{m}_1 \lesssim 10^{-3} \text{ eV}$.

It is possible to show an important constraint on this parameter $\widetilde{m}_1$ [67]:

$$\widetilde{m}_1 > m_{\nu 1}. \quad (2.29)$$

To show this, let us rewrite the $\widetilde{m}_1$ parameter in terms of the rotated Yukawa couplings $\widehat{h}_{ik}$ defined in Eq. (2.12);

$$\widetilde{m}_1 = \sum_k |\widehat{h}_{1k}|^2 \frac{\langle H_u \rangle^2}{M_1}. \quad (2.30)$$

On the other hand, from the seesaw formula Eq. (2.2), we obtain

$$m_{\nu j} \delta_{jk} = - \sum_i \widehat{h}_{ij} \widehat{h}_{ik} \frac{\langle H_u \rangle^2}{M_i}. \quad (2.31)$$

Let us define here a matrix X_{ij} as follows:

$$X_{ij} \equiv i \widehat{h}_{ij} \frac{\langle H_u \rangle}{\sqrt{M_i m_{\nu j}}}. \quad (2.32)$$

Then from Eq. (2.31) one can show that

$$\left(X^T X\right)_{ij} = \delta_{ij} = \left(XX^T\right)_{ij}, \quad (2.33)$$

while Eq. (2.30) gives rise to

$$\begin{aligned} \widetilde{m}_1 &= \sum_k m_{\nu k} |X_{1k}|^2 \\ &> \min_j \{m_{\nu j}\} \sum_k |X_{1k}|^2 \\ &\geq \min_j \{m_{\nu j}\} \left| \sum_k X_{1k}^2 \right| = \min_j \{m_{\nu j}\}. \end{aligned} \quad (2.34)$$

In the last equation, we have used Eq. (2.33). Therefore, the $\widetilde{m}_1$ parameter is bounded from below as $\widetilde{m}_1 > m_{\nu 1}$ [67].

Lepton asymmetry Now let us discuss the amount of produced lepton asymmetry in the present scenario. If the decaying right-handed neutrinos are as abundant as in thermal equilibrium [see Eq. (2.16)], and if there is no wash-out process of the produced lepton asymmetry afterwards, the final lepton asymmetry would be given by a very simple formula,

$$\frac{n_L}{s} = \epsilon_1 \frac{n_{N_1}}{s} \simeq \frac{\epsilon_1}{240}, \quad (2.35)$$

with the asymmetry parameter ϵ_1 given in Eq. (2.15). In the actual case, however, a suppression factor κ should be multiplied:

$$\frac{n_L}{s} = \kappa \epsilon_1 \frac{n_{N_1}}{s} \simeq \kappa \frac{\epsilon_1}{240}. \quad (2.36)$$

The suppression factor κ represents two effects. The first one is the wash-out effect of the produced lepton asymmetry (and the strength of the deviation from thermal equilibrium). If the interactions of right-handed neutrinos N_1 are too strong, the produced lepton asymmetry would be washed out by the interactions mediated by the N_1 itself. [See the second term in Eq. (2.25).] Weakness of the interaction is also required from the out-of-equilibrium condition for N_1 in Eq. (2.27).

The second effect represented by κ is the efficiency of the production of the right-handed (s)neutrinos N_1 . Although we have assumed that N_1 are produced as abundant as in thermal equilibrium $Y_{N_1} \simeq Y_{N_1}^{\text{eq}}$ for $T > M_1$, if the N_1 's interactions are too weak, the thermal scatterings cannot produce enough amount of N_1 and hence the number density

n_{N_1} cannot become as abundant as that in thermal equilibrium, $n_{N_1}^{\text{eq}}$.⁶ Notice that those two effects have double-edged behavior, i.e., the right-handed neutrino N_1 should have couplings $h_{1\alpha}$ with intermediate strength.

In order to determine the precise value of the suppression factor κ , one has to solve the Boltzmann equations including production, decay and inverse decay of the right-handed (s)neutrinos, and all the relevant lepton-number violating (and conserving) scatterings of lepton fields [62, 22]. A detailed numerical calculation solving the coupled Boltzmann equations in the case of SUSY has been done in Ref. [58]. (See also Ref. [22] and references therein.) It was shown that both of the production rate of the right-handed (s)neutrinos and the rate of the wash-out process of the lepton asymmetry are proportional to the mass parameter $\widetilde{m}_1$ defined in Eq. (2.28). It turns out that the suppression factor κ becomes as large as $\kappa \simeq 0.05\text{--}0.3$ for [58]

$$10^{-5} \text{ eV} \lesssim \widetilde{m}_1 \lesssim 5 \times 10^{-3} \text{ eV}. \quad (2.37)$$

For a smaller value of $\widetilde{m}_1$, the N_1 is not produced enough and generated lepton asymmetry is suppressed. On the other hand, for a larger value of $\widetilde{m}_1$, enough amount of N_1 is produced but wash-out of the lepton asymmetry is too strong, and hence final lepton asymmetry is reduced.

It is quite interesting to observe that the favored range of the mass parameter $\widetilde{m}_1$ is just below the neutrino mass scale observed by the atmospheric [10] and solar [11] (+KamLAND [12]) neutrino-oscillation experiments, $m_\nu \sim \mathcal{O}(10^{-3})\text{--}\mathcal{O}(10^{-1})$ eV. We should note that although $\widetilde{m}_1$ is not directly related to the neutrino masses $m_{\nu i}$, they are still indirectly related. (For instance, if we adopt a Froggatt-Nielsen (FN) model [68] which will be discussed in Sec. 2.3.2, $\widetilde{m}_1$ is estimated as $\widetilde{m}_1 \sim \mathcal{O}(m_{\nu 3}) \sim \mathcal{O}(0.05 \text{ eV})$. Although this is a bit larger than the range in Eq. (2.37), it can be consistent when we include $\mathcal{O}(1)$ ambiguities in the FN model. See also Ref. [69], where a more general case was discussed for two generations of neutrinos.)

After being produced, a part of lepton asymmetry is immediately converted to the baryon asymmetry [9] via the sphaleron effect [6] discussed in Sec. 1.4. Then the present baryon asymmetry is given by

$$\begin{aligned} \frac{n_B}{s} &= 0.35 \times \frac{n_L}{s} \\ &\simeq 0.3 \times 10^{-10} \left(\frac{\kappa}{0.1} \right) \left(\frac{M_1}{10^9 \text{ GeV}} \right) \times \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}, \end{aligned} \quad (2.38)$$

⁶One can consider that enough amount of right-handed neutrinos exist from the beginning and discuss their decay in thermal background. We do not consider such a case in this section, however, and assume that the right-handed neutrinos are produced by thermal scatterings from $n_{N_1} \simeq 0$ [22]. Notice that as long as an inflationary epoch is assumed, right-handed neutrinos should be produced at some stage from $n_{N_1} \simeq 0$.

where we have used Eqs. (2.15) and (2.36). Therefore, the present baryon asymmetry $n_B/s \simeq (0.4\text{--}1) \times 10^{-10}$ is naturally explained with the mass of the lightest right-handed neutrino $M_1 \sim 10^9\text{--}10^{10}$ GeV, for $\kappa \simeq 0.05\text{--}0.3$ and $\delta_{\text{eff}} \simeq \mathcal{O}(1)$.

In order to produce enough amount of right-handed neutrinos (i.e., $n_{N_1} \simeq n_{N_1}^{\text{eq}}$), the temperature of the universe should be higher than their mass M_1 . This leads to a lower bound on the reheating temperature as $T_R \gtrsim \mathcal{O}(10^{10})$ GeV. Thus the overproduction of gravitinos might cause a difficulty depending on the gravitino mass, as discussed in Sec. 1.5. If the gravitino is unstable, it must be relatively heavy, $m_{3/2} \gtrsim (\text{a few})$ TeV. When the gravitino is stable, a consistent thermal history can be obtained with gravitino mass 10–100 GeV, avoiding the problem of the decays of next-to-lightest SUSY particles after the big-bang nucleosynthesis [48].⁷

Finally, we comment on the absolute upper bound on the neutrino masses from thermal leptogenesis, which has been shown recently in Ref. [63]. The crucial observation in Ref. [63] is that the suppression factor κ is determined only by three parameters $\kappa = \kappa(\widetilde{m}_1, M_1, \overline{m})$, and hence the final baryon asymmetry in thermal leptogenesis depends only on four parameters: ϵ_1 , $\widetilde{m}_1$, M_1 , and $\overline{m}$. [See Eq. (2.36).] Here, $\overline{m} \equiv \sqrt{m_{\nu 1}^2 + m_{\nu 2}^2 + m_{\nu 3}^2}$. Furthermore, the maximal value of ϵ_1 is also determined by M_1 and $\overline{m}$, $\epsilon_1 < \epsilon_1^{\text{max}}(M_1, \overline{m})$, once the mass squared differences of atmospheric and solar neutrino oscillations are given [see Eq. (2.14) and Eq. (2.15)] [26, 70, 63]. Thus, the maximal baryon asymmetry is determined only by the set of three parameters $(\widetilde{m}_1, M_1, \overline{m})$. By solving the Boltzmann equations for different points in this parameter space $(\widetilde{m}_1, M_1, \overline{m})$, and using the bound $\widetilde{m}_1 > m_{\nu 1}$, it has been shown in Ref. [63] that the maximal baryon asymmetry can be larger than the empirical value only if

$$\overline{m} < 0.30 \text{ eV}. \quad (2.39)$$

Therefore, if the baryon asymmetry in the present universe was indeed generated by thermal leptogenesis, we have stringent constraints on the absolute neutrino masses,

$$m_{\nu i} < 0.18 \text{ eV}. \quad (2.40)$$

⁷See also Ref. [66].

2.3 Leptogenesis in inflaton decay

In this section,⁸ we discuss the leptogenesis scenario where the right-handed neutrino is produced non-thermally in inflaton decays [23]. We will find that this scenario is fully consistent with existing various SUSY inflation models such as hybrid, new, and topological inflation models.⁹

The crucial difference between this scenario and the case of the thermally produced right-handed neutrino discussed in the previous section is that, the heavy right-handed neutrino N can be produced with relatively low reheating temperatures T_R of the inflation. We find that the required baryon asymmetry can be obtained even for $T_R \simeq 10^6$ GeV in some of the inflation models, and hence there is no cosmological gravitino problem in the interesting wide region of the gravitino mass $m_{3/2} \simeq 10$ MeV–10 TeV.¹⁰ (See Sec. 1.5.)

On the other hand, the amount of the produced lepton asymmetry (and hence baryon asymmetry) in the present scenario crucially depends on the physics of the inflation, such as the mass of the inflaton m_χ and the reheating temperature T_R . Therefore, detailed analyses on the inflation models are necessary.

In Sec. 2.3.1, we calculate the amount of the resultant lepton asymmetry in inflaton decay. Then we introduce a Froggatt-Nielsen model in Sec. 2.3.2 to estimate the mass (and decay rate) of the right-handed neutrino. The subsequent subsections are devoted to each SUSY inflation model. The leptogenesis in a hybrid inflationary universe is discussed in Sec. 2.3.3, where we consider two different types of SUSY hybrid inflation models. In Sec. 2.3.4 we discuss the leptogenesis in a SUSY new inflation model. The case of a SUSY topological inflation is considered in Sec. 2.3.5. Finally, we will briefly comment on the production of right-handed neutrinos at preheating [74] in Sec. 2.3.6.

2.3.1 Lepton asymmetry

Let us first estimate the produced lepton asymmetry in the present scenario. The result obtained here is a generic one, which can be applied to all the inflation models discussed in subsequent subsections.

After the end of inflation, the inflaton decays into light particles and the energy of the inflaton is transferred into the thermal bath. Then it is very plausible that the right-handed neutrino is also produced in inflaton decay, if its decay channel is kinematically

⁸This section is based on the works in a collaboration with T. Asaka, M. Kawasaki, and T. Yanagida [71].

⁹Leptogenesis in a “natural chaotic inflation model” [72] was also investigated in Ref. [73].

¹⁰Here, we consider both of the unstable and stable gravitino.

allowed. We assume that the inflaton decays into two right-handed neutrinos, which leads to the following constraint:

$$m_\chi > 2M_1, \quad (2.41)$$

where m_χ is the inflaton mass. Here and hereafter, we consider only the N_1 decay, provided that the mass M_1 is much smaller than the others ($M_1 \ll M_2, M_3$). As discussed in Sec. 2.1, the decay of N_1 into leptons and anti-leptons produces a lepton asymmetry. We will consider the case where the right-handed neutrino is heavy enough compared with the reheating temperature, i.e., $M_1 \gtrsim T_R$. In this case the produced right-handed neutrino N_1 is always out of thermal equilibrium and it behaves like frozen-out, relativistic particle with energy $E_{N_1} \simeq m_\chi/2$. The ratio of the number density of the right-handed (s)neutrinos n_{N_1} to the entropy density s is then given by [23]

$$\begin{aligned} \frac{n_{N_1}}{s} &\simeq \frac{\rho_{\text{rad}}}{s} \times \frac{n_\chi}{\rho_\chi} \times \frac{n_{N_1}}{n_\chi} \\ &= \frac{3}{4} T_R \times \frac{1}{m_\chi} \times 2B_r = \frac{3}{2} B_r \frac{T_R}{m_\chi}, \end{aligned} \quad (2.42)$$

where ρ_{rad} is the energy density of the radiation just after the reheating process completes, and n_χ and $\rho_\chi = m_\chi n_\chi \simeq \rho_{\text{rad}}$ are the number and energy density of the inflaton just before its decay, respectively. $B_r = B_r(\chi \rightarrow N_1 N_1)$ denotes the branching ratio of the inflaton decay into N_1 channel.

As we will see, when we adopt the model which will be introduced in the next subsection, the N_1 decays immediately after produced by the inflaton decays. The lepton asymmetry is then given by

$$\begin{aligned} \frac{n_L}{s} &\simeq \epsilon_1 \frac{n_{N_1}}{s} \\ &\simeq 3 \times 10^{-10} B_r \left(\frac{T_R}{10^6 \text{ GeV}} \right) \left(\frac{M_1}{m_\chi} \right) \times \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}, \end{aligned} \quad (2.43)$$

where we have used the asymmetry parameter ϵ_1 given in Eq. (2.15). Notice that there is no wash-out effect of the produced lepton asymmetry,¹¹ since N_1 is out of thermal equilibrium.

Let us mention one point here. Although we will assume an explicit model in the next subsection to make the subsequent discussion concrete, the formula of the lepton asymmetry in Eq. (2.43) is a generic one, unless $M_1 \lesssim T_R$ or N_1 has a very long lifetime and dominates the energy density of the universe before it decays. If $M_1 \lesssim T_R$, the N_1

¹¹Lepton-number violating 2-body scatterings mediated by N_i are out of thermal equilibrium as long as $\sum_i m_{\nu i}^2 \lesssim (10 \text{ keV})^2 (10^9 \text{ GeV}/T_R)$ [75].

is produced more or less by thermal scatterings, and the resultant lepton asymmetry is expected to be reduced to the case discussed in Sec. 2.2. As for the case where N_1 dominates the energy density of the universe before its decay, we give a brief discussion in Appendix 2.A.

As discussed in Sec. 1.4, the sphaleron process converts this lepton asymmetry into baryon asymmetry as $n_B/s \simeq 0.35 \times n_L/s$. In order to explain the observed baryon asymmetry $n_B/s \simeq (0.4\text{--}1) \times 10^{-10}$, we should have lepton asymmetry

$$\frac{n_L}{s} \simeq (1 - 3) \times 10^{-10}. \quad (2.44)$$

From Eqs. (2.41) and (2.43), we can see that the reheating temperature of the inflation T_R is bounded from below as $T_R \gtrsim 10^6$ GeV, since otherwise the produced lepton asymmetry is too small as $n_L/s < 10^{-10}$.

2.3.2 Froggatt-Nielsen model for neutrino

As we have just shown, the amount of generated lepton asymmetry depends on the mass of the inflaton m_χ and the reheating temperature T_R (and the branching ratio B_r), as well as the mass of the right-handed neutrino M_1 . Before discussing the inflaton sector, we here introduce the Froggatt-Nielsen (FN) model [68] in order to settle the mass (and decay rate) of the right-handed neutrino and to make the discussion in the subsequent subsections concrete.

The FN model is one of the most attractive framework for explaining the observed hierarchies in quark and lepton mass matrices. This model is based on an $U(1)_{\text{FN}}$ symmetry that is broken by the vacuum-expectation value of Ξ , $\langle \Xi \rangle \neq 0$. Here Ξ is a gauge singlet field carrying the FN charge $Q_\Xi = -1$. Then, all Yukawa couplings are realized as nonrenormalizable interactions including Ξ , and are given by the following form;

$$\begin{aligned} W &= \widetilde{y}_{ij} \left(\frac{\langle \Xi \rangle}{M_G} \right)^{Q_i + Q_j} \psi_i \psi_j H_{u(d)} \\ &= \widetilde{y}_{ij} \varepsilon^{Q_i + Q_j} \psi_i \psi_j H_{u(d)}, \end{aligned} \quad (2.45)$$

where $\widetilde{y}_{ij}$ are $\mathcal{O}(1)$ coupling constants, Q_i are the FN charges of various chiral superfields ψ_i and $\varepsilon \equiv \langle \Xi \rangle / M_G$. Here, we have assumed that Higgs doublets H_u and H_d (which couple to up-type and down-type quarks, respectively) have zero FN charges. The observed mass hierarchies for quarks and charged leptons are well explained by taking suitable FN charges for them. For instance, we assign FN charges $(a + 1, a, a)$ for lepton doublets L_i , while giving charges $(2, 1, 0)$ to the right-handed charged leptons $\overline{E}_i$, with $\varepsilon \simeq 0.05\text{--}0.1$ [76, 77].

ψ_i	L_1	L_2	L_3	$\overline{E}_1$	$\overline{E}_2$	$\overline{E}_3$	N_1	N_2	N_3	Ξ
Q_i	$a+1$	a	a	2	1	0	d	c	b	-1

Table 2.1: The Froggatt-Nielsen charges of various supermultiplets. $a = 0$ or 1 . We assume $d > c \geq b \geq 0$, i.e., $M_1 \ll M_2, M_3$.

The charges are listed in Table. 2.1. We will take $a = 0$ or 1 according to Ref. [76]. The Charges of the quarks are also determined in the framework of the SU(5) grand unified theory [76].

Let us apply the above mechanism to the neutrino sector as well. The mass matrix for the heavy right-handed neutrinos N_i is given by

$$M_{Rij} = \widetilde{\xi}_{ij} \varepsilon^{Q_i+Q_j} M_0, \quad (2.46)$$

where M_0 represents some right-handed neutrino mass scale and $\widetilde{\xi}_{ij}$ are coupling constants of order unity like $\widetilde{y}_{ij}$. Charges for right-handed neutrinos are also found in Table. 2.1. We take $d > c \geq b \geq 0$, i.e., $M_1 \ll M_2, M_3$. Hereafter, we will take a basis where the mass matrix for the charged leptons is diagonal.¹² Then, the neutrino Dirac mass matrix m_D and the right-handed neutrino mass matrix M_R are given by the following forms;

$$\begin{aligned} m_D &= \langle H_u \rangle \begin{pmatrix} \varepsilon^d & 0 & 0 \\ 0 & \varepsilon^c & 0 \\ 0 & 0 & \varepsilon^b \end{pmatrix} \begin{pmatrix} \widetilde{y}_{11} & \widetilde{y}_{12} & \widetilde{y}_{13} \\ \widetilde{y}_{21} & \widetilde{y}_{22} & \widetilde{y}_{23} \\ \widetilde{y}_{31} & \widetilde{y}_{32} & \widetilde{y}_{33} \end{pmatrix} \begin{pmatrix} \varepsilon^{a+1} & 0 & 0 \\ 0 & \varepsilon^a & 0 \\ 0 & 0 & \varepsilon^a \end{pmatrix}, \\ M_R &= M_0 \begin{pmatrix} \varepsilon^d & 0 & 0 \\ 0 & \varepsilon^c & 0 \\ 0 & 0 & \varepsilon^b \end{pmatrix} \begin{pmatrix} \widetilde{\xi}_{11} & \widetilde{\xi}_{12} & \widetilde{\xi}_{13} \\ \widetilde{\xi}_{12} & \widetilde{\xi}_{22} & \widetilde{\xi}_{23} \\ \widetilde{\xi}_{13} & \widetilde{\xi}_{23} & \widetilde{\xi}_{33} \end{pmatrix} \begin{pmatrix} \varepsilon^d & 0 & 0 \\ 0 & \varepsilon^c & 0 \\ 0 & 0 & \varepsilon^b \end{pmatrix}, \end{aligned} \quad (2.47)$$

where $(m_D)_{i\alpha}$ is defined as $W = (m_D)_{i\alpha} N_i L_\alpha$ ($\alpha = e, \mu, \tau$). The neutrino mass matrix is then given by

$$\begin{aligned} m_\nu &= m_D^T \frac{1}{M_R} m_D \\ &= \frac{\varepsilon^{2a} \langle H_u \rangle^2}{M_0} \begin{pmatrix} \varepsilon & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} (\{\widetilde{y}_{ij}\})^T (\{\widetilde{\xi}_{ij}\})^{-1} (\{\widetilde{y}_{ij}\}) \begin{pmatrix} \varepsilon & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \frac{\varepsilon^{2a} \langle H_u \rangle^2}{M_0} \begin{pmatrix} \mathcal{O}(\varepsilon^2) & \mathcal{O}(\varepsilon) & \mathcal{O}(\varepsilon) \\ \mathcal{O}(\varepsilon) & \mathcal{O}(1) & \mathcal{O}(1) \\ \mathcal{O}(\varepsilon) & \mathcal{O}(1) & \mathcal{O}(1) \end{pmatrix}. \end{aligned} \quad (2.48)$$

¹²Here, one might wonder if the mixing matrix from the charged lepton sector would change the discussion, since the mass matrix for the charged leptons has off-diagonal elements in the above FN mechanism. However, the correction from this effect yields higher order terms in ε , and hence we can safely neglect it.

As shown in Ref. [76], this mass matrix can naturally lead to a large ν_μ - ν_τ mixing angle, which is suggested from the atmospheric neutrino oscillation [10]. It is remarkable that the FN charges of the right-handed neutrinos are completely canceled out in the neutrino mass matrix in Eq. (2.48) and hence the hierarchy in the neutrino mass matrix is determined only by the charges of the lepton doublets, $(a+1, a, a)$.

In this model, the mass scale of the right-handed neutrino M_0 is estimated as

$$\begin{aligned} M_0 &\sim \varepsilon^{2a} \frac{\langle H_u \rangle^2}{m_{\nu 3}} \\ &\sim \begin{cases} 0.6 \times 10^{15} \text{ GeV} & \text{for } a = 0, \\ (1-6) \times 10^{12} \text{ GeV} & \text{for } a = 1. \end{cases} \end{aligned} \quad (2.49)$$

Here, we have used $m_{\nu 3} \simeq 0.05 \text{ eV}$ and $\varepsilon \simeq 0.05 - 0.1$. Then the mass of the lightest right-handed neutrino M_1 is given by

$$\begin{aligned} M_1 &\sim M_0 \varepsilon^{2d} \\ &\sim \begin{cases} (1-6) \times 10^{12} \text{ GeV} & \text{for } a+d=1, \\ (0.3-6) \times 10^{10} \text{ GeV} & \text{for } a+d=2. \end{cases} \end{aligned} \quad (2.50)$$

In the following analysis, we only consider the case $a+d=1$ or $a+d=2$. Thus, our assumption of out-of-equilibrium condition $M_1 \gtrsim T_R$ is justified as far as $T_R \lesssim 10^9 \text{ GeV}$. On the other hand, the total decay width of the N_1 , Γ_{N_1} , is given by

$$\begin{aligned} \Gamma_{N_1} &\sim \frac{1}{4\pi} \varepsilon^{2(a+d)} M_1 \\ &\sim \begin{cases} (0.3-5) \times 10^9 \text{ GeV} & \text{for } a+d=1, \\ (0.02-5) \times 10^5 \text{ GeV} & \text{for } a+d=2. \end{cases} \end{aligned} \quad (2.51)$$

In deriving the formula of the lepton asymmetry in Eq. (2.43), we have assumed that the N_1 decays immediately after produced in inflaton decay, which corresponds to $\Gamma_{N_1} \gg \Gamma_\chi$. (Γ_χ is the decay rate of the inflaton χ .) In terms of the reheating temperature $T_R \simeq (\pi^2 g_*/90)^{-1/4} \sqrt{M_G \Gamma_\chi}$, this requirement corresponds to

$$T_R \ll 2 \times 10^{10} \text{ GeV} \left(\frac{\Gamma_{N_1}}{10^3 \text{ GeV}} \right)^{1/2}. \quad (2.52)$$

Therefore, our assumption is again guaranteed for $T_R \lesssim 10^9 \text{ GeV}$.

Finally, let us estimate the asymmetric parameter ϵ_1 in the present FN model. From Eq. (2.9), it is given by [60]

$$\begin{aligned} \epsilon_1 &\sim \frac{3}{8\pi} \frac{1}{(\varepsilon^{a+d})^2} \left[\left(\varepsilon^{a+d} \cdot \varepsilon^{a+c} \right)^2 \frac{\varepsilon^{2d}}{\varepsilon^{2c}} + \left(\varepsilon^{a+d} \cdot \varepsilon^{a+b} \right)^2 \frac{\varepsilon^{2d}}{\varepsilon^{2b}} \right] \\ &\sim \frac{3}{8\pi} \varepsilon^{2(a+d)} \\ &\sim \frac{3}{8\pi} \frac{m_{\nu 1} M_1}{\langle H_u \rangle^2}. \end{aligned} \quad (2.53)$$

We see that the above estimation agrees with the general result for ϵ_1 in Eq. (2.13), with $\delta_{\text{eff}} \sim 1$. Thus, we can use the formula of the lepton asymmetry in Eq. (2.43) consistently.

2.3.3 Hybrid inflation

In this subsection we perform a detailed analysis on hybrid inflation models and examine whether they can produce sufficient lepton asymmetry to account for the baryon asymmetry in the present universe, avoiding the overproduction of the gravitinos.

2.3.3.a hybrid inflation with a $B - L$ symmetry

Before discussing hybrid inflation models, let us first show a particle-physics model for the heavy right-handed neutrinos N_i . A simple extension of the SUSY standard model is given by considering a gauged $U(1)_{B-L}$ symmetry, in which, as mentioned in Sec. 1.1, right-handed neutrinos N_i are necessary to cancel $B - L$ gauge anomaly. We introduce standard-model gauge-singlet supermultiplets Ψ and $\bar{\Psi}$ carrying $B - L$ charges $+2$ and -2 , respectively, and suppose that the $B - L$ symmetry is spontaneously broken by the condensations $\langle \Psi \rangle = \langle \bar{\Psi} \rangle$ at high energies. ($\langle \Psi \rangle = \langle \bar{\Psi} \rangle$ is required from the D -term flatness condition of the $U(1)_{B-L}$.) Then the heavy neutrinos N_i , which carry $B - L$ charge -1 , acquire Majorana masses through the following superpotential:¹³

$$W = \frac{1}{2} \xi_i \Psi N_i N_i. \quad (2.54)$$

Here, we have assumed that Ψ and $\bar{\Psi}$ have zero FN charges. Then the right-handed neutrino scale M_0 in Eq. (2.49) is given by

$$M_0 = \langle \Psi \rangle = \langle \bar{\Psi} \rangle. \quad (2.55)$$

Thus, the right-handed neutrino mass scale M_0 derived in the FN model in Eq. (2.49) is explained in term of the $B - L$ breaking scale about $\mathcal{O}(10^{15})$ GeV or $\mathcal{O}(10^{12})$ GeV.

A superpotential causing the $B - L$ breaking is given by

$$W = \Phi \left(\lambda \Psi \bar{\Psi} - \mu^2 \right), \quad (2.56)$$

where Φ is a gauge-singlet supermultiplet, λ a coupling constant and μ a dimensionful mass parameter. (We will take a basis where μ^2 and λ are real and positive by using

¹³Leptogenesis in the hybrid inflation discussed in Ref. [78] assumes a nonrenormalizable superpotential $W = (g'_i/2) N_i N_i \Psi \Psi$, where the $B - L$ charge of Ψ is taken to be $+1$.

the phase rotations of Φ and $\Psi\bar{\Psi}$.) Notice that this superpotential possesses a $U(1)$ R -symmetry where the Φ and $\Psi\bar{\Psi}$ have $U(1)_R$ charges 2 and 0, respectively. The potential for scalar components of the supermultiplets Φ , Ψ and $\bar{\Psi}$ is given by, in supergravity,

$$V = e^{K/M_G^2} \left\{ \left(\frac{\partial^2 K}{\partial Z_i \partial Z_j^*} \right)^{-1} D_{Z_i} W D_{Z_j^*} W^* - 3 \frac{1}{M_G^2} |W|^2 \right\} + D \text{ terms}, \quad (2.57)$$

where

$$D_{Z_i} W = \frac{\partial W}{\partial Z_i} + \frac{1}{M_G^2} \frac{\partial K}{\partial Z_i} W, \quad (2.58)$$

and Z_i denote scalar components of supermultiplets. We assume the R -invariant Kähler potential for Φ , Ψ and $\bar{\Psi}$

$$K = |\Phi|^2 + |\Psi|^2 + |\bar{\Psi}|^2 + \frac{\kappa_1}{4M_G^2} |\Phi|^4 \dots, \quad (2.59)$$

where the ellipsis denotes higher-order terms which we neglect in the present analysis. From the above scalar potential, we have the following SUSY-invariant vacuum (hereafter, we denote the scalar components of the supermultiplets Φ and Ψ by the same symbols as the corresponding supermultiplets):

$$\langle \Psi \rangle = \langle \bar{\Psi} \rangle = \sqrt{\frac{\mu^2}{\lambda}}, \quad \langle \Phi \rangle = 0, \quad (2.60)$$

where we have chosen $\langle \Psi \rangle$ to be real and positive by using the $U(1)_{B-L}$ rotation.

It is quite interesting to observe that the superpotential (2.56) is nothing but the one proposed in Refs. [79, 80] for a SUSY hybrid inflation model. In this context, the real part of Φ is identified with the inflaton field $\varphi/\sqrt{2}$. The scalar potential is minimized at $\Psi = \bar{\Psi} = 0$ when φ is larger than the following critical value:

$$\varphi_C \equiv \sqrt{\frac{2\mu^2}{\lambda}}, \quad (2.61)$$

and hybrid inflation occurs for $\varphi_C < \varphi \lesssim M_G$ and $k \equiv -\kappa_1 \geq 0$. Including one-loop corrections [79], the potential for the inflaton φ is given by, for $\varphi > \varphi_C$,

$$\begin{aligned} V \simeq & \mu^4 \left[1 + \frac{k}{2} \left(\frac{\varphi}{M_G} \right)^2 + \frac{1}{16} (4k^2 + 7k + 2) \left(\frac{\varphi}{M_G} \right)^4 \right] \\ & + \frac{\lambda^4}{128\pi^2} \left[2\varphi_C^4 \ln \left(\frac{\lambda^2 \varphi^2}{2\Lambda^2} \right) + (\varphi^2 - \varphi_C^2)^2 \ln \left(1 - \frac{\varphi_C^2}{\varphi^2} \right) + (\varphi^2 + \varphi_C^2)^2 \ln \left(1 + \frac{\varphi_C^2}{\varphi^2} \right) \right], \end{aligned} \quad (2.62)$$

where Λ denotes the renormalization scale. Here, we have included the higher order terms which come from the Kähler potential (2.59) up to $\mu^4(\varphi/M_G)^4$ term, since the initial value of the inflaton field is close to M_G .¹⁴

Let us now discuss the inflation dynamics. The slow-roll conditions for inflation are given by [28]

$$\begin{aligned} \frac{1}{2} \left(M_G \frac{V'}{V} \right)^2 &< 1, \\ M_G^2 \frac{|V''|}{V} &< 1, \end{aligned} \quad (2.63)$$

where the prime denotes the derivative with the inflaton field φ . These conditions are satisfied when $\varphi > \varphi_C$, $\lambda < 1$ and $0 \leq k < 1$. Therefore, while the inflaton φ rolls down along the potential in Eq. (2.62) from φ_I ($\varphi_C < \varphi_I \lesssim M_G$) to φ_C , the vacuum energy μ^4 of the potential dominates the energy of the universe and hence the hybrid inflation takes place [81].

After the inflation ends, the vacuum energy is transferred into the energies of the coherent oscillations of the following two fields: the inflaton φ and a scalar field $\Sigma = (\Psi + \bar{\Psi})/\sqrt{2}$. Notice that the inflaton φ forms a massive supermultiplet together with the Σ field in the vacuum in Eq. (2.60), whose masses are given by

$$\begin{aligned} m_\varphi = m_\Sigma &= \sqrt{2\lambda} \mu \\ &= \sqrt{2} \lambda \langle \Psi \rangle. \end{aligned} \quad (2.64)$$

The radiations of the universe are produced by the decays of the φ and/or Σ field. In order to estimate the reheating temperature T_R we should know the total decay rates of these scalar fields. Through the interactions in the superpotentials in Eqs. (2.54) and (2.56), the inflaton φ decays into scalar components $\widetilde{N}_1$ of the N_1 supermultiplet, if kinematically allowed, with the rate

$$\begin{aligned} \Gamma_\varphi &\simeq \Gamma(\varphi \rightarrow \widetilde{N}_1 \widetilde{N}_1) \\ &= \frac{1}{64\pi} m_\varphi \left(\frac{M_1}{\langle \Psi \rangle} \right)^2 \left(1 - \frac{4M_1^2}{m_\varphi^2} \right)^{1/2}, \end{aligned} \quad (2.65)$$

while the Σ field decays into scalar ($\widetilde{N}_1$) and fermionic (N_1) components of the N_1 with the rates

$$\Gamma(\Sigma \rightarrow \widetilde{N}_1 \widetilde{N}_1) = \frac{1}{16\pi} \frac{1}{m_\Sigma} \left(\frac{M_1^2}{\langle \Psi \rangle} \right)^2 \left(1 - \frac{4M_1^2}{m_\Sigma^2} \right)^{1/2}, \quad (2.66)$$

¹⁴However, we have neglected the $\mu^4(\varphi/M_G)^4$ terms coming from the higher order interactions in the Kähler potential (e.g., $K = \kappa' |\Phi|^6/M_G^4$), for simplicity.

$$\Gamma(\Sigma \rightarrow N_1 N_1) = \frac{1}{64\pi} m_\Sigma \left(\frac{M_1}{\langle \Psi \rangle} \right)^2 \left(1 - \frac{4M_1^2}{m_\Sigma^2} \right)^{3/2}. \quad (2.67)$$

Here, we have used $M_1 = \xi_1 \langle \Psi \rangle$ and $m_\varphi = m_\Sigma = \sqrt{2} \lambda \langle \Psi \rangle$.

Since the Σ field has a non-zero vacuum expectation value, it can decay also through nonrenormalizable interactions in the Kähler potential,

$$K = \sum_i \frac{c_i}{M_G^2} |\Sigma|^2 |\psi_i|^2, \quad (2.68)$$

where ψ_i denote supermultiplets of the SUSY standard-model particles including the right-handed neutrinos N_i , and c_i are coupling constants of order unity. Then the partial decay rate through these interactions is estimated as

$$\Gamma(\Sigma \rightarrow \psi_i \bar{\psi}_i) \simeq \frac{1}{8\pi} C \left(\frac{\langle \Psi \rangle}{M_G} \right)^2 m_\Sigma^3, \quad (2.69)$$

where $C = \sum c_i^2$ is a parameter of order unity. In the following analysis we take $C \simeq 1$, for simplicity. Comparing this rate (2.69) with the rates in Eqs. (2.66) and (2.67), we see that the total decay rate of the Σ field is determined by that in Eq. (2.67), i.e.,

$$\Gamma_\Sigma \simeq \Gamma(\Sigma \rightarrow N_1 N_1), \quad (2.70)$$

since $\Gamma(\Sigma \rightarrow N_1 N_1) > \Gamma(\Sigma \rightarrow \widetilde{N}_1 \widetilde{N}_1) \gg \Gamma(\Sigma \rightarrow \psi_i \bar{\psi}_i)$ for $2M_1 < m_\Sigma \ll M_1 (M_G / \langle \Psi \rangle)^2$.

We assume $2M_1 < m_\varphi = m_\Sigma$ as discussed in Sec. 2.3.1. Then the inflaton φ and the Σ field decay at the almost same time because of $\Gamma_\varphi \simeq \Gamma_\Sigma$. Thus, the reheating temperature T_R is estimated by the total decay width of the Σ field as

$$T_R \simeq \left(\frac{\pi^2}{90} \pi^2 \right)^{-1/4} \sqrt{\Gamma_\Sigma M_G}. \quad (2.71)$$

Here it should be noted that the branching ratio of the decay rate of the Σ field and that of the inflaton φ into two N_1 ($\widetilde{N}_1$) is almost unity, and hence we have $B_r \simeq 1$ in Eq. (2.43).

The above hybrid inflation must explain the following two observations:¹⁵

- (i) the e -fold number N_e of the present horizon, and
- (ii) the density fluctuations observed by the cosmic background explorer (COBE) satellite [82].

¹⁵The spectrum index in the hybrid inflation is predicted to be scale-invariant, $n_s \simeq 1$.

First, while the inflaton φ rolls down along the potential (2.62) from φ_{N_e} to φ_C , the scale factor of the universe increases by e^{N_e} . This e -fold number N_e is given by

$$N_e \simeq \int_{\varphi_C}^{\varphi_{N_e}} d\varphi \frac{V(\varphi)}{M_G^2 V'(\varphi)}. \quad (2.72)$$

In order to explain the present horizon scale, the e -fold number should be

$$N_e = 67 + \frac{1}{3} \ln \left(\frac{H_{\text{inf}} T_R}{M_G^2} \right), \quad (2.73)$$

where $H_{\text{inf}} = \sqrt{V}/(\sqrt{3}M_G) \simeq \mu^2/(\sqrt{3}M_G)$ denotes the Hubble parameter during the inflation. Next, the amplitude of the primordial density fluctuations $\delta\rho/\rho$ predicted by the hybrid inflation,

$$\frac{\delta\rho}{\rho} \simeq \frac{1}{5\sqrt{3}\pi} \frac{V^{3/2}(\varphi_{N_e})}{M_G^3 |V'(\varphi_{N_e})|}, \quad (2.74)$$

should be normalized by the data on anisotropies of the cosmic microwave background radiation observed by the COBE satellite [82], which gives the following constraint:

$$\frac{V^{3/2}(\varphi_{N_e})}{M_G^3 |V'(\varphi_{N_e})|} \simeq 5.3 \times 10^{-4}. \quad (2.75)$$

The parameters of the inflation are determined by the constraints in Eqs. (2.73) and (2.75). We have performed numerical calculations using the full scalar potential in Eq. (2.62). The scale μ of the hybrid inflation is determined for given couplings λ and k , and it is shown in Fig. 2.1.¹⁶ Here and hereafter, we exclude the region where $\varphi_{N_e} \gtrsim M_G$, because in that region the higher order terms of $(\varphi/M_G)^n$ become large and our effective treatment of the inflaton potential in Eq. (2.62) would be invalid.

In Fig. 2.2 and Fig. 2.3, we also show the $B-L$ breaking scale $\langle\Psi\rangle$ and the inflaton mass m_φ . It is interesting that, as shown in Fig. 2.2, the scale of the $B-L$ breaking is predicted as $\langle\Psi\rangle \simeq (1-5) \times 10^{15}$ GeV in a wide parameter region, $10^{-6} \lesssim \lambda \lesssim 10^{-2}$ and $k \lesssim 10^{-3}$, which is consistent with $M_0 \sim 10^{15}$ GeV (i.e., $a=0$) derived from the observed neutrino mass [see Eq. (2.49)]. On the other hand, the lower value of the $B-L$ breaking scale of $M_0 \sim 10^{12}$ GeV ($a=1$) cannot be obtained in the present hybrid inflation model.

The reheating temperature T_R depends on the mass of the right-handed neutrino, M_1 , since the decay rate of the Σ (and φ) depends on M_1 . We take $M_1 \simeq 10^{12}$ GeV ($a+d=1$) and $M_1 \simeq 3 \times 10^9$ GeV ($a+d=2$) for representation. The obtained reheating temperatures T_R are shown in Fig. 2.4 and Fig. 2.5.

¹⁶Although the scale μ slightly depends on the mass of the right-handed neutrino M_1 , we find no sizable difference in the scale μ between the cases of $a+d=1$ and $a+d=2$, which are discussed below. This is because M_1 enters in the calculation only through the small correction of $\ln(T_R)$ in Eq. (2.73).




Figure 2.1: The contour lines of the scale μ in the hybrid inflation model. The contour lines are shown by the dotted lines and corresponding values of μ are represented in unit of GeV. The upper bound on k from the requirement $\varphi_{N_e} < M_G$ is shown by the thick solid line.

Here, we require the reheating temperatures T_R to be lower than 10^8 GeV to avoid overproduction of the gravitinos in a relatively wide range of gravitino mass (see Sec. 1.5). It is found that, for the region of the inflaton mass $m_\varphi = m_\Sigma \gg 2M_1$, the reheating temperature $T_R \lesssim 10^8$ GeV is obtained only for the case $M_1 \simeq 3 \times 10^9$ GeV ($a + d = 2$). In the case of $M_1 \simeq 10^{12}$ GeV ($a + d = 1$), the desired low reheating temperature is obtained for the region $m_\varphi \leq 2M_1$ because the decay into N_1 is kinematically forbidden and the decay rate is determined by the suppressed decay width in Eq. (2.69). However, such cases are not interesting since the N_1 are not produced in the φ and Σ decays and leptogenesis does not take place.¹⁷

Now let us examine whether the leptogenesis works well or not in the above hybrid inflation model. Since the heavy Majorana neutrinos N_1 are produced in the decays of the inflaton φ and the Σ field, the mass of the inflaton should satisfy $m_\varphi = m_\Sigma > 2M_1$. As derived in Sec. 2.3.1, the ratio of the produced lepton number to the entropy n_L/s is given by Eq. (2.43). Here, we stress again that the branching ratio is automatically given by $B_r \simeq 1$ in the present hybrid inflation. We show in Fig. 2.6 and Fig. 2.7 the n_L/s for the cases $a + d = 1$ and $a + d = 2$, respectively.

¹⁷See, however, also Sec. 2.3.6.




Figure 2.2: The contour lines of the $\langle \Psi \rangle$ in the hybrid inflation model. The contour lines are shown by the dotted lines and corresponding values of $\langle \Psi \rangle$ are represented in unit of 10^{15} GeV.




Figure 2.3: The contour lines of the inflaton mass m_{φ} in the hybrid inflation model. The contour lines are shown by the dotted lines and corresponding values of m_{φ} are represented in unit of GeV.




Figure 2.4: The contour lines of the reheating temperature T_R in the hybrid inflation model for the case $M_1 \simeq 10^{12}$ GeV ($a + d = 1$). The contour lines are shown by the dotted lines and corresponding values of T_R are represented in unit of GeV. The upper bound on k from the requirement $\varphi_{N_e} < M_G$ is shown by the thick solid line. The k yielding $m_\varphi = m_\Sigma = 2M_1$ is also shown by the thick solid line.




Figure 2.5: The same as Fig. 2.4, but for the case $M_1 \simeq 3 \times 10^9$ GeV ($a + d = 2$).




Figure 2.6: The contour lines of the lepton asymmetry n_L/s in the hybrid inflation model for the case $M_1 \simeq 10^{12}$ GeV ($a+d=1$). The contour lines are shown by the thin solid lines and corresponding values of n_L/s are represented. We also show the contour lines of the reheating temperature by the dotted lines and corresponding values of T_R are represented in unit of GeV. The upper bound on k from the requirement $\varphi_{N_e} < M_G$ and the lower bound on k from $m_\varphi = m_\Sigma > 2M_1$ are both shown by the thick solid lines.




Figure 2.7: The same as Fig. 2.6, but for the case $M_1 \simeq 3 \times 10^9$ GeV ($a+d=2$).

First, we consider the case of $M_1 \simeq 10^{12}$ GeV ($a + d = 1$). We find from Fig. 2.6 that the lepton asymmetry enough to explain the present baryon asymmetry can be generated in a wide parameter region. However, we have a too high reheating temperature of $T_R \simeq 10^9\text{--}10^{12}$ GeV as mentioned before. Therefore, only the small region of $m_\Sigma \simeq 2M_1$, where the reheating temperature becomes as small as $T_R \simeq 10^6\text{--}10^8$ GeV due to the phase space suppression,¹⁸ is free from the cosmological gravitino problem for a wide range of gravitino mass. We can obtain in this very narrow region the required lepton asymmetry of $n_L/s \simeq (1\text{--}3) \times 10^{-10}$ to account for the present baryon asymmetry.

Next, we consider the case of $M_1 \simeq 3 \times 10^9$ GeV ($a + d = 2$). It is found from Fig. 2.7 that the required lepton asymmetry of $n_L/s \simeq 10^{-10}$ as well as the low enough reheating temperature of $T_R \simeq 10^7\text{--}10^8$ GeV are naturally offered in the region of $k \lesssim 10^{-3}$ and $\lambda \simeq 10^{-6}\text{--}10^{-3}$. Therefore, we conclude that the hybrid inflation with $M_1 \simeq 3 \times 10^9$ GeV can produce a sufficient baryon asymmetry, giving a reheating temperature low enough to solve the cosmological gravitino problem. However, even lower reheating temperature of $T_R \simeq 10^6$ GeV, which is required to avoid the cosmological difficulty for a very wide range of gravitino mass $m_{3/2} \simeq 10$ MeV–10 TeV, is achieved only in the narrow parameter region of $m_\varphi = m_\Sigma \simeq 2M_1$, where T_R is reduced due to the phase suppression as in the previous case ($a + d = 1$).

2.3.3.b hybrid inflation without a $B - L$ symmetry

We have, so far, identified the U(1) gauge symmetry in the hybrid inflation model with the $B - L$ symmetry. We now consider the case where the U(1) symmetry is not related to the $B - L$ symmetry and even completely decoupled from the SUSY standard-model sector. The role of the U(1) gauge symmetry here is only to eliminate an unwanted flat direction in the superpotential in Eq. (2.56).

We reanalyze the leptogenesis in hybrid inflation in the absence of the superpotential in Eq. (2.54). In this case the Σ field decays through the nonrenormalizable interactions in Eq. (2.68). On the other hand, the decay of the inflaton φ is much suppressed due to the absence of the interaction in Eq. (2.54). Thus we introduce a new interaction in the Kähler potential as

$$K = \frac{h}{M_G} \Phi^* H_u H_d + \text{H.c.} . \quad (2.76)$$

Through this interaction the inflaton φ can decay faster than the Σ field for the coupling $h \gtrsim C (\langle \Psi \rangle / M_G)^2 \simeq 10^{-6}$, and the reheating temperature T_R is given by the decay of the

¹⁸In this region, we should include the decay rates in Eqs. (2.66) and (2.69) in estimating T_R .

Σ field. (Notice that the reheating temperature is determined by the slower decay rate.) Since the decay rate of the Σ field in Eq. (2.69) is very small compared with Eq. (2.67), the reheating temperature T_R becomes much lower than in the previous model. We show the obtained reheating temperature T_R in Fig. 2.8.

The inflation dynamics is almost the same as in the previous hybrid inflation model. The differences in the results for parameters, μ , $\langle\Psi\rangle$ and m_φ , only come from the small correction of $\ln(T_R)$ in Eq. (2.73), and we find no sizable difference in these values between the present and the previous models.

Now let us turn to estimate the lepton asymmetry in this hybrid inflation model. First, we find that too small lepton asymmetry is obtained for $M_1 \simeq 3 \times 10^9$ GeV ($a + d = 2$). This is because the T_R and M_1 are too low to produce enough lepton asymmetry [see Eq. (2.43)]. Hence, we concentrate on the case of $M_1 \simeq 10^{12}$ GeV ($a + d = 1$) in the following analysis.

We show the obtained lepton asymmetry n_L/s in Fig. 2.9. Here, we have assumed $B_r = 1$. Notice that the Σ field decays not only into the heavy neutrinos N_1 but also into the SUSY standard-model particles through the nonrenormalizable interactions in Eq. (2.68) and hence $B_r \simeq 1$ is not automatic in this model. Therefore, the lepton asymmetry shown in Fig. 2.9 is understood as a maximal value. It is found that the required lepton asymmetry to account for the empirical baryon asymmetry can be generated in a wide parameter region $k \lesssim 10^{-2}$ and $\lambda \simeq 10^{-3} - 10^{-2}$ with the reheating temperature of $T_R \simeq 10^6 - 10^8$ GeV. It is remarkable that we can obtain $n_L/s \simeq 10^{-10}$ and $T_R \simeq 10^6$ GeV simultaneously for $\lambda \simeq 10^{-3}$ and $k \lesssim 10^{-3}$. The overproduction of gravitinos can be avoided in the full gravitino mass region of $m_{3/2} \simeq 10$ MeV–10 TeV with such a low reheating temperature $T_R \simeq 10^6$ GeV.¹⁹

2.3.4 New inflation

In the previous subsection, we have seen that the SUSY hybrid inflation can successfully produce the lepton asymmetry to explain the baryon number in the present universe, even with low reheating temperatures of $T_R \simeq 10^6 - 10^8$ GeV. It was found, however, that relatively small couplings of $k \lesssim 10^{-2}$ and $\lambda \lesssim 10^{-2}$ are necessary to realize such low reheating temperatures. Although this is not so problematic, it is important to consider also other inflation models which naturally realize a low reheating temperature. The new inflation [83] is a well-known candidate for such an inflation. In this subsection, therefore,

¹⁹Although gravitinos are also produced in the reheating process of the inflaton φ decay, they are diluted by the subsequent entropy production of the Σ decays and become negligible.




Figure 2.8: The contour lines of the reheating temperature T_R in the hybrid inflation model without the $B-L$ symmetry. The contour lines are shown by the dotted lines and corresponding values of T_R are represented in unit of GeV.




Figure 2.9: The contour lines of the lepton asymmetry n_L/s in the hybrid inflation model without the $B-L$ symmetry for the case $M_1 \simeq 10^{12}$ GeV ($a+d=1$). The contour lines are shown by the thin solid lines and corresponding values of n_L/s are represented. Here we have assumed the branching ratio $B_r(\Sigma \rightarrow N_1 N_1)$ is $B_r = 1$ for the estimation of n_L/s . We also show the contour lines of the reheating temperature by the dotted lines and corresponding values of T_R in units of GeV. The upper bound on k from the requirement $\varphi_{N_e} < M_G$ and the lower bound on k from $m_\varphi = m_\Sigma > 2M_1$ are both shown by the thick solid lines.

we investigate the leptogenesis in the new inflation.

In order to make the discussion concrete, we propose a SUSY new inflation model which has the following superpotential W and Kähler potential K :

$$W = X(v^2 - g \frac{\Phi^n}{M_G^{n-2}}), \quad (2.77)$$

$$K = |\Phi|^2 + |X|^2 + \frac{\kappa_1}{4M_G^2}|\Phi|^4 + \frac{\kappa_2}{M_G^2}|\Phi|^2|X|^2 + \frac{\kappa_3}{4M_G^2}|X|^4 + \dots, \quad (2.78)$$

where Φ and X denote supermultiplets, v is the energy scale of the inflation, g , κ_1 , κ_2 and κ_3 are constants of order unity, and the ellipsis denotes higher order terms. The superpotential in Eq. (2.77) is naturally obtained, for example, by imposing $U(1)_R \times Z_n$ symmetry. Hereafter, we will take g and v^2 real and positive by using the phase rotations of X and Φ .

The SUSY vacuum of the scalar potential is obtained from Eqs. (2.57), (2.77) and (2.78), which is given by

$$\langle \Phi \rangle = M_G \left(\frac{v^2}{gM_G^2} \right)^{1/n}, \quad \langle X \rangle = 0. \quad (2.79)$$

Here and hereafter, we use the same symbols Φ and X for the scalar components of corresponding supermultiplets. The scalar potential for $|\Phi|, |X| \ll M_G$ is given by

$$\begin{aligned} V \simeq & v^4 + (1 - \kappa_2) v^4 \left(\frac{|\Phi|}{M_G} \right)^2 - g v^2 \frac{\Phi^n + \Phi^{*n}}{M_G^{n-2}} + g^2 \frac{|\Phi|^{2n}}{M_G^{2n-4}} \\ & - \kappa_3 v^4 \left(\frac{|X|}{M_G} \right)^2. \end{aligned} \quad (2.80)$$

Here, we have neglected irrelevant higher order terms. As for the X field, it is found that if $\kappa_3 < -1/3$, X receives a positive mass squared which is larger than H_{inf}^2 , where $H_{\text{inf}} = v^2/(\sqrt{3}M_G)$ is the Hubble parameter during the new inflation. Thus X settles down at $X = 0$ for $\kappa_3 < -1/3$. Hereafter, we assume $\kappa_3 < -1/3$ and set $X = 0$.

For $g > 0$ and $k \equiv \kappa_2 - 1 > 0$, we can identify the inflaton field φ with the real part of the scalar field Φ . The inflaton potential near the origin ($\varphi \simeq 0$) is obtained as,²⁰

$$V(\varphi) \simeq v^4 - \frac{k}{2} v^4 \left(\frac{\varphi}{M_G} \right)^2 - \frac{g}{2^{n/2-1}} v^2 \frac{\varphi^n}{M_G^{n-2}}. \quad (2.81)$$

The slow-roll conditions for the inflation in Eq. (2.63) are satisfied when

$$0 < k \lesssim 1, \quad (2.82)$$

$$0 < \varphi \lesssim \varphi_f \equiv \sqrt{2} M_G \left[\frac{1-k}{gn(n-1)} \frac{v^2}{M_G^2} \right]^{1/(n-2)}, \quad (2.83)$$

²⁰The new inflation model proposed in Ref. [84] offers a scalar potential for the inflaton field similar to Eq. (2.81). Thus, our discussion in this subsection can also be applied to that model.

and the new inflation takes place when the inflaton φ rolls down along the potential in Eq. (2.81) from $\varphi \simeq 0$ to φ_f .²¹

Now let us discuss the decay of the inflaton. We assume that the inflaton decays through nonrenormalizable interactions in the Kähler potential such as

$$K = \sum_i c'_i |\Phi|^2 |\psi_i|^2, \quad (2.84)$$

where ψ_i denote supermultiplets for SUSY standard-model particles including the right-handed neutrinos, and c'_i are coupling constants of order unity. (Here, a Kähler potential $K = (1/M_G)\Phi^* H_u H_d$ is forbidden by suitable charge assignment of the Z_n symmetry for the Higgs supermultiplets H_u and H_d .) With the interactions in the above Kähler potential, the decay rate Γ_φ is estimated as

$$\Gamma_\varphi \simeq \frac{1}{8\pi} C' \left(\frac{\langle \Phi \rangle}{M_G^2} \right)^2 m_\varphi^3, \quad (2.85)$$

where $C' \equiv \sum_i c_i'^2$ and we take $C' = 1$ in the following discussion. Here, the inflaton mass m_φ in the true vacuum Eq. (2.79) is given by

$$m_\varphi \simeq n g^{1/n} M_G \left(\frac{v}{M_G} \right)^{2-2/n}. \quad (2.86)$$

Then the reheating temperature T_R is given by $T_R \simeq (\pi^2 g_*/90)^{-1/4} \sqrt{\Gamma_\varphi M_G}$ with the decay rate in Eq. (2.85).

The scale v of the new inflation is determined from (i) the e -fold number N_e of the present horizon, (ii) the amplitude and (iii) the spectrum index n_s of the primordial density fluctuations $\delta\rho/\rho$. First, the number of e -foldings N_e should satisfy Eq. (2.73) to explain the present horizon scale. In the present new inflation model, N_e is given by

$$\begin{aligned} N_e &= \int_{\varphi_{N_e}}^{\varphi_f} d\varphi \frac{V(\varphi)}{M_G^2 |V'(\varphi)|} \\ &\simeq \int_{\varphi_{N_e}}^{\tilde{\varphi}} d\varphi \frac{v^4}{k v^4 \frac{\varphi}{M_G}} + \int_{\tilde{\varphi}}^{\varphi_f} d\varphi \frac{v^4}{\frac{ng}{2^{n/2-1}} v^2 \frac{\varphi^{n-1}}{M_G^n}}. \end{aligned} \quad (2.87)$$

where

$$\tilde{\varphi} = \sqrt{2} M_G \left(\frac{k}{ng} \frac{v^2}{M_G^2} \right)^{1/(n-2)}, \quad (2.88)$$

²¹For a successful new inflation, the initial value of the inflaton field φ should be taken near the local maximum of the potential $\varphi \simeq 0$ [85]. Although this seems unnatural, there have been found some dynamical mechanisms [86] to solve this initial value problem.




Figure 2.10: A Schematic behavior of the new inflation potential for $\varphi \ll M_G$.

and φ_{N_e} is the value of the φ field when the present universe crossed the horizon. Here we have assumed that $\varphi_{N_e} < \tilde{\varphi}$ (see Fig. 2.10). This condition corresponds to

$$k \gtrsim 1/[N_e(n-2)]. \quad (2.89)$$

From Eqs. (2.83), (2.87), and (2.88) we obtain the field value of φ_{N_e} :

$$\varphi_{N_e} \simeq \sqrt{2}M_G \left(\frac{k}{ng} \frac{v^2}{M_G^2} \right)^{1/(n-2)} \exp \left[-k \left(N_e + \frac{nk-1}{(n-2)k(1-k)} \right) \right]. \quad (2.90)$$

This φ_{N_e} should satisfy the COBE normalization in Eq. (2.75):

$$\frac{v^2}{kM_G\varphi_{N_e}} \simeq 5.3 \times 10^{-4}. \quad (2.91)$$

On the other hand, the COBE observations show the spectrum index n_s as $n_s = 1.0 \pm 0.2$ [82], while in the present new inflation model it is given by [84]

$$n_s \simeq 1 - 2k. \quad (2.92)$$

Therefore, we take $0.01 \lesssim k \lesssim 0.1$ in the following analysis assuming the coupling k not be extremely small. (Notice that this justifies the assumption $\varphi_{N_e} < \tilde{\varphi}$ [see Eq. (2.89)].)

From the above relations, we calculate the inflation scale v for given n , g and k , which are shown in Fig. 2.11. The vacuum expectation value $\langle \Phi \rangle$, the inflaton mass m_φ and the reheating temperature T_R are found in Fig. 2.12, Fig. 2.13, and Fig. 2.14, respectively. From Fig. 2.14, we find that the new inflation model with the power indices $n = 4, 5$ and 6 naturally offer the low reheating temperature $T_R \lesssim 10^8$ GeV. For the cases $n = 7, 8$, we obtain $T_R \lesssim 10^8$ GeV for the region $k \simeq 0.05$ – 0.1 and 0.07 – 0.1 , respectively.




Figure 2.11: The scale v of the new inflation for $g = 1$. We take the index n as $n = 4, 5, 6, 7$ and 8 from the bottom to the top.




Figure 2.12: The vacuum expectation value $\langle\Phi\rangle$ of the new inflation for $g = 1$. We take the index n as $n = 4, 5, 6, 7$ and 8 from the bottom to the top.




Figure 2.13: The inflation mass m_ϕ of the new inflation for $g = 1$. We take the index n as $n = 4, 5, 6, 7$ and 8 from the bottom to the top.




Figure 2.14: The reheating temperature T_R of the new inflation for $g = 1$. We take the index n as $n = 4, 5, 6, 7$ and 8 from the bottom to the top.

Now we turn to discuss the leptogenesis in this new inflation model. The ratio of the lepton-number density n_L to the entropy density s is again given by Eq. (2.43). Notice that the new inflation model with $n = 4$ gives such a low reheating temperature as $T_R \simeq 1 \text{ GeV} - 10 \text{ GeV}$ (see Fig. 2.14) that the required amount of lepton asymmetry cannot be generated, and hence we discard this case.

The obtained lepton asymmetry for the cases $n = 5, 6, 7$ and 8 are shown in Fig. 2.15 and Fig. 2.16 by taking $M_1 \simeq 3 \times 10^{11} \text{ GeV}$ ($a + d = 1$) and $M_1 \simeq 10^9 \text{ GeV}$ ($a + d = 2$), respectively. Here we have taken relatively smaller values of M_1 [see Eq. (2.50)] to obtain a wider allowed region, taking account of $\mathcal{O}(1)$ ambiguities in the FN model.

Interesting results are given for $M_1 \simeq 3 \times 10^{11} \text{ GeV}$. (Here, we find that the condition $m_\varphi > 2M_1$ excludes the regions $k < 1.2 \times 10^{-2}$ and $k > 4.8 \times 10^{-2}$ for the case $n = 5$.) From Fig. 2.14 and Fig. 2.15, we see that the sufficient lepton asymmetry can be generated for $n = 5, 6, 7$ and 8 with the low enough reheating temperature of $T_R \simeq 10^6 - 10^8 \text{ GeV}$. In particular, for $n = 5$, the required lepton asymmetry $n_L/s \simeq -10^{-10}$ is obtained for $B_r \simeq 1$ with low reheating temperature $T_R \simeq 10^6 \text{ GeV}$. Thus, we can avoid the overproduction of gravitinos for a wide range of gravitino mass $m_{3/2} \simeq 10 \text{ MeV} - 10 \text{ TeV}$ in this case.

On the other hand, when $M_1 \simeq 10^9 \text{ GeV}$ ($a + d = 2$), we find from Fig. 2.16 that the leptogenesis does not work well.




Figure 2.15: The lepton asymmetry produced via the decays of N_1 in the new inflation. The index n is taken as $n = 5, 6, 7$ and 8 from the bottom to the top. We take $M_1 \simeq 3 \times 10^{11}$ GeV ($a + d = 1$). For $n = 5$, the regions $k < 1.2 \times 10^{-2}$ and $k > 4.8 \times 10^{-2}$ are excluded since $m_\varphi \leq 2M_1$.




Figure 2.16: The same as Fig. 2.15, but for $M_1 \simeq 10^9$ GeV ($a + d = 2$).

2.3.5 Topological inflation

In this subsection, we discuss the leptogenesis in a topological inflation. If the vacuum expectation value of the inflaton exceeds the gravitational scale, the inflaton potential of the new inflation in Eq. (2.81) becomes nothing but the inflaton potential for the topological inflation [87].

We adopt the SUSY topological inflation model proposed in Ref. [88]. The superpotential and Kähler potential in that model are given by

$$W = v^2 X \left(1 - g \frac{\Phi^2}{M_G^2}\right), \quad (2.93)$$

$$K = |X|^2 + |\Phi|^2 + \frac{\kappa_1}{4M_G^2} |\Phi|^4 + \frac{\kappa_2}{M_G^2} |X|^2 |\Phi|^2 + \frac{\kappa_3}{4M_G^2} |X|^4 + \dots, \quad (2.94)$$

where v is the energy scale of the inflation and g , κ_1 , κ_2 and κ_3 are coupling constants of order unity. Hereafter, we take g and v to be real and positive by using the redefinition of the field X and Φ . These potentials possess $U(1)_R \times Z_2$ symmetry; the $U(1)_R$ charge of Φ and X are 0 and 2, respectively, and Φ (X) is odd (even) under the Z_2 . This discrete symmetry is crucial to allow a topological defect (in this case, a domain wall) and hence to realize the topological inflation [87].

From Eqs. (2.57), (2.93) and (2.94) we find a SUSY-invariant vacuum:

$$\langle X \rangle = 0, \quad \langle \Phi \rangle = \frac{1}{\sqrt{g}} M_G. \quad (2.95)$$

where the scalar components of the supermultiplets are denoted by the same letters as the corresponding supermultiplets. The potential for the region $|X|, |\Phi| \ll M_G$ is written approximately as

$$V \simeq v^4 \left\{ \left| 1 - g \frac{\Phi^2}{M_G^2} \right|^2 + (1 - \kappa_2) \frac{|\Phi|^2}{M_G^2} - \kappa_3 \frac{|X|^2}{M_G^2} \right\}. \quad (2.96)$$

Hereafter, we set $X = 0$ assuming $\kappa_3 < -1/3$ as we have done in the new inflation.

For $g > 0$ and $\kappa_2 < 1$, the inflaton field φ is identified with the real part of Φ and the potential around the origin is given by

$$V(\varphi) \simeq v^4 \left(1 - \frac{k}{2} \frac{\varphi^2}{M_G^2} \right), \quad (2.97)$$

where $k \equiv 2g + \kappa_2 - 1$.

A topological inflation takes place if the vacuum expectation value of Φ is of order of the gravitational scale M_G [87]. It was shown in Ref. [89] that, in order for a topological

defect (in our case, a domain wall) to inflate, its vacuum expectation value should be larger than the following critical value.

$$\langle \Phi \rangle \geq \langle \Phi \rangle_C \simeq 1.7 M_G. \quad (2.98)$$

The slow-roll conditions in Eq. (2.63) are satisfied when $0 < k < 1$ and $\varphi \lesssim \varphi_f$, where φ_f is expected to be of order of the gravitational scale. Hereafter, we take $\varphi_f = M_G$.

As for the decay of the inflaton φ , we assume that φ decays through the interactions in the Kähler potential as shown in Eq. (2.84), and hence the reheating temperature T_R of the topological inflation is again given by $T_R \simeq (\pi^2 g_*/90)^{-1/4} \sqrt{\Gamma_\varphi M_G}$ with the decay rate in Eq. (2.85). Here, the inflaton mass in the true vacuum in Eq. (2.95) is estimated as

$$m_\varphi \simeq 2\sqrt{g} \frac{v^2}{M_G} = \frac{2v^2}{\langle \Phi \rangle}. \quad (2.99)$$

As in the other inflation models, the scale of this topological inflation is determined by (i) the e -fold number N_e , (ii) the density fluctuations $\delta\rho/\rho$ and (iii) the spectrum index n_s . The e -fold number in the present model is estimated as

$$N_e \simeq \int_{\varphi_f}^{\varphi_{N_e}} d\varphi \frac{V}{M_G^2 V'} \simeq \frac{1}{k} \ln \left(\frac{\varphi_f}{\varphi_{N_e}} \right), \quad (2.100)$$

which should satisfy the present Hubble horizon scale in Eq. (2.73), while the COBE normalization in Eq. (2.75) requires

$$\frac{v^2}{k M_G \varphi_{N_e}} \simeq 5.3 \times 10^{-4}. \quad (2.101)$$

The spectrum index n_s is again given by $n_s \simeq 1 - 2k$, and hence we take $0.01 \lesssim k \lesssim 0.1$ as in the new inflation model.

From the above relations, we can calculate the inflation scale v and the mass m_φ , and hence the reheating temperature T_R . We show the results in Fig. 2.17. Here, we have taken $\langle \Phi \rangle = \langle \Phi \rangle_C \simeq 1.7 M_G$. A crucial difference between this model and the previous new inflation model is the reheating temperature. In this topological model, as shown in Fig. 2.17, relatively high reheating temperatures are obtained compared with those in the new inflation. This is because the large value of the vacuum expectation value $\langle \Phi \rangle$. (See Eq. (2.85).)

Finally, we show the obtained lepton asymmetry n_L/s in Fig. 2.18. Here, we have taken $M_1 \simeq 3 \times 10^{11}$ GeV ($a + d = 1$) and $M_1 \simeq 10^9$ GeV ($a + d = 2$) as in the new inflation model discussed in the previous subsection. We can see that for the both cases enough lepton asymmetry $n_L/s > 10^{-10}$ is obtained for $B_r \simeq 1$. On the other




Figure 2.17: The inflaton mass m_φ and the reheating temperature T_R of the topological inflation model. We take $\langle\Phi\rangle = \langle\Phi\rangle_C \simeq 1.7M_G$. The solid and dot-dashed lines represent m_φ and T_R , respectively.




Figure 2.18: The lepton asymmetries, n_L/s , in the topological inflation model for the cases of $M_1 \simeq 3 \times 10^{11}$ GeV ($a + d = 1$, the solid line) and $M_1 \simeq 10^9$ GeV ($a + d = 2$, the dot-dashed line).

hand, the requirement of a low enough reheating temperature ($T_R \lesssim 10^8$ GeV) is satisfied only in a small range of the coupling $0.092 \lesssim k \lesssim 0.1$. (The lowest reheating temperature $T_R = 6.3 \times 10^7$ GeV is obtained for $k = 0.1$.) For $k \simeq 0.1$, the spectrum index n_s deviates from the scale invariant one as $n_s \simeq 0.8$, and it will be testable in current and future satellite experiments on anisotropies of the cosmic microwave background radiation [90].

One final comment: in the case of $M_1 \simeq 3 \times 10^{11}$ GeV, we may accommodate a late-time entropy production of order 10^2 – 10^3 since the produced lepton asymmetry is as large as $n_L/s \simeq 10^{-7}$ as shown in Fig. 2.18. If this is the case, the energy density of the gravitinos is diluted by the factor 10^2 – 10^3 , which allows the region of $T_R \simeq 10^9$ GeV in Fig. 2.17.

2.3.6 Production of right-handed neutrinos at preheating

Here, we briefly mention the production of right-handed neutrinos at preheating epoch, which was investigated in Ref. [74].

So far, we have assumed that the reheating process after the inflation occurs in such a way that energy density of the zero mode of the inflaton decays perturbatively into light particles. However, the heavy right-handed neutrinos might be predominantly produced nonperturbatively, through the parametric resonance [91, 92].

In Ref. [74], production of heavy right-handed neutrinos via preheating of fermion [92] was discussed, adopting a chaotic inflation with the inflaton potential $V(\varphi) = (1/2)m_\varphi\varphi^2$. (m_φ is the inflaton mass.) The point is that, if there is a Yukawa coupling between the inflaton φ and heavy fermion X

$$\mathcal{L} = g\varphi\bar{X}X, \quad (2.102)$$

then the effective mass $m_{X\text{eff}}$ of the fermion X depends on time t through the oscillating inflaton $\varphi(t)$ during the inflaton-oscillating period:

$$m_{X\text{eff}}(t) = m_X + g\varphi(t). \quad (2.103)$$

Therefore, the effective mass $m_{X\text{eff}}(t)$ oscillates with time, and can even be zero if the amplitude of the oscillating inflaton is large enough, which is very crucial to produce very massive fermions heavier than the inflaton itself.

The production of the X fermion crucially depends on the parameter

$$q \equiv g^2 \frac{\varphi(0)^2}{4m_\varphi^2}. \quad (2.104)$$

where $\varphi(0)$ denotes the initial amplitude of the inflaton oscillation. An efficient production of very massive fermions requires a large value of q [74], and hence large enough initial amplitude $\varphi(0)$. In the case of chaotic inflation, $\varphi(0) \simeq M_G$ and $m_\varphi \simeq 10^{13}$ GeV can lead to $q \simeq g \times \mathcal{O}(10^{10}) \gg 1$.

The maximum mass of the produced fermion is given by [74]

$$(m_X)_{\max} \simeq 0.25g \varphi(0). \quad (2.105)$$

If $m_X \lesssim (m_X)_{\max}$, the effective mass $m_{X\text{eff}}(t)$ can vanish more than once [see Eq. (2.103)]²² and the production of X through parametric resonance occurs efficiently, whereas if otherwise the production is much less effective. The production continues during vanishing effective mass can be achieved. Namely, the production stops when the amplitude of the oscillating inflaton φ_0 becomes below the following critical value:

$$(\varphi_0)_{\text{crit}} = \frac{1}{g} m_X. \quad (2.106)$$

The energy density of the inflaton at this time is thus given by $\rho_\varphi = (1/2)m_\varphi^2 (\varphi_0)_{\text{crit}}^2$. On the other hand, the energy density of the produced fermion X at this frozen-out time is given by [74]

$$\rho_X \simeq \frac{1}{6\pi^2} m_X^2 m_\varphi^2. \quad (2.107)$$

After that, i.e., for $\varphi_0 < (\varphi_0)_{\text{crit}}$, the ratio of the energy density of the X to the inflaton φ , ρ_X/ρ_φ , is fixed. From the above equations, this ratio is simply given by

$$\frac{\rho_X}{\rho_\varphi} \simeq \frac{g^2}{3\pi^2}. \quad (2.108)$$

Now let us identify the X field to the right-handed neutrino N_1 . From above equation, the resultant lepton asymmetry after the reheating process of the inflation completes is given by

$$\frac{n_L}{s} \simeq \frac{3}{4} \epsilon_1 \left(\frac{T_R}{M_1} \right) \frac{g^2}{3\pi^2}. \quad (2.109)$$

Thus, the baryon asymmetry $n_B/s \simeq 0.35 \times n_L/s$ is given by

$$\frac{n_B}{s} \simeq 2g^2 \times 10^{-10} \left(\frac{T_R}{10^8 \text{ GeV}} \right) \times \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}, \quad (2.110)$$

²²Their numerical calculation shows [74] that the minimum of the inflaton amplitude $\varphi_0^{(-)}$ during the first oscillation is given by $\varphi_0^{(-)} \simeq -0.25\varphi(0)$.

where we have used Eq. (2.15).

Before closing this section, let us see what happens if we apply the above mechanism to the first one of the SUSY hybrid inflation models discussed in Sec. 2.3.3.a, where the inflaton has a Yukawa coupling to the right-handed neutrino. In this case, the parameter q is estimated as

$$\begin{aligned} q &\simeq \xi_1^2 \frac{\langle \Psi \rangle^2}{4m_\varphi^2} \\ &\simeq \frac{1}{16} \left(\frac{2M_1}{m_\varphi} \right)^2. \end{aligned} \quad (2.111)$$

Therefore, in the region we have studied, where the perturbative decay is kinematically allowed ($m_\varphi > 2M_1$), preheating takes place only through a narrow resonance, and hence only small fraction of the inflaton oscillating energy can be converted to heavy right-handed neutrinos via preheating. For the case of $M_1 \simeq 10^{12}$ GeV ($a + d = 1$), in the small coupling region $k \lesssim 10^{-4}$ and $\lambda \lesssim 10^{-4}$ where $m_\varphi \lesssim (1/2)M_1$ [see Fig. 2.3], preheating mechanism might work effectively, although it requires another detailed analysis.

2.4 Leptogenesis from $\widetilde{N}$ -dominant universe

In the previous sections, we have discussed the leptogenesis mechanisms by the decay of right-handed (s)neutrino, which is produced as a *particle*. However, there is a new possibility once we introduce the SUSY, that is, the condensation of the scalar component of the right-handed neutrino, $\widetilde{N}$ [25].²³ The lepton asymmetry is produced in the same way as the previous mechanisms, by the asymmetric decay of the right-handed “s”neutrino into leptons and anti-leptons. However, the thermal history of the universe in this scenario is quite different from the others. In this section,²⁴ we investigate this scenario in detail. In particular, we mainly discuss the case in which the coherent oscillation of the right-handed sneutrino $\widetilde{N}$ dominates the energy density of the universe before it decays. Remarkably, the amount of produced baryon asymmetry is determined almost only by the decay rate of the right-handed neutrino, whatever happened before the coherent oscillation dominates the universe. (Thus, the baryon asymmetry is independent of the details of the inflation !)

Furthermore, as a big bonus, the abundance of the gravitinos which are created in the reheating phase after inflation are diluted by the entropy production due to the decay of the coherent right-handed sneutrino, so that the cosmological gravitino problems can be avoided even when the reheating temperature T_R of the inflation is higher than 10^{11} GeV, in a wide range of the gravitino mass $m_{3/2} \sim 10$ MeV–10 TeV. In particular, this dilution of the gravitinos has great advantages in the gauge-mediated SUSY breaking (GMSB) models [45].

We will also comment on the cosmic density perturbation caused by the quantum fluctuation of the right-handed sneutrino, which may be detected in the future.

2.4.1 Lepton asymmetry from $\widetilde{N}$ -dominant universe

The main character in the present scenario is the right-handed sneutrino, $\widetilde{N}$. During inflation, it can acquire a large amplitude [25, 24, 93] if the Hubble expansion rate of the inflation H_{inf} is larger than the mass of the $\widetilde{N}$. (The mechanism which makes a scalar field develop a large initial amplitude during inflation will be explained in Sec. 3.2.2, in the context of the Affleck-Dine condensation.) Let us assume that there exists (at least) one right-handed neutrino with a mass lighter than H_{inf} , and that it develops a large expectation value during the inflation. Hereafter, we focus on the lightest right-handed sneutrino $\widetilde{N}_1$ for simplicity. (Possible contributions from the heavier right-handed

²³See also Refs. [24, 93], where the right-handed sneutrino is considered as an inflaton of the chaotic inflation.

²⁴This section is based on the work in a collaboration with H. Murayama and T. Yanagida [26].

sneutrinos $\widetilde{N}_2$ and $\widetilde{N}_3$ will be discussed in Sec. 2.4.4.b.) It is assumed here that the potential for the right-handed neutrino is given simply by the mass term

$$V = M_1^2 |\widetilde{N}_1|^2. \quad (2.112)$$

We do not settle the initial amplitude of the $\widetilde{N}_1$ at this stage, which will be discussed later. Because of the large amplitude of the $\widetilde{N}_1$, the lepton L and Higgs H_u doublets receive large masses, and the expectation values of their scalar components $\langle \widetilde{L} \rangle$ and $\langle H_u \rangle$ vanish.

The evolution of the $\widetilde{N}_1$ is as follows. After the end of the inflation, the Hubble parameter H decreases with cosmic time t as $H \propto t^{-1}$, and $\widetilde{N}_1$ begins to oscillate around the origin when H becomes smaller than the mass of the right-handed sneutrino M_1 . ($H \simeq M_1$.) Then, the coherent oscillation eventually decays when the Hubble expansion rate H becomes comparable to the decay rate $\Gamma_{\widetilde{N}_1}$ of $\widetilde{N}_1$. ($H \simeq \Gamma_{\widetilde{N}_1}$.) As explained in Sec. 2.1, the asymmetric decay of $\widetilde{N}_1$ into leptons and anti-leptons produces a net lepton asymmetry. The generated lepton number density is given by

$$n_L = \epsilon_1 M_1 |\widetilde{N}_{1d}|^2, \quad (2.113)$$

where $|\widetilde{N}_{1d}|$ is the amplitude of the oscillation when it decays, and ϵ_1 is given by Eq. (2.15).

The fate of the generated lepton asymmetry depends on whether or not the coherent oscillation of $\widetilde{N}_1$ dominates the energy density of the universe before it decays [25]. Here, we focus on the leptogenesis scenario from the universe dominated by $\widetilde{N}_1$. (We will mention the case where $\widetilde{N}_1$ does not dominate the universe in Sec. 2.4.4.d.) As we shall show soon, once the $\widetilde{N}_1$ dominant universe is realized, the present baryon asymmetry is determined only by the properties of the right-handed neutrino, whatever happened before the $\widetilde{N}_1$ dominates the universe. We first derive the amount of the generated lepton asymmetry just assuming that the $\widetilde{N}_1$ dominates the universe, and after that we will discuss the necessary conditions of the present scenario.

Once $\widetilde{N}_1$ dominates the universe before it decays, the universe is *reheated again* at $H \simeq \Gamma_{\widetilde{N}_1}$ by the decay of $\widetilde{N}_1$. The energy density of the resulting radiation, with a temperature T_{N_1} , is given by the following relation;

$$\begin{aligned} \frac{\pi^2}{30} g_* T_{N_1}^4 &= M_1^2 |\widetilde{N}_{1d}|^2 \\ &= 3M_G^2 \Gamma_{N_1}^2, \end{aligned} \quad (2.114)$$

while the entropy density is given by [see Eq. (B.17)]

$$s = \frac{2\pi^2}{45} g_* T_{N_1}^3. \quad (2.115)$$

From the Eqs. (2.15), (2.113), (2.114) and (2.115), the ratio of the lepton number density to the entropy density is given by the following very simple form;

$$\begin{aligned}\frac{n_L}{s} &= \frac{3}{4}\epsilon_1 \frac{T_{N_1}}{M_1} \\ &\simeq 1.5 \times 10^{-10} \left(\frac{T_{N_1}}{10^6 \text{ GeV}} \right) \times \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}.\end{aligned}\quad (2.116)$$

Here, it is assumed that the decay of the $\widetilde{N}_1$ occurs in an out-of-equilibrium way, namely, $T_{N_1} < M_1$, so that the produced lepton-number asymmetry not be washed out by lepton-number violating interactions mediated by N_1 . As explained in Sec. 1.4, this lepton asymmetry is partially converted into the baryon asymmetry through the sphaleron effect:

$$\begin{aligned}\frac{n_B}{s} &\simeq 0.35 \times \frac{n_L}{s} \\ &\simeq 0.5 \times 10^{-10} \left(\frac{T_{N_1}}{10^6 \text{ GeV}} \right) \times \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}.\end{aligned}\quad (2.117)$$

Therefore, as stressed previously, the baryon asymmetry in the present universe is indeed determined only by the decay temperature of the right-handed sneutrino T_{N_1} (and the effective CP violating phase δ_{eff}). Thus it is independent of unknown parameters of the inflation such as the reheating temperature T_R . Assuming the effective CP violating phase $\delta_{\text{eff}} (\leq 1)$ to be not too small, the empirical baryon asymmetry $n_B/s \simeq (0.4\text{--}1) \times 10^{-10}$ is obtained by taking

$$T_{N_1} \simeq 10^6 - 10^7 \text{ GeV}.\quad (2.118)$$

2.4.2 Necessary conditions

So far, we have required the following two conditions;

- (i) $\widetilde{N}_1$ dominates the universe before it decays.
- (ii) $\widetilde{N}_1$ decays in an out-of-equilibrium way.

Let us first consider the easier one, the second condition (ii). By taking the T_{N_1} in Eq. (2.118), the condition of the out-of-equilibrium decay is simply given by²⁵

$$M_1 > T_{N_1} \simeq 10^6 - 10^7 \text{ GeV}.\quad (2.119)$$

²⁵ The condition $M_1 > T_{N_1}$ is formally equivalent to

$$H|_{T=M_1} > \Gamma_{N_1},$$

if we take the usual relation $3M_G H^2 = (\pi^2/30)g_* T^4$. However, this interpretation is not appropriate in the present case, since the universe is not radiation-dominant for $H > \Gamma_{N_1}$. In any case, the condition Eq. (2.119) is necessary to avoid the thermal production of N_1 after the decay of the coherent $\widetilde{N}_1$.

Notice that this condition gives rise to a constraint on the Yukawa couplings $h_{1\alpha}$, since the temperature T_{N_1} is determined by the decay rate of the $\widetilde{N}_1$, and hence is related to the mass and couplings of $\widetilde{N}_1$. Actually, from Eq. (2.114) and $\Gamma_{N_1} = (1/4\pi) \sum_{\alpha} |h_{1\alpha}|^2 M_1$, the following relation is derived:

$$\sqrt{\sum_{\alpha} |h_{1\alpha}|^2} \simeq 5 \times 10^{-6} \left(\frac{T_{N_1}}{10^6 \text{ GeV}} \right)^{1/2} \left(\frac{T_{N_1}}{M_1} \right)^{1/2}. \quad (2.120)$$

Hence, it is found from Eq. (2.119) that Yukawa couplings $h_{1\alpha}$ of N_1 should be smaller than $\mathcal{O}(10^{-5})$. (Possible explanations for the smallness of $h_{1\alpha}$ will be discussed in Sec. 2.4.4.a.)

Now let us turn to discuss the first condition (i). In order to discuss whether or not $\widetilde{N}_1$ dominates the universe, it is necessary to consider the history of the universe before it decays. Here, we assume that the potential of the $\widetilde{N}_1$ is “flat” up to the Planck scale, namely, the potential is just given by the mass term $M_1^2 |\widetilde{N}_1|^2$ up to the Planck scale. (This may not be the case when the masses of the right-handed neutrinos are induced by a breaking of an additional gauge symmetry. We will discuss such a case in Sec. 2.4.4.d.)

Assuming the flatness of the $\widetilde{N}_1$ ’s potential up to the Planck scale (i.e., only the mass term), the initial amplitude of the oscillation is naturally given by $|\widetilde{N}_{1i}| \simeq M_G$, since above the Planck scale the scalar potential is expected to be exponentially lifted by the supergravity effects [see Eq. (2.57)].²⁶ Then the energy density of $\widetilde{N}_1$ when it starts the coherent oscillation is given by $\rho_{N_1} = M_1^2 |\widetilde{N}_{1i}|^2 \simeq M_1^2 M_G^2$.

The rest of the total energy density of the universe at $H = M_1$ is dominated by (I) the radiation or (II) the oscillating inflaton χ , depending on the M_1 and the decay rate of the inflaton Γ_{χ} . (See Fig. 2.19.)

If $\Gamma_{\chi} > M_1$ [case (I)], the inflaton decay has already completed before $H = M_1$, and the energy density of the radiation at $H = M_1$ is given by $\rho_{\text{rad}} = (\rho_{\text{total}} - \rho_{N_1}) \simeq 2M_1^2 M_G^2$. In this case, the oscillating $\widetilde{N}_1$ dominates the universe soon after it starts the oscillation and hence before its decay.

On the other hand, if $M_1 > \Gamma_{\chi}$ [case (II)], the reheating process of the inflation has not completed yet at $H = M_1$, and the inflaton χ is still oscillating around its minimum, whose energy density is given by $\rho_{\chi} \simeq 2M_1^2 M_G^2$. The ratio of the energy density of $\widetilde{N}_1$ to that of the inflaton, $\rho_{N_1}/\rho_{\chi} \simeq 1/2$, takes a constant value until either of these oscillations decays. Because the energy density of the radiation ρ_{rad} resulting from the inflaton decay is diluted faster than ρ_{N_1} , the oscillating $\widetilde{N}_1$ can dominate the universe if its decay rate Γ_{N_1} is slow enough compared with that of the inflaton Γ_{χ} ; $\Gamma_{\chi} \gg \Gamma_{N_1}$.²⁷

²⁶Even though it is possible that $\widetilde{N}_1$ has a larger initial amplitude $|\widetilde{N}_{1i}| > M_G$ (see, e.g., Ref [24, 93]), it depends on the scalar potential beyond the Planck scale, so that we do not discuss this possibility here.

²⁷More precisely, $\rho_{N_1} \geq \rho_{\text{rad}}$ occurs for $\Gamma_{N_1} \leq (1/2\sqrt{3})\Gamma_{\chi}$ in case (II).




Figure 2.19: Schematic behaviors of the energy densities. R denotes the scale factor of the expanding universe. It is found that the coherent oscillation $\tilde{N}_1$ can dominate the energy density of the universe if the condition $\Gamma_\chi \gg \Gamma_{N_1}$ ($T_R \gg T_{N_1}$) is satisfied.

Therefore, the condition for $\tilde{N}_1$ to dominate the universe is just given by $\Gamma_\chi \gg \Gamma_{N_1}$. In terms of the reheating temperature T_R , it is

$$T_R \gg T_{N_1} \simeq 10^6 - 10^7 \text{ GeV}, \quad (2.121)$$

which is easily satisfied in various SUSY inflation models. (Recall that the SUSY inflation models in Sec. 2.3 naturally lead to $T_R \gg 10^6 - 10^7 \text{ GeV}$.) Thus, the present leptogenesis scenario from $\tilde{N}_1$ dominated early universe is almost automatic as long as the right-handed neutrino has suitable mass and couplings given in Eqs. (2.119) and (2.120).

2.4.3 Dilution of the gravitinos

A very attractive feature in the present “ $\tilde{N}_1$ -dominant universe scenario” is the dilution of the gravitino due to the entropy production caused by the $\tilde{N}_1$ decay. We discuss this point in this subsection.

Let us first calculate the dilution factor of the entropy production caused by the $\tilde{N}_1$ ’s decay. It depends on the two cases discussed in the previous subsection, $M_1 < \Gamma_\chi$ [case (I)] and $M_1 > \Gamma_\chi$ [case (II)]. We first consider the latter case, $M_1 > \Gamma_\chi$ [see Fig. 2.19 (II)]. At the time of inflaton decay ($H = \Gamma_\chi$), the ratio of the energy density of $\tilde{N}_1$ to the entropy density is given by

$$\left. \frac{\rho_{N_1}}{s} \right|_{\text{before } \tilde{N}_1 \text{ decay}} = \frac{\rho_{N_1}}{\rho_{\text{rad}}} \times \frac{\rho_{\text{rad}}}{s} \simeq \frac{3}{8} T_R, \quad (2.122)$$

where we have used $\rho_{N_1}/\rho_{\text{rad}} \simeq 1/2$ and $\rho_{\text{rad}}/s = (3/4)T_R$ at $H = \Gamma_\chi$. This ratio takes a constant value until the $\widetilde{N}_1$ decays, since ρ_{N_1} and s are diluted at the same rate R^{-3} as the universe expands. Then at the time of the $\widetilde{N}_1$'s decay, the energy density of the $\widetilde{N}_1$ is converted to the radiation. The ratio of the energy density to the entropy density in the *new* radiation is given by

$$\left. \frac{\rho_{\text{rad}}}{s} \right|_{\text{after } \widetilde{N}_1 \text{ decay}} = \frac{3}{4}T_{N_1}. \quad (2.123)$$

By equating the ρ_{N_1} (just before the $\widetilde{N}_1$'s decay) in Eq. (2.122) with the ρ_{rad} (just after the $\widetilde{N}_1$'s decay) in Eq. (2.123), we obtain the dilution factor of the entropy production:

$$\Delta \equiv \frac{s_{\text{after}}}{s_{\text{before}}} = \frac{T_R}{2T_{N_1}}. \quad (2.124)$$

Next, we consider the case $\Gamma_\chi > M_1$ [case (I)]. In this case, the ratio ρ_{N_1}/s takes a constant value after the $\widetilde{N}_1$ starts its oscillation, i.e., for $H \lesssim M_1$. This ratio is evaluated in the same way as Eq. (2.122):

$$\left. \frac{\rho_{N_1}}{s} \right|_{\text{before } \widetilde{N}_1 \text{ decay}} = \frac{3}{8}T_{M_1} \quad (2.125)$$

where T_{M_1} is the temperature at $H = M_1$ in the case (I), which is given by

$$T_{M_1} \equiv 7 \times 10^{11} \left(\frac{M_1}{10^6 \text{ GeV}} \right)^{1/2} \text{ GeV}. \quad (2.126)$$

Then the dilution factor at $H = \Gamma_{N_1}$ is obtained in the same way as in Eq. (2.124), which results in $\Delta = T_{M_1}/(2T_{N_1})$.

To summarize, the dilution factor Δ is given by

$$\Delta \simeq \begin{cases} \frac{T_R}{2T_{N_1}} & (\text{for } T_R < T_{M_1}) \\ \frac{T_{M_1}}{2T_{N_1}} & (\text{for } T_R > T_{M_1}) \end{cases}, \quad (2.127)$$

where we have used the fact that $\Gamma_\chi < M_1$ ($\Gamma_\chi > M_1$) corresponds to $T_R < T_{M_1}$ ($T_R > T_{M_1}$).

As discussed in Sec. 1.5, the abundance of the gravitino produced at the reheating epoch after the inflation is proportional to the reheating temperature of the inflation T_R . Therefore, there are usually severe upper bounds on T_R depending on the gravitino mass. However, in the present scenario, the number density of the gravitino is diluted thanks to the entropy production caused by the $\widetilde{N}_1$'s decay, which was discussed just

above. Namely, the constraint coming from the gravitino problems should apply, not to the reheating temperature T_R , but to an effective temperature given by

$$T_{R\text{eff}} \equiv \frac{1}{\Delta} T_R \simeq \begin{cases} 2 T_{N_1} \\ 2 T_{N_1} \left(\frac{T_R}{T_{M_1}} \right) \end{cases} \simeq \begin{cases} 2 \times 10^6 - 2 \times 10^7 \text{ GeV} & (\text{for } T_R < T_{M_1}) \\ 2 \times 10^6 - 2 \times 10^7 \text{ GeV} \times \left(\frac{T_R}{T_{M_1}} \right) & (\text{for } T_R > T_{M_1}) \end{cases} \quad (2.128)$$

which is much below the original reheating temperature T_R . Actually, for $T_R < T_{M_1}$, the gravitino abundance which was originally produced at the reheating after primordial inflation is diluted and becomes comparable to the abundance which is produced at the “reheating” after $\widetilde{N}_1$ ’s decay.

In the case of unstable gravitino, Eq. (2.128) means that the cosmological gravitino problem coming from the big-bang nucleosynthesis bounds can be avoided in a wide range of the gravitino mass $m_{3/2} \simeq 100 \text{ GeV} - 10 \text{ TeV}$, even if the reheating temperature T_R of the inflation is much higher than 10^{11} GeV . (Recall that the T_{M_1} is given by Eq. (2.126).) We should stress that the fact that such high reheating temperature is allowed makes it very easy to construct SUSY inflation models (see Sec. 2.3).

In the case of stable gravitino as well, we see from Eq. (2.128) that the overclosure problem can be avoided in a wide range of the gravitino mass. Actually, in our scenario, the relic abundance of the gravitino is obtained by dividing the original abundance in Eq. (1.23) by the dilution factor Δ in Eq. (2.127):

$$\begin{aligned} \Omega_{3/2} h^2 \Big|_{\text{with } \widetilde{N}_1 \text{ decay}} &\simeq \frac{1}{\Delta} \Omega_{3/2} h^2 \Big|_{\text{without } \widetilde{N}_1 \text{ decay}} \\ &\simeq 0.3 \times \left(\frac{m_{\tilde{g}}}{1 \text{ TeV}} \right)^2 \left(\frac{m_{3/2}}{10 \text{ MeV}} \right)^{-1} \left(\frac{T_{R\text{eff}}}{10^6 \text{ GeV}} \right). \end{aligned} \quad (2.129)$$

Therefore, the overclosure problem can be avoided almost independently of the reheating temperature T_R , and a reheating temperature even higher than 10^{11} GeV is possible for $m_{3/2} \gtrsim 10 \text{ MeV}$. Moreover, it is found from this equation that the present energy density of the gravitino is independent of the reheating temperature T_R , in a very wide range of $T_{N_1} \ll T_R < T_{M_1}$. Thus, we can *predict* the gravitino mass by requiring that the gravitino is the dominant component of the dark matter:²⁸

$$m_{3/2} \simeq 20 \text{ MeV} \times \left(\frac{m_{\tilde{g}}}{1 \text{ TeV}} \right)^2 \left(\frac{\Omega_{\text{matter}} h^2}{0.15} \right)^{-1} \left(\frac{T_{R\text{eff}}}{10^6 \text{ GeV}} \right)$$

²⁸Notice that with such a small gravitino mass as in Eq. (2.130), the problem of the decay of the next-to-lightest SUSY particle into LSP gravitino after the big-bang nucleosynthesis is also avoided [94, 48]. See Sec. 1.5.

$$\simeq (40 - 400) \text{ MeV} \times \left(\frac{m_{\tilde{g}}}{1 \text{ TeV}} \right)^2 \left(\frac{\Omega_{\text{matter}} h^2}{0.15} \right)^{-1}, \quad (2.130)$$

for $T_{N_1} \ll T_R < T_{M_1}$. Here, we have taken the present matter density $\Omega_{\text{matter}} \simeq 0.3$ and $h \simeq 0.7$ [29].

Such a light (stable) gravitino as in Eq. (2.130) naturally exists in gauge-mediated SUSY breaking (GMSB) models [45]. The GMSB mechanism has been regarded as a very attractive candidate for the SUSY breaking, since it suppresses quite naturally the flavor changing processes, which are inherent problems in the SUSY standard model. It is remarkable that the gravitino with mass $m_{3/2} \simeq 40\text{--}400$ MeV becomes naturally the dominant component of the dark matter in the present scenario, independently of the reheating temperature T_R for $T_{N_1} \ll T_R < T_{M_1}$. This prediction comes from the fact that the present energy density of the gravitino is determined by the effective temperature $T_{R\text{eff}} = 2T_{N_1}$ (for $T_R < T_{M_1}$), while the decay temperature of the right-handed neutrino T_{N_1} is fixed by the baryon asymmetry in the present universe [see Eq. (2.118)].

2.4.4 Remarks

We give several comments on the present scenario.

2.4.4.a neutrino masses $m_{\nu i}$

The first one is about the neutrino mass m_ν . Let us first rewrite the neutrino masses in terms of the three contributions from N_1 , N_2 and N_3 :

$$m_{\nu j} = m_{\nu j}^{\text{from } N_1} + m_{\nu j}^{\text{from } N_2} + m_{\nu j}^{\text{from } N_3}, \quad (2.131)$$

where

$$m_{\nu j}^{\text{from } N_i} \equiv -\hat{h}_{ij}\hat{h}_{ij}\frac{\langle H_u \rangle^2}{M_i}, \quad (2.132)$$

and the $\hat{h}_{ij}$ denote the Yukawa couplings of the neutrinos in the basis where mass matrices of heavy and light neutrinos are diagonal, which are defined in Eq. (2.12). Then we see that the contribution from N_1 satisfies the following relation:²⁹

$$\begin{aligned} \left| m_{\nu j}^{\text{from } N_1} \right| &= \left| \hat{h}_{1j}\hat{h}_{1j} \right| \frac{\langle H_u \rangle^2}{M_1} \\ &\leq \sum_k |\hat{h}_{1k}|^2 \frac{\langle H_u \rangle^2}{M_1} \end{aligned}$$

²⁹The last expression in Eq. (2.133) is nothing but the $\widetilde{m}_1$ parameter in Eq. (2.28).

$$= \sum_{\alpha} |h_{1\alpha}|^2 \frac{\langle H_u \rangle^2}{M_1} \simeq 8 \times 10^{-4} \text{ eV} \left(\frac{T_{N_1}}{M_1} \right)^2, \quad (2.133)$$

Here, we have used the relation in Eq. (2.120). Therefore, it is understood that the mass scale of the neutrinos suggested from the atmospheric [10] and solar [11] (+KamLAND [12]) neutrino oscillations, $m_{\nu} \sim \mathcal{O}(10^{-3})\text{--}\mathcal{O}(10^{-1})$ eV, should be induced from the heavier right-handed neutrinos, N_2 and N_3 .

The relative hierarchy between the mass and couplings of N_1 and those of the N_2 and N_3 might be naturally explained by a broken flavor symmetry. For example, a broken discrete Z_6 symmetry [95]³⁰ with a breaking parameter $\varepsilon \simeq 1/17$ and charges $Q(L_e, L_{\mu}, L_{\tau}) = (a+1, a, a)$ and $Q(N_1, N_2, N_3) = (3+d, c, b)$ gives rise to the following superpotential;

$$W = \frac{1}{2} \widetilde{\xi}_{11} M_0 \varepsilon^{2d} N_1 N_1 + \frac{1}{2} \sum_{(i,j) \neq (1,1)} \widetilde{\xi}_{ij} M_0 \varepsilon^{Q_i+Q_j} N_i N_j + \widetilde{h}_{i\alpha} \varepsilon^{Q_i+Q_{\alpha}} N_i L_{\alpha} H_u,$$

where $\widetilde{\xi}_{ij}$ and $\widetilde{h}_{i\alpha}$ are $\mathcal{O}(1)$ couplings. The above charge assignments for lepton doublets naturally lead to the realistic neutrino mass matrix including the maximal mixing for the atmospheric neutrino oscillation [76]. (See Sec. 2.3.2.) The overall mass scale of the right-handed neutrino M_0 is determined by $m_{\nu 3} \sim \varepsilon^{2a} \langle H_u \rangle^2 / M_0$. By taking $a+d=2$, this model gives $M_1 \sim \varepsilon^{2d} M_0 \sim 7 \times 10^9$ GeV, $\sqrt{\sum_{\alpha} |h_{1\alpha}|^2} \sim \varepsilon^5 \sim 7 \times 10^{-7}$, and hence $T_{N_1} \sim 1 \times 10^7$ GeV.³¹

Recently, it has been shown [96] that the small mass and the couplings of the N_1 , required in the present scenario, naturally arise if the right-handed neutrino is a bulk field in higher dimensional theory.

2.4.4.b contributions from $\widetilde{N}_2$ and $\widetilde{N}_3$

In the previous subsections, we have only considered the lightest right-handed sneutrino, $\widetilde{N}_1$. The heavier right-handed sneutrino $\widetilde{N}_{2(3)}$ can also develop a large amplitude during the inflation (if $M_{2(3)} < H_{\text{inf}}$) and it may produce lepton asymmetry in a similar way to the $\widetilde{N}_1$. However, it is found that the decay temperatures of the $\widetilde{N}_2$ and $\widetilde{N}_3$ cannot satisfy the out-of-equilibrium condition $T_{2(3)} < M_1$, since N_2 and N_3 must explain the mass scales of the neutrino oscillations. (In other words, if $T_{N_{2(3)}}$ would also satisfy the condition $T_{N_{2(3)}} < M_1 < M_{2(3)}$, the mass scale for the neutrino oscillations could no longer be explained by the contributions from N_2 and N_3 [see Eq. (2.133)].) Therefore, even if the $\widetilde{N}_{2(3)}$'s decay produces additional lepton asymmetry, it is washed out and hence it cannot contribute to the resultant total lepton asymmetry.

³⁰The model here looks similar to the one in Sec. 3.3.1, but different from that.

³¹In terms of $\widetilde{m}_1$ parameter, this corresponds to $\widetilde{m}_1 \sim \varepsilon^6 m_{\nu 3} \sim \mathcal{O}(10^{-9})$ eV.

2.4.4.c thermal effects

We should also comment on the effects of the thermal plasma, which might cause an early oscillation of the right-handed sneutrino $\widetilde{N}_1$ before $H = M_1$. (Notice that there is a dilute plasma with a temperature $T \simeq (T_R^2 M_G H)^{1/4}$ even before the reheating process of the inflation completes [28].) There are basically two possible thermal effects [97, 98]. (Those thermal effects will be also explained in detail in Sec. 3.2.3.)

First, when the temperature T is higher than the effective mass for L and H_u , i.e., $T > m_{\text{eff}} = \sqrt{\sum_\alpha |h_{1\alpha}|^2 |\widetilde{N}_1|}$, the $\widetilde{N}_1$ receives an additional thermal mass $\delta M_{\text{th}}^2 = (1/4) \sum_\alpha |h_{1\alpha}|^2 T^2$ from the Yukawa coupling to L and H_u [97]. Thus, the $\widetilde{N}_1$ field would start an early oscillation if the additional thermal mass becomes larger than the Hubble expansion rate before $H = M_1$. However, even if $\widetilde{N}_1$ receives the thermal mass, the ratio of the thermal mass to the Hubble expansion rate is given by

$$\frac{\delta M_{\text{th}}^2}{H^2} \simeq \begin{cases} 0.07 \times \left(\frac{10T_{N_1}}{M_1}\right)^2 \left(\frac{M_1}{H}\right)^{3/2} \left(\frac{T_R}{T_{M_1}}\right) & \text{for } T_R < T_{M_1}, \\ 0.03 \times \left(\frac{10T_{N_1}}{M_1}\right)^2 \left(\frac{M_1}{H}\right) & \text{for } T_R > T_{M_1}, \end{cases} \quad (2.134)$$

where we have used the relation given in Eq. (2.120). Therefore, we can safely neglect the above thermal effect as long as M_1 is a bit larger than T_{N_1} .

There is also another thermal effect [98]. If the temperature is lower than the effective mass for L and H_u , i.e., $T < m_{\text{eff}} = \sqrt{\sum_\alpha |h_{1\alpha}|^2 |\widetilde{N}_1|}$, the evolution of the running gauge and/or Yukawa coupling constants $f(\mu)$ which couple to them are modified below the scale $\mu = m_{\text{eff}}$. Thus, these running coupling constants depend on $|\widetilde{N}_1|$, and there appears an additional thermal potential for $\widetilde{N}_1$;

$$\delta V(\widetilde{N}_1) = aT^4 \log \left(\frac{|\widetilde{N}_1|^2}{T^2} \right), \quad (2.135)$$

where a is a constant of order $\mathcal{O}(f^4)$. However, again, it turns out that the effective thermal mass for $\widetilde{N}_1$ is less than the Hubble expansion rate;

$$\begin{aligned} \frac{\delta M_{\text{th}}'^2}{H^2} &= \frac{aT^4}{H^2 |\widetilde{N}_1|^2} \\ &\simeq \begin{cases} 0.2 \times a \left(\frac{M_G}{|\widetilde{N}_1|}\right)^2 \left(\frac{M_1}{H}\right) \left(\frac{T_R}{T_{M_1}}\right)^2 & \text{for } T_R < T_{M_1}, \\ 0.05 \times a \left(\frac{M_G}{|\widetilde{N}_1|}\right)^2 & \text{for } T_R > T_{M_1}, \end{cases} \end{aligned} \quad (2.136)$$

and hence this thermal effect is also irrelevant to the present scenario.

2.4.4.d more general cases

So far, we have assumed that the initial amplitude of the $\widetilde{N}_1$'s oscillation is $|\widetilde{N}_{1i}| \simeq M_G$. This can be realized when the right-handed neutrino has only the mass term up to the Planck scale. We have also assumed that the $\widetilde{N}_1$ dominates the energy density of the universe before its decay. Here, we consider more general cases for completeness.

Let us assume that the masses of the right-handed neutrinos are dynamically induced by a spontaneously broken gauge symmetry, for example, by the $U(1)_{B-L}$ symmetry, which is an attractive candidate for such a gauge symmetry as discussed in Sec. 1.1. We denote the chiral superfields whose vacuum expectation values break the $U(1)_{B-L}$ by Ψ and $\overline{\Psi}$. (We need two fields with opposite charges in order to cancel $U(1)_{B-L}$ gauge anomalies.) Due to the D -term and the F -term coming from the superpotential which gives the right-handed neutrino masses, the scalar potential of the right-handed sneutrino $\widetilde{N}_1$ is lifted above the $U(1)_{B-L}$ breaking scale $\langle \Psi \rangle$ [25]. Therefore, the initial amplitude of the $\widetilde{N}_1$'s oscillation at $H \simeq M_1$ is given by $|\widetilde{N}_{1i}| \sim \langle \Psi \rangle$.

The breaking scale of the $U(1)_{B-L}$ gauge symmetry is model dependent. If it is broken at the Planck scale, $\langle \Psi \rangle \simeq M_G$, the discussion so far does not change at all.³² On the other hand, if $\langle \Psi \rangle$ is below the Planck scale, the initial amplitude of the $\widetilde{N}_1$'s oscillation is reduced and its energy density when it starts the oscillation is given by

$$\rho_{N_1} = M_1^2 |\widetilde{N}_{1i}|^2 \ll \rho_{\text{total}} = 3M_1^2 M_G^2 \quad \text{for } H = M_1. \quad (2.137)$$

Compared with the previous case (see Fig. 2.19), we can see that the initial energy density is much suppressed.

In Fig. 2.20, We show the behavior of the energy densities of the inflaton χ , the right-handed sneutrino $\widetilde{N}_1$ and the radiation, for generic cases.³³ Figure (Ia) and (Ib) represent the case of $\Gamma_\chi > M_1$ ($T_R > T_{M_1}$), (IIa) and (IIb) represent the case of $M_1 > \Gamma_\chi > \Gamma_{N_1}$ ($T_{M_1} > T_R > T_{N_1}$) and (III) represents that of $\Gamma_{N_1} > \Gamma_\chi$ ($T_{N_1} > T_R$). Among them, (Ia) and (IIa) show the case of $\widetilde{N}_1$ dominant universe, corresponding to (I) and (II) in Fig. 2.19, respectively. Figures (Ib), (IIb), and (III) represent the cases in which the coherent oscillation of the right-handed sneutrino $\widetilde{N}$ does not dominate the energy density of the universe.

³²In this case, we need small couplings in order to explain the intermediate right-handed neutrino mass scale. For example, a superpotential $W = (1/2)g_i \Psi N_i N_i$ with $\langle \Psi \rangle \simeq M_G$ and $g_3 \sim 10^{-4}$ gives the mass $M_3 \sim 10^{14}$ GeV to the heaviest right-handed neutrino. Such a small Yukawa coupling could well be a consequence of broken flavor symmetries.

³³For the reduced initial amplitude, the thermal effect from the $aT^4 \log(|\widetilde{N}_1|^2)$ potential becomes larger than the case of $|\widetilde{N}_{1i}| \simeq M_G$. However, it is still irrelevant as long as the coupling $a \sim \mathcal{O}(f^4)$ is small enough. [See Eq. (2.136).]




Figure 2.20: Schematic behaviors of the energy densities of the inflaton χ , right-handed sneutrino $\tilde{N}_1$, and radiation for general cases.




Figure 2.21: The condition of the $\tilde{N}_1$ dominant universe is represented in $(\Gamma_\chi, |\tilde{N}_{1i}|)$ plane. The behaviors of the energy densities for the regions of (Ia), (Ib), (IIa), (IIb) and (III) are shown in the corresponding figures in Fig. 2.20.

Let us first discuss the condition of the $\tilde{N}_1$ dominant universe, which was given by [Eq. (2.121)] for $|\tilde{N}_1| \simeq M_G$. Because of the smaller initial amplitude, the condition now becomes

$$|\tilde{N}_{1i}| > \sqrt{3}M_G \left(\frac{\Gamma_{N_1}}{M_1} \right)^{1/4} \quad \text{for } \Gamma_\chi > M_1 \quad \textbf{(Ia)}, \quad (2.138)$$

$$|\tilde{N}_{1i}| > \sqrt{3}M_G \left(\frac{\Gamma_{N_1}}{\Gamma_\chi} \right)^{1/4} \quad \text{for } \Gamma_{N_1} < \Gamma_\chi < M_1 \quad \textbf{(IIa)}. \quad (2.139)$$

These conditions are summarized in Fig. 2.21. Notice that the $\tilde{N}_1$ cannot dominate the universe for $\Gamma_\chi < \Gamma_{N_1}$ [the case (III)].

Next we estimate the baryon asymmetry in the above general cases. It can be calcu-

lated in the same manner as before, and is given by, for each case,

$$\begin{aligned}
\text{(Ia)(IIa)} \quad \frac{n_B}{s} &= 0.35 \times \frac{3}{4} \epsilon_1 \frac{T_{N_1}}{M_1} = 0.5 \times 10^{-10} \left(\frac{T_{N_1}}{10^6 \text{ GeV}} \right) \eta, \\
\text{(Ib)} \quad \frac{n_B}{s} &= 0.35 \times \frac{3}{4} \epsilon_1 \left(\frac{|\widetilde{N}_{1i}|}{\sqrt{3} M_G} \right)^2 \frac{T_{M_1}}{M_1} = 0.5 \times 10^{-10} \left(\frac{T_{M_1}}{10^6 \text{ GeV}} \right) \left(\frac{|\widetilde{N}_{1i}|}{\sqrt{3} M_G} \right)^2 \eta, \\
\text{(IIb)(III)} \quad \frac{n_B}{s} &= 0.35 \times \frac{3}{4} \epsilon_1 \left(\frac{|\widetilde{N}_{1i}|}{\sqrt{3} M_G} \right)^2 \frac{T_R}{M_1} = 0.5 \times 10^{-10} \left(\frac{T_R}{10^6 \text{ GeV}} \right) \left(\frac{|\widetilde{N}_{1i}|}{\sqrt{3} M_G} \right)^2 \eta,
\end{aligned} \tag{2.140}$$

where $\eta \equiv (m_{\nu 3}/0.05 \text{ eV}) \delta_{\text{eff}}$. Notice that the baryon asymmetry which was given in Eq. (2.117) does not change and independent of the initial amplitude $|\widetilde{N}_{1i}|$ as long as the $\widetilde{N}_1$ dominant universe is realized [(Ia) and (IIa)].

Finally, we comment on the gravitino problem. The effective temperature of the cosmological gravitino problem, which was given by [Eq. (2.128)], is now given by

$$\Omega_{3/2} h^2 \propto T_{\text{Reff}} = \begin{cases} T_{N_1} \left(\frac{\sqrt{3} M_G}{|\widetilde{N}_{1i}|} \right)^2 \left(\frac{T_R}{T_{M_1}} \right) & \text{(Ia)}, \\ T_{N_1} \left(\frac{\sqrt{3} M_G}{|\widetilde{N}_{1i}|} \right)^2 & \text{(IIa)}, \\ T_R & \text{(Ib)(IIb)(III)}. \end{cases} \tag{2.141}$$

2.4.4.e cosmic microwave background (CMB) anisotropy from $\widetilde{N}_1$

Let us mention the cosmic density perturbation coming from the fluctuation of the initial amplitude of the right-handed sneutrino, $|\widetilde{N}_{1i}|$, according to the recent study in Ref. [99].

In the present scenario, the CMB radiation we observe today has two origins; the inflaton χ and the right-handed sneutrino $\widetilde{N}_1$. Furthermore, if the Hubble expansion rate of the inflation is larger than the mass of the right-handed sneutrino M_1 , as we have assumed, the initial amplitude of the $\widetilde{N}_1$ has a non-vanishing fluctuation during the inflation. Therefore, the CMB anisotropy also has two origins, the primordial fluctuations of χ and that of $\widetilde{N}_1$. If the $\widetilde{N}_1$ completely dominates the energy density of the universe, the perturbation becomes adiabatic [100]. On the other hand, if the radiation coming from $\widetilde{N}_1$ is negligible compared to that from χ , purely baryonic isocurvature perturbation is produced from $\widetilde{N}_1$. Particularly interesting is between these two cases (i.e., the region near the border line between the two regions (Ia)-(IIa) and (Ib)-(IIb)-(III) in Fig. 2.21), in which correlated mixture of the adiabatic and isocurvature fluctuation is generated, that can be observed in the current and future satellite experiments on the CMB anisotropies [90].

2.A N -dominant universe from inflaton decay

In this appendix, we briefly discuss the case where the N_1 which are produced in inflaton decay dominates the universe before it decays. (This case was also discussed in Ref. [74].)

The baryon asymmetry produced by the N_1 decay in this case is the same as that of the $\widetilde{N}$ -dominant universe (Sec. 2.4), and is given by Eq. (2.117):

$$\frac{n_B}{s} \simeq 0.5 \times 10^{-10} \left(\frac{T_{N_1}}{10^6 \text{ GeV}} \right) \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}, \quad (2.142)$$

which requires the decay temperature of the N_1 to be $T_{N_1} \simeq 10^6\text{--}10^7$ GeV for $\delta_{\text{eff}} \simeq 0.1\text{--}1$.³⁴

On the other hand, the condition for the N_1 to dominate the energy density of the universe in this case is derived as follows. The point is that the ratio n_{N_1}/s , which is given by Eq. (2.42), takes a constant value until the time of N_1 's decay. Thus, the ratio of the energy density of the N_1 to that of the radiation at N_1 's decay is given by

$$\begin{aligned} \frac{\rho_{N_1}}{\rho_{\text{rad}}} &= M_1 \frac{n_{N_1}}{s} \times \frac{s}{\rho_{\text{rad}}} \\ &= M_1 \frac{3}{2} B_r \frac{T_R}{m_\chi} \times \frac{4}{3T_{N_1}} \\ &= 2B_r \frac{M_1}{m_\chi} \frac{T_R}{T_{N_1}}. \end{aligned} \quad (2.143)$$

Thus, the dominant condition $\rho_{N_1}/\rho_{\text{rad}} > 1$ gives a lower bound on the M_1 as follows:

$$M_1 > \frac{m_\chi}{2B_r T_R} T_{N_1}, \quad (2.144)$$

which is stronger than the out-of-equilibrium condition ($M_1 > T_{N_1}$) for $m_\chi > 2B_r T_R$. Then using the relation in Eq. (2.120), we obtain a constraint on the Yukawa couplings:

$$\sum_\alpha |h_{1\alpha}|^2 \lesssim 5 \times 10^{-15} B_r \left(\frac{T_R}{10^9 \text{ GeV}} \right) \left(\frac{10^{13} \text{ GeV}}{m_\chi} \right) \times \left(\frac{T_{N_1}}{10^6 \text{ GeV}} \right). \quad (2.145)$$

If this is satisfied, N_1 produced from inflaton decay can dominate the energy density of the universe, and the produced baryon asymmetry is given by Eq. (2.142).

³⁴This point was not discussed in Ref. [74].

Chapter 3

Leptogenesis via LH_u flat direction

In the previous chapter, we have investigated various leptogenesis mechanisms in which lepton asymmetry is produced by the CP -violating, asymmetric decay of the heavy right-handed (s)neutrino into leptons and anti-leptons. In this chapter,¹ we will discuss a completely different leptogenesis scenario. The main character in this chapter is not a heavy right-handed (s)neutrino, but a flat direction scalar field including the charged lepton doublet L . Actually, as we will see, most of the necessary ingredients for this scenario exist in the minimal supersymmetric standard model (MSSM), and basically it does not rely on any new physics beyond supersymmetry, the existence of small Majorana neutrino masses and inflation.

¹This chapter is based on the works with T. Asaka, M. Fujii and T. Yanagida [101, 95, 102, 103].

3.1 Overview

Affleck and Dine [27] proposed a completely new mechanism to produce cosmological baryon asymmetry, which is inherent in the SUSY framework. In the SUSY theories, there are many flat directions in the scalar potential which contain scalar quark (squark) and scalar lepton (slepton) fields, carrying baryon and/or lepton charges. During the inflation, such a scalar field (which will be denoted by ϕ) can acquire a large vacuum expectation value along the flat direction, since the potential is flat. After the inflationary stage ends, the flat direction field ϕ starts a coherent oscillation. At this stage, if there exists baryon (or lepton) number violating operator in the scalar potential, ϕ field can have a nonzero phase rotational motion: $\frac{d}{dt}[\arg(\phi)] \neq 0$, which means, a nonzero ϕ -number

$$n_\phi \equiv i(\dot{\phi}^* \phi - \phi^* \dot{\phi}) \neq 0, \quad (3.1)$$

is produced. (The dot denotes a derivative with time t .) Since the ϕ field carries baryon (or lepton) number, this represents nothing but baryon (or lepton) number production.

Among various flat directions, the LH_u flat direction proposed by Murayama and Yanagida [25] is especially attractive and had been investigated in detail later as well [104, 105, 101, 95, 102, 103]. The most interesting point is that it is closely related to the neutrino physics. Actually, as we will see, the most important parameter which determines the present baryon asymmetry is the mass of the lightest neutrino, $m_{\nu 1}$. Furthermore, the LH_u flat direction is very special among various flat directions, since it is free from a serious problem of the Q-ball formation.²

Meantime, it has been pointed out [104, 97, 98] that the dynamics of a flat direction field is affected by thermal effects. Although production of baryon and/or lepton number usually takes place before the reheating process of the inflation completes, there is a dilute plasma even in this epoch. This dilute plasma can modify the scalar potential [97, 98], affect the dynamics of ϕ , and change (actually, suppress) the resultant baryon asymmetry.

In Sec. 3.2, therefore, we will perform a detailed analysis on the dynamics of the LH_u flat direction field ϕ , including all the relevant thermal effects, in order to calculate the amount of produced lepton (and hence baryon) asymmetry. We see that evolution of the ϕ field is indeed affected by the thermal effects, and that the produced lepton asymmetry is suppressed by those thermal effects. On the other hand, we will find a very interesting result: the amount of produced baryon asymmetry is almost independent of the reheating temperature T_R , and is determined mainly by the mass of the lightest

²We do not discuss the Q-ball problem in this thesis. For details and references, see, for example, Refs. [106, 102, 107].

neutrino $m_{\nu 1}$. We show that the empirical baryon asymmetry $n_B/s \simeq (0.4-1) \times 10^{-10}$ predicts an ultralight neutrino $m_{\nu 1} \simeq (0.3-1) \times 10^{-9}$ eV for a wide range of reheating temperature $T_R \gtrsim 10^5$ GeV. In Sec. 3.3 we show some explicit models to explain such a small mass of the lightest neutrino.

Then we study in Sec. 3.4 what happens in this leptogenesis scenario via LH_u flat direction when we introduce a gauged $U(1)_{B-L}$ symmetry. We will see that if the initial amplitude of the flat direction field exceeds the $B - L$ breaking scale, we can avoid the suppression of the lepton-asymmetry production due to the thermal effects.

Finally, in Sec. 3.5, we show that if we assume a very small mass of the lightest neutrino as suggested by the present leptogenesis mechanism, the rate of the neutrino-less double-beta decay can be predicted in a high accuracy.

A comment: we do not discuss other lepton-number carrying flat directions (see, e.g., early works in Refs. [108, 57], other general directions in Ref. [104], and a recent one in Ref. [109]) nor the case of multi-scalar manifolds [110]. Notice that the LH_u direction is special not only because it is a leptonic direction, but also because it is directly related to observable physics: the neutrino mass.

3.2 Lepton asymmetry from LH_u flat direction

In this section, we will discuss the dynamics of the LH_u flat direction field in detail and calculate the amount of generated lepton asymmetry.

3.2.1 The LH_u flat direction

Let us start by writing down the effective dimension-five operator in the superpotential

$$W = \frac{m_{\nu i}}{2 \langle H_u \rangle^2} (L_i H_u) (L_i H_u) = \frac{1}{2 M_{\text{eff},i}} (L_i H_u) (L_i H_u) , \quad (3.2)$$

which induces neutrino masses $m_{\nu i}$ after the neutral component of the Higgs field H_u obtains its vacuum expectation value $\langle H_u \rangle = 174 \text{ GeV} \times \sin \beta$. Here, we have taken a basis where the neutrino mass matrix is diagonal. As we will see, the effective mass scales $M_{\text{eff},i}$ play crucial roles in the present scenario. Notice that they are directly related to the neutrino masses $m_{\nu i}$ by the following equation:

$$M_{\text{eff},i} \equiv \frac{\langle H_u \rangle^2}{m_{\nu i}} = 3 \times 10^{13} \text{ GeV} \left(\frac{1 \text{ eV}}{m_{\nu i}} \right) . \quad (3.3)$$

Here and hereafter, we take $\sin \beta \simeq 1$, for simplicity.³

We adopt the following scalar field ϕ_i as the Affleck-Dine flat direction [27] for leptogenesis, which was originally proposed in Ref. [25]:

$$\widetilde{L}_i = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_i \\ 0 \end{pmatrix} , \quad H_u = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \phi_i \end{pmatrix} , \quad (3.4)$$

where the factor $\sqrt{2}$ is introduced to ensure the canonical kinetic term for ϕ_i . Here, we have chosen the neutral components of the $\widetilde{L}_i$ and H_u for illustration, which is always possible by using the $SU(2)_L$ gauge transformation. We also used an $U(1)_Y$ gauge rotation to make their phases to be the same.

There are three flat directions ($i = 1, 2, 3$). (See Appendix 3.A for more details.) Hereafter, we will suppress the family index i for simplicity. As we will see later, it turns out that the relevant flat direction ϕ_i for the most effective leptogenesis corresponds to the first family $i = 1$. Thus, our flat direction field is parameterized by one complex scalar field, $\phi \equiv \phi_1$. We will use the same symbol ϕ to denote the supermultiplet corresponding to the scalar field ϕ , hereafter. Its superpotential is then given by

$$W = \frac{1}{8 M_{\text{eff}}} \phi^4 , \quad (3.5)$$

³This is the case for $\tan \beta > 1$. Even for $\tan \beta \simeq 1$, the final lepton (and hence baryon) asymmetry changes (decreases) by a factor of 2.

while the scalar potential is given by

$$V_0 = m_\phi^2 |\phi|^2 + \frac{m_{3/2}}{8M_{\text{eff}}} (a_m \phi^4 + \text{H.c.}) + \frac{1}{4M_{\text{eff}}} |\phi|^6. \quad (3.6)$$

Here, m_ϕ and $m_{3/2}a_m$ are SUSY-breaking mass parameters,⁴ and the last term directly comes from the superpotential Eq. (3.5). Here and hereafter, we assume that the SUSY breaking is mediated by the gravity. Therefore, it is expected that $m_\phi \sim m_{3/2} \sim 1$ TeV and $|a_m| \sim 1$.

3.2.2 Initial amplitude

The potential for ϕ is D -flat, and it is also F -flat in the limit of $m_\nu \rightarrow 0$. Thus, it is expected that the ϕ field can develop a large amplitude during the inflaton. It depends on the coupling between ϕ and the inflaton χ whether or not the ϕ field can have a large initial amplitude. Suppose that there are general nonminimal couplings of the ϕ field to the inflaton χ in the Kähler potential:

$$\begin{aligned} K &= \phi^\dagger \phi + \chi^\dagger \chi \\ &+ \left(\frac{c_\phi}{M_G} \chi \phi^\dagger \phi + \text{H.c.} \right) + \frac{b_\phi}{M_G^2} \chi^\dagger \chi \phi^\dagger \phi + \cdots, \end{aligned} \quad (3.7)$$

where c_ϕ and b_ϕ are complex and real coupling constants, respectively, and the ellipsis denotes higher order terms which are irrelevant to the following discussion. During inflation, the energy density of the universe is dominated by the inflaton χ . Thus, the total energy density of the universe is given by $3H_{\text{inf}}^2 M_G^2 \simeq |F_\chi|^2$, where H_{inf} is the Hubble parameter during the inflation and F_χ is the F -component of the inflaton supermultiplet χ . Because of this nonzero energy density of χ , there appears additional SUSY-breaking effects during the inflation [104]. Using the scalar potential in the supergravity in Eq. (2.57), we obtain

$$\delta V = 3 \left(1 - b_\phi + |c_\phi|^2 \right) H_{\text{inf}}^2 |\phi|^2 - \frac{\sqrt{3}}{2} \frac{H_{\text{inf}}}{M_{\text{eff}}} (c_\phi \phi^4 + \text{H.c.}) + \cdots, \quad (3.8)$$

where we redefined the phase of c_ϕ to include the phase of F_χ . A crucial observation here is that the ϕ field gets a large mass term of the order of the Hubble parameter. (We will call it a Hubble mass term.) In general, the coefficient of this Hubble mass term, $3(1 - b_\phi + |c_\phi|^2)$, is expected to be order unity. If it is positive, therefore, the ϕ field is driven exponentially towards the origin during the inflation, and the present mechanism

⁴Here, we have included the possible contribution from the μ -term, $W = \mu H_u H_d$, which gives a mass term $(1/2)\mu^2 |\phi|^2$, in the mass m_ϕ^2 .

cannot work. (Notice that this would happen even if the Kähler potential is minimal, i.e., $c_\phi = b_\phi = 0$.) Hence, we will assume that it is negative, and hereafter take

$$\delta V = -c_H H_{\text{inf}}^2 |\phi|^2 + \frac{H_{\text{inf}}}{8M_{\text{eff}}} (a_H \phi^4 + \text{H.c.}) , \quad (3.9)$$

with $c_H \simeq |a_H| \simeq 1$.

The negative Hubble mass term causes an instability of ϕ at the origin. From Eqs. (3.6) and (3.9), the minima of the potential are given by

$$|\phi| \simeq \sqrt{M_{\text{eff}} H_{\text{inf}}} , \quad (3.10)$$

$$\arg(\phi) \simeq \frac{-\arg(a_H) + (2n+1)\pi}{4} , \quad n = 0 \cdots 3 , \quad (3.11)$$

where we have used $m_\phi, m_{3/2}|a_m| \ll H_{\text{inf}}$. Since curvatures of the potential around the minimum along both the radius and phase directions are positive and of the order of the Hubble parameter H_{inf} , the flat direction ϕ runs to one of the four minima from any given initial value and is settled down it.

Here, we comment on another possibility that the flat direction field does not receive a mass term of the order of the Hubble parameter, which occurs with a Kähler potential of the no-scale type supergravity [111]. The evolution of the ϕ field in this case was discussed in Ref. [105], and it turns out that the initial amplitude is likely to be given by Eq. (3.10) in this case as well.

3.2.3 Evolution of the flat direction

Before discussing the detailed dynamics of the ϕ field, we first roughly describe the evolution of ϕ and note the relevant epoch for the leptogenesis. During the inflation, the ϕ field takes a large value $\sim \sqrt{M_{\text{eff}} H_{\text{inf}}}$ as discussed in the previous subsection. After the end of inflation, the value of ϕ gradually decreases as the Hubble parameter H decreases and then, at some time, the ϕ starts its coherent oscillation. As we will see, the net lepton number is fixed when the flat direction field ϕ starts the coherent oscillation.

Hereafter, we will assume that the net lepton number is fixed before the reheating process of the inflation completes, namely, during the epoch when the energy density of the universe is dominated by the oscillating inflaton χ . (This assumption will be justified after we calculate the amount of the generated lepton asymmetry in Sec. 3.2.4.)

3.2.3.a potential for ϕ

Let us first show the total effective potential for the flat direction field ϕ relevant to the leptogenesis:

$$\begin{aligned}
V_{\text{total}} = & \left(m_\phi^2 - H^2 + \sum_{f_k|\phi|<T} c_k f_k^2 T^2 \right) |\phi|^2 \\
& + \frac{m_{3/2}}{8M_{\text{eff}}} (a_m \phi^4 + H.c.) + \frac{H}{8M_{\text{eff}}} (a_H \phi^4 + H.c.) \\
& + a_g \alpha_S(T)^2 T^4 \ln \left(\frac{|\phi|^2}{T^2} \right) \\
& + \frac{1}{4M_{\text{eff}}^2} |\phi|^6.
\end{aligned} \tag{3.12}$$

We explain each term in turn.

First of all, the F -term potential and the SUSY-breaking terms give rise to the scalar potential at vacuum V_0 , which is given by Eq. (3.6).

Next, because the energy density of the universe is still dominated by the inflaton χ during the inflaton-oscillation epoch, the ϕ field receives additional SUSY-breaking terms caused by the finite energy density of χ :

$$\delta V_H = -c_H H^2 |\phi|^2 + \frac{H}{8M_{\text{eff}}} (a_H \phi^4 + \text{H.c.}) . \tag{3.13}$$

This can be derived in the same way as Eq. (3.9). Recall that we have taken $c_H \simeq |a_H| \simeq 1$. We will call the second term Hubble A -term.

Finally, the rests in Eq. (3.12) correspond to the thermal effects which we discuss now. The crucial point here is that the decay of the inflaton χ occurs during the coherent oscillation of χ , while it completes much later than the beginning of the oscillation. Thus, even in the χ oscillation period, there is a dilute plasma consisting of the decay products of the inflaton χ , although most of the energy density of the universe is carried by the coherent oscillation of χ . The temperature of this dilute plasma is given by [28]

$$T \simeq (T_R^2 M_G H)^{1/4} . \tag{3.14}$$

Notice that in this dilute plasma the temperature decreases as $T \propto H^{1/4}$, rather than as $T \propto H^{1/2}$ like in the usual radiation dominated universe.

This dilute plasma has crucial effects on the dynamics of the ϕ field [104, 97, 98]. There are basically two possible thermal effects. First, the fields ψ_k which couple to ϕ are

produced by the inflaton decay and/or by thermal scatterings if their effective masses are less than the temperature,

$$m_{\text{eff},k} = f_k |\phi| < T, \quad (3.15)$$

and hence they must be included in the plasma. Here, f_k denote the Yukawa or gauge coupling constants between the flat direction ϕ and ψ_k (we take f_k real and positive). Therefore, the flat direction ϕ receives a thermal mass of order $\sim f_k T$, if the condition Eq. (3.15) is satisfied. The induced mass term is given by [97]

$$\delta V_1^{\text{th}} = \sum_{f_k |\phi| < T} c_k f_k^2 T^2 |\phi|^2, \quad (3.16)$$

where c_k are real positive constants of order unity (see below). The summation means that only the fields in thermal plasma [i.e., ψ_k whose couplings f_k satisfy the condition Eq. (3.15)] can induce the thermal mass for ϕ .

We include all the couplings relevant for the flat direction $H_u = \widetilde{L}_1 = \phi/\sqrt{2}$ in the SUSY standard model, i.e., the gauge couplings for the $\text{SU}(2)_L$ and $\text{U}(1)_Y$ gauge groups, and Yukawa couplings for up-type quarks and charged leptons. The couplings f_k , which are redefined so that the effective masses for the fields ψ_k become $m_{\text{eff},k} = f_k |\phi|$, are listed in Table. 3.1 with the coefficients c_k of the thermal mass for ϕ . Among the gauge supermultiplets, the flat direction ϕ does not couple to one $\text{U}(1)$ group which remains unbroken by the condensation of ϕ . Thus, there are one Z -like and two W -like massive gauge multiplets as in the SUSY standard model. For the couplings of the charged leptons, we should be careful. The flat direction ϕ receives its thermal mass from only one linear combination of charged leptons. The effective Yukawa coupling y_{L1} is determined by the Yukawa couplings for the charged leptons and the mixing matrix among gauge eigenstates of the neutrinos. (Note that the relevant ϕ is the flattest direction which corresponds to the first family ν_1 of mass eigenstates.) Taking large mixing angles both for atmospheric and solar neutrino oscillations suggested by the recent data [10, 11, 12], we find $y_{L1} \simeq \mathcal{O}(0.1-1) \times y_\tau$. We will see in Sec. 3.2.4 that the Yukawa coupling for the up quark induces a relevant thermal effect, while other couplings turn out to be too large to satisfy the condition Eq. (3.15) in most of the parameter space.

Next, there is another thermal effect which was pointed out in Ref. [98]. Along the LH_u flat direction, the $\text{SU}(3)_C$ gauge symmetry remains unbroken and hence gluons and gluinos are massless. Furthermore, the down type (s)quarks also remain massless since they have no direct coupling to the ϕ field. These massless fields produce the free energy which depends on the $\text{SU}(3)_C$ gauge coupling constant g_S , and we obtain an effective potential $\delta V \propto g_S(T)^2 T^4$ at two loop level.⁵ At first sight, there seems no dependence on

⁵For the coefficient of this free energy, see, for example, Ref. [112].

f_k (coupling for ϕ)	$\sqrt{\frac{g_1^2 + g_2^2}{2}}$	$\frac{g_2}{\sqrt{2}}$	$\frac{y_{u,a}}{\sqrt{2}}$ ($a = \text{up, charm, top}$)	$\frac{y_{L1}}{\sqrt{2}}$
c_k	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{3}{4}$	$\frac{1}{4}$

Table 3.1: The couplings of the flat direction field ϕ to other fields and the coefficients c_k of the thermal mass of ϕ induced by these fields. g_1 and g_2 denote the gauge couplings for the $U(1)_Y$ and $SU(2)_L$ gauge groups, respectively, and $y_{u,a}$ ($a = \text{up, charm, top}$) and y_{L1} are the Yukawa couplings for up-type quarks and charged lepton, respectively.

the ϕ field in this free energy. However, we can see this is not the case by means of the following arguments.

The evolution of the running coupling g_S is given by

$$\frac{d}{d \ln \mu} g_S(\mu) = \frac{g_S^3}{16\pi^2} \left(-3C_2 + \sum_{m_i < \mu} C(R_i) \right), \quad (3.17)$$

where $C_2 = 3$ and $C(R_i) = 1/2$ for fundamental representations R_i . Notice that the evolution changes due to the decoupling when the scale μ passes through a mass of a field m_i . Since the up type (s)quarks get large masses from the couplings to the ϕ field $f_{u,a} = y_{u,a}/\sqrt{2}$ ($a = \text{up, charm, top}$), they change the trajectory of the running at $\mu = m_{u,a} = f_{u,a}|\phi|$, as shown in Fig. 3.1. Hence, if $f_{u,a}|\phi| > T$, the running coupling $g_S(T)$ depends on the value of the ϕ field:

$$g_S(T) = g_S(T)|_{\phi=0} + \frac{g_S(M_U)^3}{32\pi^2} \sum_{f_{u,a}|\phi| > T} C(R_{u,a}) \ln \left(\frac{f_{u,a}^2 |\phi|^2}{T^2} \right), \quad (3.18)$$

where M_U is the ultraviolet scale where g_S is fixed. Then, there appears an additional potential through the modification of the gauge coupling constant in Eq. (3.18)

$$\delta V_2^{\text{th}} = a_g \alpha_S(T)^2 T^4 \left(\frac{2}{3} \sum_{f_{u,a}|\phi| > T} C(R_{u,a}) \right) \ln \left(\frac{|\phi|^2}{T^2} \right), \quad (3.19)$$

where a_g is a constant which is a bit larger than unity⁶ and $\alpha_S(T) \equiv g_S(T)_{\phi=0}^2/(4\pi)$. (We will omit the subscript $\phi = 0$ for simplicity.) Hereafter, we take the factor $(2/3) \sum C(R)$ to be unity since it does not change the result much.⁷

⁶By using the free energy at two-loop order given in Ref. [112], a_g is given by $a_g = 1.125$ in the case of the LH_u flat direction. The author thanks Masaaki Fujii for helpful discussion.

⁷At least the top Yukawa coupling $f_{u,\text{top}}$ always satisfies $f_{u,\text{top}}|\phi| > T$ before the oscillation of ϕ .




Figure 3.1: A schematic behavior of the $SU(3)_C$ gauge coupling g_S . The dashed line represents the running coupling when the ϕ field does not have a vacuum-expectation value.

3.2.3.b evolution of ϕ

Now we eventually obtain the total effective potential Eq. (3.12):

$$V_{\text{total}} = V_0 + \delta V_H + \delta V_1^{\text{th}} + \delta V_2^{\text{th}}. \quad (3.20)$$

The evolution of ϕ is described by the equation of motion with this V_{total} as

$$\ddot{\phi} + 3H\dot{\phi} + \frac{\partial V_{\text{total}}}{\partial \phi^*} = 0, \quad (3.21)$$

where the dot denotes a derivative with time. During the inflation, there is no thermal plasma and hence no thermal potentials. The vacuum expectation value of the ϕ field at this stage is given by Eqs. (3.10) and (3.11). Let us call the valley to which the ϕ field rolls down $n = n_0$.

After the inflation ends, the inflaton χ starts to oscillate and its decay produces a dilute plasma. However, the potential is still dominated by Hubble-induced terms [in Eq. (3.13)] and $|\phi|^6$ term [in Eq. (3.6)] at the first stage of the oscillation. Thus, the flat direction field ϕ is trapped in the following minimum:

$$|\phi| \simeq \sqrt{M_{\text{eff}} H}, \quad (3.22)$$

$$\arg(\phi) \simeq \frac{-\arg(a_H) + (2n_0 + 1)\pi}{4}, \quad (3.23)$$

Thus, the resultant baryon asymmetry changes by only a factor of $\sqrt{3}$ at most. [See Eqs. (3.34) and (3.41).]

This is because the curvatures around the minimum along both radius and phase directions are of the order of H also in this period, and hence the ϕ field always catches up the instantaneous minimum [104].

Then, as the Hubble parameter decreases, the negative Hubble-induced mass term is eventually exceeded by one of the following three terms in the potential:⁸

$$H^2 \lesssim m_\phi^2 + \sum_{f_k|\phi|<T} c_k f_k^2 T^2 + a_g \alpha_S^2(T) \frac{T^4}{|\phi|^2}. \quad (3.24)$$

As we show now, it is this time when the oscillation of ϕ starts. Let us denote the Hubble parameter at this time by H_{osc} . The evolution of the ϕ after $H \simeq H_{\text{osc}}$ depends on which term in Eq. (3.12) dominates the effective potential. There are basically three cases; the potential is dominated by (i) $m_\phi^2|\phi|^2$ term, (ii) $T^2|\phi|^2$ term, or (iii) $T^4 \ln(|\phi|^2)$ term.

(i) First, if the potential is dominated by the $m_\phi^2|\phi|^2$ term, the equation of motion Eq. (3.21) is given by

$$\ddot{\phi} + 3H\dot{\phi} + m_\phi^2\phi \simeq 0. \quad (3.25)$$

It is clear in this case that the field ϕ oscillates around the origin ($\phi = 0$) and the amplitude of the oscillation damps as $|\phi| \propto H \propto t^{-1}$.

(ii) Second, when the potential is dominated by the thermal mass term $c_k f_k^2 T^2 |\phi|^2$, the equation of motion becomes

$$\ddot{\phi} + 3H\dot{\phi} + c_k f_k^2 T^2 \phi \simeq 0. \quad (3.26)$$

This equation can be solved analytically and the solution is given by

$$\begin{aligned} \phi(t) &= \phi_1 \left[\frac{1}{z^{2/3}} J_{2/3}(z) \right] + \phi_2 \left[\frac{1}{z^{2/3}} J_{-2/3}(z) \right] \\ z &= \frac{4}{3} \left(\frac{2}{3} c_k^2 f_k^4 M_G T_R^2 t^3 \right)^{1/4}, \end{aligned} \quad (3.27)$$

where ϕ_1 and ϕ_2 are constants and J_ν is the Bessel function. We see that the ϕ oscillates around the origin in this case as well. The time scale of the oscillation is $\sim (f_k^4 M_G T_R^2)^{-1/3}$ and the amplitude of the oscillation is damped as $|\phi| \sim t^{-7/8} \sim H^{7/8}$.

(iii) The third case is given when the $a_g \alpha_S^2 T^4 \ln(|\phi|^2/T^2)$ term dominates the potential. If we neglect the time dependence of T^4 , the damping rate of the oscillation amplitude due to such a flat potential, $V \sim \ln(|\phi|^2)$, can be estimated by using the virial theorem, and it

⁸Here, we have compared $(1/\phi)(\partial V/\partial\phi^*)$ instead of the potential terms themselves, since the evolution of ϕ is governed by the equation of motion Eq. (3.21).

is given by $|\phi| \propto H^2 \propto t^{-2}$ [113]. In the actual case, however, the potential itself gradually decreases with time as $T^4 \propto t^{-1}$. We have numerically checked that the amplitude damps as $|\phi| \propto H^\alpha \propto t^{-\alpha}$ with $\alpha \simeq 1.5$.

Thus, in any case, the ϕ field starts a coherent oscillation. Notice that, in all the above cases, the damping rate is faster than the rate before the beginning of the ϕ 's oscillation ($|\phi| \propto H^{1/2} \propto t^{-1/2}$).

Now let us estimate the cosmic time t_{osc} when the ϕ field starts its oscillation, or equivalently the Hubble parameter at that time, $H_{\text{osc}} = (2/3)t_{\text{osc}}^{-1}$. As we will see in Sec. 3.2.4, this oscillation time plays an important role to determine the final lepton asymmetry. H_{osc} is determined by the term which exceeds the Hubble mass term at first. (Therefore, again, there are three cases.)

(i) If the soft mass term dominates the potential at first, H_{osc} is just given by $H_{\text{osc}} = m_\phi$.

(ii) The case when the thermal mass term dominates at first is a bit complicated. Let us first consider a simple case in which there is only one field ψ_1 with a coupling f_1 . From Eqs. (3.15) and (3.24), it is found that the following two conditions must be satisfied in order to cause the ϕ 's oscillation by the thermal mass term:

$$H < \frac{1}{f_1^4} \frac{M_G T_R^2}{M_{\text{eff}}^2}, \quad (3.28)$$

$$H < \left(c_1^2 f_1^4 M_G T_R^2 \right)^{1/3}, \quad (3.29)$$

where we have used the relations $|\phi| \simeq \sqrt{M_{\text{eff}} H}$ and $T = (T_R^2 M_G H)^{1/4}$. The meanings of these conditions are as follows. If the coupling f_1 is small enough, the ψ_1 can easily enter in the thermal plasma ($f_1 |\phi| < T$), but the induced thermal mass for ϕ is smaller than the Hubble-induced mass H at the beginning ($c_1 f_1^2 T^2 < H^2$). However, since the temperature decreases more slowly than the Hubble parameter H ($T \propto H^{1/4}$) as the universe expands, the thermal mass eventually exceeds H ($c_1 f_1^2 T^2 > H^2$). Then the ϕ 's oscillation starts at the time when the condition Eq. (3.29) is satisfied. On the other hand, if f_1 is very large, the would-be thermal mass can be large enough to exceed the Hubble mass ($c_1 f_1^2 T^2 > H^2$). However, the thermal mass does not appear until the $|\phi|$ becomes small enough to satisfy $f_1 |\phi| < T$. In this case the ϕ starts to oscillate at the time when the condition Eq. (3.28) is satisfied. Therefore, we find the Hubble parameter H_{osc} to be

$$H_{\text{osc}} = \min \left[\frac{1}{f_1^4} \frac{M_G T_R^2}{M_{\text{eff}}^2}, \left(c_1^2 f_1^4 M_G T_R^2 \right)^{1/3} \right]. \quad (3.30)$$

It is easy to apply the above discussion to the case of more than one couplings. If there

is another coupling f_i which can satisfy the both conditions Eqs. (3.28) and (3.29) earlier than f_1 , the flat direction ϕ starts its oscillation earlier. Therefore, the Hubble parameter H_{osc} is given by⁹

$$H_{\text{osc}} = \max_i (H_i) , \quad (3.31)$$

where

$$H_i = \min \left[\frac{1}{f_i^4} \frac{M_G T_R^2}{M_{\text{eff}}^2} , \left(c_i^2 f_i^4 M_G T_R^2 \right)^{1/3} \right] . \quad (3.32)$$

(iii) The third case is again given when the $a_g \alpha_s^2 T^4 \ln(|\phi|^2/T^2)$ term dominates the potential at first. In this case, it is found from Eq. (3.24) that the oscillation time of the ϕ field is given by¹⁰

$$H_{\text{osc}} = \alpha_S T_R \left(\frac{a_g M_G}{M_{\text{eff}}} \right)^{1/2} . \quad (3.33)$$

After all, the Hubble parameter at the time when the ϕ field starts its coherent oscillation is given by

$$H_{\text{osc}} \simeq \max \left[m_\phi , H_i , \alpha_S T_R \left(\frac{a_g M_G}{M_{\text{eff}}} \right)^{1/2} \right] , \quad (3.34)$$

where H_i is given by Eq. (3.32). We stress here that the thermal effects always make the oscillation time earlier. (Notice that if there is no thermal effect, the oscillation time is just given by $H_{\text{osc}} = m_\phi$.) As we shall show soon, this “early oscillation” drastically affect the amount of the produced lepton asymmetry.

3.2.4 Lepton asymmetry

Now we are at the point to calculate the lepton asymmetry produced by the ϕ field. Since the ϕ field carries lepton charge, its number density is related to the lepton number density n_L as

$$n_L = \frac{1}{2} i \left(\dot{\phi}^* \phi - \phi^* \dot{\phi} \right) . \quad (3.35)$$

From Eqs. (3.12), (3.21) and (3.35), the evolution of n_L is described by the following equation:

$$\dot{n}_L + 3H n_L = \frac{m_{3/2}}{2M_{\text{eff}}} \text{Im} \left(a_m \phi^4 \right) + \frac{H}{2M_{\text{eff}}} \text{Im} \left(a_H \phi^4 \right) . \quad (3.36)$$

⁹We assume here hierarchical couplings and neglect effects of the summation.

¹⁰The running coupling $\alpha_S(T)$ here is obtained by solving the equation $H^2 = a_g \alpha_S(T) T^4 / |\phi|^2$ iteratively.

The production of the lepton asymmetry occurs as follows. Suppose that the original A -term would vanish, i.e., $m_{3/2} a_m = 0$. If this is the case, the flat direction ϕ is always trapped in one of the valleys induced by the Hubble A -term [see Eq. (3.23)] during the both periods $t < t_{\text{osc}}$ and $t > t_{\text{osc}}$, and the direction of the valley does not change with time. Therefore, there is no force which causes the motion of ϕ along the phase direction and no lepton-number asymmetry is produced (i.e., the right-hand side of Eq. (3.36) vanishes.) However, we have the original A -term, and the phase of ϕ is kicked by the relative phase difference between a_m and a_H in Eq. (3.12). The phase of ϕ changes during its rolling towards the origin, because the Hubble parameter H decreases and the direction of the true valleys changes with time. Therefore, the original A -term, which corresponds to the first term in Eq. (3.36), plays a role of the source of the lepton asymmetry.

One might wonder if the Hubble A -term, which corresponds to the second term in Eq. (3.36), gives a larger contribution to the lepton asymmetry since $H \gg m_{3/2}$. However, since the ϕ almost traces one of the valleys determined mainly by the Hubble A -term, $\text{Im}(a_H \phi^4)$ in Eq. (3.36) is highly suppressed compared with $\text{Im}(a_m \phi^4)$. In fact, we have found numerically that the contribution of the second term in Eq. (3.36) is always comparable or less than that from the first term. Thus, we neglect the lepton asymmetry produced from the Hubble A -term in our analytic calculation, for simplicity. By integrating Eq. (3.36), we obtain the resultant lepton number at the time t ,

$$\left[R^3 n_L \right] (t) \simeq \frac{m_{3/2}}{2M_{\text{eff}}} \int^t dt' R^3 \text{Im} \left(a_m \phi^4 \right), \quad (3.37)$$

where R denotes the scale factor of the expanding universe, which scales as $R^3 \propto H^{-2} \propto t^2$ in the universe dominated by the oscillation energy of the inflaton χ . We can see that the total lepton number increases with time as $R^3 n_L \propto t$ until the oscillation of ϕ starts ($t < t_{\text{osc}}$), since $\phi^4 \propto H^2$ and hence $R^3 \phi^4 \sim \text{const}$ in this stage.

On the other hand, after the ϕ starts its oscillation, the production of lepton number is strongly suppressed. This is because $\text{Im}(a_m \phi^4)$ changes its sign rapidly due to the oscillation of ϕ , and also because the amplitude of ϕ 's oscillation is damped as $R^3 \phi^4 \sim t^{-n}$ with $n > 1$. [See discussion around Eqs. (3.25)–(3.27).] Therefore, the net lepton asymmetry is fixed when the oscillation of ϕ starts. The generated lepton number at this epoch ($t = t_{\text{osc}}$) is given approximately by

$$\begin{aligned} n_L(t_{\text{osc}}) &= \frac{m_{3/2}}{2M_{\text{eff}}} \text{Im} \left[a_m \phi^4(t_{\text{osc}}) \right] t_{\text{osc}} \\ &\simeq \frac{1}{3} \left(m_{3/2} |a_m| \right) M_{\text{eff}} H_{\text{osc}} \delta_{\text{ph}}, \end{aligned} \quad (3.38)$$

where $\delta_{\text{ph}} \simeq \sin(4 \arg \phi + \arg a_m)$ represents an effective CP violating phase. Here, we have used $\phi(t_{\text{osc}}) \simeq \sqrt{M_{\text{eff}} H_{\text{osc}}}$ and $t_{\text{osc}} = (2/3) H_{\text{osc}}^{-1}$. Recall that we have assumed that the

production of net lepton asymmetry occurs before the reheating process of the inflation completes, i.e., $H_{\text{osc}} > \Gamma_\chi$ where Γ_χ is the decay rate of the inflaton χ . Thus, the lepton number when the reheating process completes ($t = t_R$, $H \simeq \Gamma_\chi$) is given by

$$\begin{aligned} n_L(t_R) &= n_L(t_{\text{osc}}) \left(\frac{R(t_{\text{osc}})}{R(t_R)} \right)^3 \\ &= n_L(t_{\text{osc}}) \left(\frac{\Gamma_\chi}{H_{\text{osc}}} \right)^2. \end{aligned} \quad (3.39)$$

Then, the lepton-to-entropy ratio is estimated as

$$\frac{n_L}{s} = \frac{M_{\text{eff}} T_R}{12 M_G^2} \left(\frac{m_{3/2} |a_m|}{H_{\text{osc}}} \right) \delta_{\text{ph}}, \quad (3.40)$$

when the reheating process of inflation completes. Here, we have used $3M_G^2 \Gamma_\chi^2 = \rho_{\text{rad}}(t_R) = (3/4)T_R \times s(t_R)$.

As explained in Sec. 1.4, this lepton asymmetry is partially converted into the baryon asymmetry through the sphaleron effect. Thus, after all, the present baryon asymmetry is given by

$$\begin{aligned} \frac{n_B}{s} &\simeq 0.35 \times \frac{n_L}{s} \\ &= 0.029 \times \frac{M_{\text{eff}} T_R}{M_G^2} \left(\frac{m_{3/2} |a_m|}{H_{\text{osc}}} \right) \delta_{\text{ph}}. \end{aligned} \quad (3.41)$$

From Eq. (3.41), we see that the baryon asymmetry is a monotonically increasing function of the scale M_{eff} . [Notice that the H_{osc} depends on M_{eff} as shown in Eq. (3.34).] Since M_{eff} is directly related to the neutrino mass m_ν as $M_{\text{eff}} = \langle H_u \rangle^2 / m_\nu$, this means that the baryon asymmetry becomes larger as the neutrino mass m_ν decreases. Therefore, as mentioned at the beginning of this chapter, the most effective flat direction corresponds to the lightest neutrino ν_1 , i.e., the first family field $\phi/\sqrt{2} = L_1 = H_u$.

The factor H_{osc}^{-1} in Eq. (3.41) represents the suppression of the lepton asymmetry due to the thermal effects. When the ϕ 's oscillation is caused by the thermal effects, i.e., $H_{\text{osc}} > m_\phi$, the production of lepton asymmetry stops earlier than the case without thermal effects, and hence the resultant baryon asymmetry becomes suppressed.

We should also note that the amount of the produced baryon asymmetry is proportional to the gravitino mass, $m_{3/2}$. This is because the force which rotates the ϕ field along the phase direction comes from the SUSY-breaking A -term $(m_{3/2}/8M_{\text{eff}})(a_m \phi^4 + \text{H.c.})$. Thus, the resultant baryon asymmetry would be very much suppressed if the gravitino




Figure 3.2: The contour plot of the baryon asymmetries n_B/s in the $m_{\nu 1}$ - T_R plane. The lines represent the contour plots for $n_B/s = 10^{-9}$, 10^{-10} , 0.4×10^{-10} , 10^{-11} , and 10^{-12} from the left to the right.

is light, like in gauge-mediated SUSY breaking models [45].¹¹ This is the reason why we have assumed gravity-mediated SUSY-breaking and $m_{3/2} \simeq 1$ TeV.

Fig. 3.2 shows the contour plot of the produced baryon asymmetry in the $m_{\nu 1}$ - T_R plane, given by the analytic formula Eq. (3.41). Here, we have taken $m_\phi = m_{3/2} = 1$ TeV, $|a_m| = 1$ and $\delta_{\text{ph}} = 1$, and we have used the relation $m_{\nu 1} = \langle H_u \rangle^2 / M_{\text{eff}}$. (Although the big-bang nucleosynthesis constraint on the gravitino abundance requires $T_R \lesssim 10^9$ GeV for $m_{3/2} = 1$ TeV, we have plotted the contour in Fig. 3.2 up to $T_R \leq 10^{11}$ GeV, keeping in mind that even $T_R \simeq 10^{11}$ GeV can be allowed for a slightly heavier gravitino, say, $m_{3/2} \sim 3$ TeV. See Sec. 1.5.)

A remarkable observation here is that the present baryon asymmetry n_B/s is determined almost independently of the reheating temperature for a wide range of $T_R \gtrsim 10^5$ GeV. In particular, for a relatively high reheating temperature $T_R \gtrsim 10^8$ GeV, the baryon asym-

¹¹If we introduce a gauged $U(1)_{B-L}$, however, the LH_u flat direction can produce enough lepton asymmetry even with a light gravitino. See Sec. 3.4.

metry derived from the Eqs. (3.34) and (3.41) is given by the following simple form;

$$\frac{n_B}{s} \simeq 10^{-11} \delta_{\text{eff}} \times \left(\frac{m_{\nu 1}}{10^{-8} \text{ eV}} \right)^{-3/2} \left(\frac{m_{3/2} |a_m|}{1 \text{ TeV}} \right). \quad (3.42)$$

The reason why it is independent of the reheating temperature T_R is that the oscillation time H_{osc} is determined by the thermal potential $T^4 \ln(|\phi|^2)$ in the higher temperature regime and becomes proportional to the reheating temperature T_R . [See Eq. (3.34).] Thus, the T_R dependence is canceled out in Eq. (3.41).¹² Even in the lower reheating temperature region $10^5 \lesssim T_R \lesssim 10^8 \text{ GeV}$, where H_{osc} is determined by the thermal-mass term potential $T^2 |\phi|^2$, T_R dependence is still mild, i.e., $n_B/s \propto T_R^{1/3}$. The “reheating-temperature independence” discussed here is a very attractive feature of the present mechanism since the produced baryon asymmetry crucially depends on T_R in many other baryogenesis scenarios.

For the analytic calculation in Fig. 3.2, we have taken $\delta_{\text{ph}} = 1$. It is expected that $\delta_{\text{ph}} \sim \mathcal{O}(1)$, say, $\delta_{\text{ph}} \simeq 0.1$ – 1 , unless there is an unnatural cancellation between $\arg(a_m)$ and $\arg(a_H)$. Thus, it is found from Fig. 3.2 that the present baryon asymmetry in our universe $n_B/s \simeq (0.4$ – $1) \times 10^{-10}$ suggests an ultralight neutrino of a mass,

$$m_{\nu 1} \simeq (0.1 - 3) \times 10^{-9} \text{ eV}, \quad (3.43)$$

for $T_R \gtrsim 10^5 \text{ GeV}$ and $\delta_{\text{ph}} \simeq 0.1$ – 1 .

• remarks

Before closing this section, we give several comments. First, let us justify our assumption that the production of the lepton asymmetry stops before the reheating process of the inflation completes, namely, $H_{\text{osc}} > \Gamma_\chi$. We can see this condition is satisfied in all the parameter region in Fig. 3.2 by using the fact that H_{osc} is bounded from below, as shown in Eq. (3.34). For example, it is enough to show $\alpha_S T_R (a_g M_G / M_{\text{eff}})^{1/2} > \Gamma_\chi$. By using $\Gamma_\chi = (\pi^2 g_*/90)^{1/2} T_R^2 / M_G$, it is found that this condition is indeed satisfied for $T_R \lesssim 10^{14} \text{ GeV} \times (m_{\nu 1} / 10^{-9} \text{ eV})^{1/2}$.

Next, let us investigate the behavior of Fig. 3.2 in more detail. Fig. 3.3 shows the dependence of the baryon asymmetry on the Yukawa coupling of the up quark. Notice that the up-quark Yukawa coupling is given by $y_{u,\text{up}} = m_{\text{up}} / \langle H_u \rangle = m_{\text{up}} / (174 \text{ GeV} \times \sin \beta)$. In the previous figure 3.2, we have taken $m_{\text{up}} = 3 \text{ MeV}$ (and $\tan \beta = 10$). In Fig. 3.3, we have taken $m_{\text{up}} = 1 \text{ MeV}$ and $m_{\text{up}} = 5 \text{ MeV}$ for comparison. It is found that the final

¹²Precisely speaking, H_{osc} depends on T_R through the running coupling $\alpha_S(T)$ even in this case. However, as can be seen in Fig. 3.2, this dependence is negligibly mild.




Figure 3.3: The contour plot of the baryon asymmetries n_B/s in the $m_{\nu 1}$ - T_R plane. The lines represent the contour plots for $n_B/s = 10^{-9}$, 10^{-10} , 10^{-11} , and 10^{-12} from the left to the right. The solid lines are the same as those in Fig. 3.2 ($m_{\text{up}} = 3$ MeV). The left (right) dashed lines correspond to $m_{\text{up}} = 5(1)$ MeV.

baryon asymmetry changes at most one order of magnitude. However, the dependence on the reheating temperature is still mild, and hence our conclusion does not change much.

Meantime, we can also see from Fig. 3.3 that the up-quark Yukawa coupling is the only coupling which can give a dominant contribution to the early oscillation of ϕ , since below this “up-quark Yukawa” region the oscillation is caused by the soft mass term m_ϕ (where $n_B/s \propto T_R m_{\nu 1}^{-1}$), while in the higher T_R region the oscillation is caused by the thermal logarithmic term $T^4 \ln(|\phi|^2)$ (where $n_B/s \propto m_{\nu 1}^{-3/2}$). This is because other couplings f_k are too large and the fields with those couplings cannot satisfy the condition $f_k |\phi| < T$.

We have also solved the equation of motion (3.21) numerically with the full scalar potential Eq. (3.12). The result is shown in Fig. 3.4, where the final baryon asymmetry $n_B/s \simeq 0.35 \times n_L/s$ is plotted. (n_L is given by Eq. (3.35), $n_L = (i/2)(\dot{\phi}^* \phi - \phi^* \dot{\phi})$.) Here, we have taken $m_\phi = m_{3/2} |a_m| = 1$ TeV, $\arg(a_m) = \pi/3$ and $\arg(a_H) = 0$. As for the initial condition, we have taken $H_{\text{inf}} = 10^{12}$ GeV, and put the ϕ field at the minimum of the potential during the inflation. From Fig. 3.4 we confirm that the analytic estimation

discussed above reproduces very well the result obtained by the numerical calculation.

Finally, we comment on the possibility to avoid the early oscillation caused by the thermal effects discussed in this section, which has been recently pointed out in Ref. [114]. The point is that, if the scale of the inflation is very low, the Hubble parameter during the inflation H_{inf} can be below the H_{osc} which is given by Eq. (3.34). If $H_{\text{inf}} < H_{\text{osc}}$, our (implicit) assumption that the Hubble parameter of the universe gradually decreases from H_{inf} to H_{osc} breaks down. What happens in this case is that the flat direction field ϕ starts its oscillation just after the inflation ends, i.e., at $H \simeq H_{\text{inf}}$. Thus lepton asymmetry is also fixed at this time. The final baryon asymmetry is then given by replacing H_{osc} in Eq. (3.41) with H_{inf} :

$$\left. \frac{n_B}{s} \right|_{H_{\text{inf}} < H_{\text{osc}}} \simeq 0.029 \times \frac{M_{\text{eff}} T_R}{M_G^2} \left(\frac{m_{3/2} |a_m|}{H_{\text{inf}}} \right) \delta_{\text{ph}}. \quad (3.44)$$

It was shown [114] that $H_{\text{inf}} \lesssim H_{\text{osc}}$ is indeed satisfied for very low H_{inf} and high enough reheating temperature T_R , e.g, $H_{\text{inf}} \simeq 10^5$ GeV and $T_R \gtrsim 10^7$ – 10^8 GeV. For example, in an extreme case where $H_{\text{inf}} \simeq m_\phi$ and $T_R \simeq (\pi^2 g_*/90)^{-1/4} \sqrt{M_G H_{\text{inf}}}$ (sudden reheating), baryon asymmetry is given by [114]

$$\frac{n_B}{s} \simeq 0.3 \times 10^{-10} \left(\frac{10^{-4} \text{ eV}}{m_{\nu 1}} \right) \left(\frac{m_{3/2}}{H_{\text{inf}}} \right)^{1/2} \left(\frac{m_{3/2}}{1 \text{ TeV}} \right)^{1/2} |a_m| \delta_{\text{ph}}. \quad (3.45)$$

Thus the empirical baryon asymmetry can be explained even with $m_{\nu 1} \simeq 10^{-4}$ eV in this extreme case.




Figure 3.4: The plots of the baryon asymmetries n_B/s in the $m_{\nu 1}$ - T_R plane, which are obtained by numerically solving the equation of motion Eq. (3.21). The regions with points show the result of the numerical calculation. They represent $10^{-9} > n_B/s > 10^{-10}$, $10^{-10} > n_B/s > 10^{-11}$ and $10^{-11} > n_B/s > 10^{-12}$ from the left to the right. The solid lines are the same as those in Fig. 3.2, which represent the contour plots for $n_B/s = 10^{-9}$, 10^{-10} , 10^{-11} , and 10^{-12} from the left to the right, given by the analytic formula Eq. (3.41).

3.3 Some models

In Sec. 3.2.4 we have shown that the baryon asymmetry in the present universe predicts the mass for the lightest neutrino in a narrow region, $m_{\nu 1} \simeq (0.1\text{--}3) \times 10^{-9}$ eV. Together with the neutrino masses required to solve the solar [11] and atmospheric [10] neutrino anomalies, this suggests a very large mass hierarchy between the lightest and the heavier two neutrinos.

Although the present leptogenesis mechanism via LH_u direction does not rely on the origin of the effective operator Eq. (3.2), it is natural to consider the seesaw mechanism [13] as the origin of that operator. In this case, the effective mass scale M_{eff} is related to the masses of the right-handed neutrinos M_i . Then, one might wonder if the discussion so far is reliable, since successful leptogenesis requires the effective mass scale M_{eff} to be far beyond the Planck scale M_G , i.e., $M_{\text{eff}} = \langle H_u \rangle^2 / m_{\nu 1} \simeq 3 \times 10^{22} \text{ GeV} \times (m_{\nu 1} / 10^{-9} \text{ eV})^{-1}$. Here, we first comment on this point. In the framework of the seesaw mechanism, the neutrino masses are given by Eq. (2.2):

$$m_{\nu i} = - \sum_k \hat{h}_{ki} \hat{h}_{ki} \frac{\langle H_u \rangle^2}{M_k}, \quad (3.46)$$

where we have rotated the Yukawa coupling so that the neutrino mass matrix becomes diagonal [see Sec. 2.1 and Eq. (2.12)]. Then, from Eqs. (3.3) and (3.46) we obtain

$$M_{\text{eff}} \equiv M_{\text{eff},1} = - \left(\sum_i \frac{\hat{h}_{i1} \hat{h}_{i1}}{M_i} \right)^{-1}. \quad (3.47)$$

Therefore, the large mass scale M_{eff} (or equivalently small neutrino mass $m_{\nu 1}$) can be obtained by taking the Yukawa couplings $\hat{h}_{i1}$ to be small, keeping the masses of the right-handed neutrinos M_i to be below the Planck scale.

Nonetheless, the reader might still consider that the mass scale M_{eff} is too large. In other words, the mass of the lightest neutrino $m_{\nu 1} \simeq (0.1\text{--}3) \times 10^{-9}$ eV might seem to be too small, compared with the masses of the two heavier neutrinos, $m_\nu \sim \mathcal{O}(10^{-3})\text{--}\mathcal{O}(10^{-1})$ eV suggested from the recent neutrino oscillation experiments [10, 11, 12]. In this section, we propose some models in which such an ultralight neutrino (or large mass scale M_{eff}) can be naturally implemented.

3.3.1 Model for the ultralight neutrino with a broken discrete symmetry

Here, we show an explicit model based on a Froggatt-Nielsen (FN) mechanism [68], in which the required large mass hierarchy is naturally obtained. We adopt a discrete Z_6 as

the FN symmetry instead of a continuous $U(1)_{\text{FN}}$. We see that the discrete symmetry is crucial to produce the required large mass hierarchy in the neutrino sector. To see this, let us first recall the case of the continuous $U(1)_{\text{FN}}$, which is described in Sec. 2.3.2. In that case, the mass matrix of the neutrino in Eq. (2.48) gives rise to a mild hierarchy

$$m_{\nu 1} : m_{\nu 2} : m_{\nu 3} \sim \varepsilon^2 : 1 : 1 = \mathcal{O}(10^{-2}) : \mathcal{O}(1) : \mathcal{O}(1), \quad (3.48)$$

and hence it cannot explain the large hierarchy required in the present scenario. To change the above point, we suppose that the broken FN symmetry is not a $U(1)_{\text{FN}}$ but a discrete symmetry Z_n with $n = 2d$. Then, the mass matrix for the right-handed neutrino M_R changes into the following form;

$$M_R = M_0 \begin{pmatrix} \widetilde{\xi}_{11} & \widetilde{\xi}_{12} \varepsilon^{c+d} & \widetilde{\xi}_{13} \varepsilon^{b+d} \\ \widetilde{\xi}_{12} \varepsilon^{c+d} & \widetilde{\xi}_{22} \varepsilon^{2c} & \widetilde{\xi}_{23} \varepsilon^{b+c} \\ \widetilde{\xi}_{13} \varepsilon^{b+d} & \widetilde{\xi}_{23} \varepsilon^{b+c} & \widetilde{\xi}_{33} \varepsilon^{2b} \end{pmatrix}. \quad (3.49)$$

Here, we have assumed $d(= n/2) > c \geq b \geq 0$. Notice that the Majorana mass for N_1 is no longer suppressed by the power of ε , which is a basic point to yield an extremely small neutrino mass $m_{\nu 1}$. Although the mass matrix M_R is modified, the structure of the neutrino mass matrix looks similar to the original one:

$$\begin{aligned} m_\nu &= m_D^T \frac{1}{M_R} m_D \\ &= \frac{\varepsilon^{2a} \langle H_u \rangle^2}{M_0} \begin{pmatrix} \varepsilon & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} (\{\widetilde{y}_{ij}\})^T \begin{pmatrix} \widetilde{\xi}_{11} \varepsilon^{-2d} & \widetilde{\xi}_{12} & \widetilde{\xi}_{13} \\ \widetilde{\xi}_{12} & \widetilde{\xi}_{22} & \widetilde{\xi}_{23} \\ \widetilde{\xi}_{13} & \widetilde{\xi}_{23} & \widetilde{\xi}_{33} \end{pmatrix}^{-1} (\{\widetilde{y}_{ij}\}) \begin{pmatrix} \varepsilon & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \frac{\varepsilon^{2a} \langle H_u \rangle^2}{M_0} \begin{pmatrix} \mathcal{O}(\varepsilon^2) & \mathcal{O}(\varepsilon) & \mathcal{O}(\varepsilon) \\ \mathcal{O}(\varepsilon) & \mathcal{O}(1) & \mathcal{O}(1) \\ \mathcal{O}(\varepsilon) & \mathcal{O}(1) & \mathcal{O}(1) \end{pmatrix}. \end{aligned} \quad (3.50)$$

However, in spite of the large components of the mass matrix, we see that one of the mass eigenvalue of this mass matrix strongly suppressed as $\sim \varepsilon^{2(1+d)}$. (This suppression is also understood directly by taking the determinant of the above mass matrix.) Thus, the required large mass hierarchy between $m_{\nu 1}$ and $m_{\nu 2,3}$ can be naturally obtained by taking $d = 3$ (Z_6).

Notice that the neutrino mass matrix Eq. (3.50) can still lead to a large $\nu_\mu - \nu_\tau$ mixing angle. The mass hierarchy of the charged leptons also remains unchanged.¹³ Hence, this model preserves all the desirable features in the continuous $U(1)_{\text{FN}}$ model, explaining the mass of the ultralight neutrino $m_{\nu 1}$ as well.

¹³This is because the charges of the Yukawa couplings of the charged leptons are smaller than $n = 6$, $Q(\overline{E}_i) + Q(L_j) \leq (a + 3) < 6$.




Figure 3.5: The plot for $r = \delta m_{\text{sol}}^2 / \delta m_{\text{atm}}^2$ and $m_{\nu 1}$ in a Froggatt-Nielsen model with a discrete Z_6 symmetry.

To demonstrate our point, we randomly generate $\mathcal{O}(1)$ couplings $\widetilde{y}_{ij}$ and $\widetilde{\xi}_{ij}$. Namely we calculate the mass matrix for neutrinos, taking the magnitudes of the couplings $\widetilde{y}_{ij}$ and $\widetilde{\xi}_{ij}$ to be in a range $0.5 - 1.5$ and their phases to be $0 - 2\pi$. We also take $\varepsilon = 0.05 - 0.1$ randomly.¹⁴ From the obtained neutrino mass matrix, we calculated the mass eigenstates $m_{\nu i}$ and the mixing matrix $U_{\alpha i}$,¹⁵ and then required the parameters $\delta m_{\text{sol}}^2 \equiv m_{\nu 3}^2 - m_{\nu 2}^2$, $\delta m_{\text{atm}}^2 = m_{\nu 2}^2 - m_{\nu 1}^2$, $\sin^2 2\theta_{\text{atm}} \equiv 4|U_{\mu 3}|^2(1 - |U_{\mu 3}|^2)$, and $\tan^2 \theta_{\text{sol}} \equiv |U_{e 2}/U_{e 1}|^2$ to be consistent with the data of neutrino-oscillation experiments [10, 11, 12]. Here, we take large-angle MSW solution to the solar neutrino deficits, since it is the only solution allowed by the recently announced KamLAND results [12], and adopt the following constraints [10, 115]:

$$\begin{aligned}
0.9 &\leq \sin^2 2\theta_{\text{atm}} \leq 1.0, \\
r &\equiv \delta m_{\text{sol}}^2 / \delta m_{\text{atm}}^2 = 0.008 - 0.15 \\
\tan^2 \theta_{\text{sol}} &= 0.2 - 0.9.
\end{aligned} \tag{3.51}$$

We also required

$$|U_{e 3}| < 0.15, \tag{3.52}$$

to satisfy CHOOZ limit [116]. Fig. 3.5 shows the obtained mass of the lightest neutrino, $m_{\nu 1}$. We can see that an ultralight neutrino of mass $m_{\nu 1} \simeq (0.1-3) \times 10^{-9}$ eV is naturally obtained.

¹⁴A similar calculation was done in Ref. [77], where they adopted the continuous $U(1)_{\text{FN}}$ model.

¹⁵The matrix $U_{\alpha i}$ is defined in Sec. 2.1.

3.3.2 Large effective masses for right-handed neutrinos

In Sec. 3.3.1, we have shown a model in which an ultralight neutrino with mass $m_{\nu 1} \simeq (0.1-3) \times 10^{-9}$ can be naturally implemented. Here, we discuss another possibility. The point is that the large effective mass scale M_{eff} is required *only during the leptogenesis*, not necessarily at the true vacuum. Therefore, if the effective masses of the right-handed neutrinos in the early universe are dynamical values and they are different from the values in the true vacuum, we might naturally obtain a large effective mass scale M_{eff} during the leptogenesis.

Suppose that the right-handed neutrino masses M_i are given by a vacuum expectation value of some field X with a superpotential

$$W = \frac{1}{2} \xi_i X N_i N_i, \quad (3.53)$$

which gives $M_i = \xi_i \langle X \rangle$. If $\langle X \rangle$ takes a large value during the leptogenesis, therefore, M_i can be much larger than the values in the true vacuum. To demonstrate this point, let us consider that the field X is responsible to the Peccei-Quinn symmetry breaking [117]. Then, as we will see, there automatically exists a flat direction containing the X field. The X field can have a large value as $X \simeq M_G$ during the inflation whereas it has a vacuum expectation value of $\langle X \rangle \simeq F_a$ in the true vacuum. Here, the Peccei-Quinn breaking scale F_a is constrained by laboratory experiments, astrophysics, and cosmology as $F_a \simeq 10^{10}-10^{12}$ GeV [118].

We consider the following superpotential for the Peccei-Quinn symmetry breaking sector:

$$W = \lambda Y (X \overline{X} - F_a^2), \quad (3.54)$$

where λ is a coupling constant, and Y , X , and $\overline{X}$ are supermultiplets which are singlets under the standard-model gauge groups, which have 0, +1, and -1 Peccei-Quinn charges, respectively. From Eq. (3.54) we see that there exists a flat direction $X \overline{X} = F_a^2$. We parameterize this direction by the scalar field σ , which is called “saxion”. This saxion σ receives a soft SUSY breaking mass of $m_\sigma \simeq m_{3/2}$, and the X and $\overline{X}$ have vacuum expectation values of the order of F_a .¹⁶

Along this flat direction the X field can have a large value during the inflation, if the additional SUSY breaking effects during the inflation induce a negative mass squared for X (see Sec. 3.2.2). The initial value of X is then expected to be $X \simeq M_G$, since the

¹⁶We assume here that the soft SUSY breaking masses for X and $\overline{X}$ are almost the same, i.e., $m_X \simeq m_{\overline{X}}$ ($\simeq m_{3/2}$).

supergravity effects prevent the X field from running over M_G [see Eq. (2.57)], while the initial value of the $\bar{X}$ field is given by $\bar{X} \simeq F_a^2/M_G$.

In order to fix the mass scale M_{eff} at the true vacuum, let us take the mass of the lightest neutrino to be $m_{\nu 1} = 1 \times 10^{-4}$ eV for a representative value, with which the hierarchy of the neutrino masses becomes mild. Then the scale M_{eff} is given by

$$M_{\text{eff}} \simeq 3 \times 10^{17} \text{ GeV} \left(\frac{10^{-4} \text{ eV}}{m_{\nu 1}} \right) \left(\frac{\langle X \rangle}{F_a} \right). \quad (3.55)$$

Here, we have used $M_{\text{eff}} \propto \langle X \rangle$ and $M_{\text{eff}} = \langle H_u \rangle^2 / m_{\nu 1}$ at the true vacuum $\langle X \rangle \simeq F_a$. Although this scale M_{eff} at the true vacuum is too low to produce the enough baryon asymmetry (see Fig. 3.2), the large initial value of X , $X_0 \simeq M_G$ gives rise to

$$M_{\text{eff}} = 7 \times 10^{25} \text{ GeV} \left(\frac{10^{-4} \text{ eV}}{m_{\nu 1}} \right) \left(\frac{10^{10} \text{ GeV}}{F_a} \right) \left(\frac{X_0}{M_G} \right). \quad (3.56)$$

For such a high scale M_{eff} , the evolution of the ϕ field is completely free from the thermal effects and the early oscillation can be avoided, even if the reheating temperature T_R is as high as $T_R \simeq 10^8$ GeV.¹⁷

The baryon asymmetry is then obtained by using Eqs. (3.41) and (3.56) as

$$\frac{n_B}{s} = 4 \times 10^{-5} \delta_{\text{ph}} \left(\frac{T_R}{10^8 \text{ GeV}} \right) \left(\frac{10^{-4} \text{ eV}}{m_{\nu 1}} \right) \left(\frac{10^{10} \text{ GeV}}{F_a} \right) \left(\frac{X_0}{M_G} \right), \quad (3.57)$$

where we have used $H_{\text{osc}} \simeq m_\phi \simeq m_{3/2}|a_m|$. This shows that too large amount of the lepton asymmetry is produced. However, this asymmetry is sufficiently diluted, since there exists substantial entropy production by the decay of the saxion σ .

The saxion begins the coherent oscillation with the initial amplitude $X_0 \simeq M_G$ at $H \simeq m_\sigma \simeq m_{3/2}$, just after the production of the lepton asymmetry ends. The oscillation energy at that time is given by $\rho_\sigma \simeq m_\sigma^2 X_0^2 / 2$. When the reheating process of the inflation completes, the ratio of ρ_σ to the energy density ρ_{rad} of the radiation of the universe is estimated as

$$\frac{\rho_\sigma}{\rho_{\text{rad}}} \simeq \frac{1}{6} \left(\frac{X_0}{M_G} \right)^2. \quad (3.58)$$

Notice that ρ_σ decreases at the rate R^{-3} as the universe expands, while ρ_{rad} decreases as R^{-4} . Therefore, the oscillation energy of the saxion dominates the energy of the universe soon after the reheating process completes. Here, it is useful to take the ratio of ρ_σ to the entropy density s ,

$$\frac{\rho_\sigma}{s} \simeq \frac{1}{8} T_R \left(\frac{X_0}{M_G} \right)^2, \quad (3.59)$$

¹⁷This can be seen by calculating the oscillation time H_{osc} in Eq. (3.34) with $M_{\text{eff}} \simeq 10^{26}$ GeV.

which takes a constant value until the saxion decays.

This energy density of the saxion is transferred into the thermal bath when it decays. Here, it should be noted that the saxion might decay dominantly into two axions. If it is the case, the extra energy of the axion at the BBN epoch raises the Hubble expansion of the universe, which leads to overproduction of ${}^4\text{He}$. To avoid this difficulty, the branching ratio of the saxion decay into two axion should be smaller than about 0.1. Here, we simply assume this is the case.¹⁸ Then, the saxion decays dominantly into two gluons with the decay rate

$$\Gamma(\sigma \rightarrow 2g) = \frac{\alpha_s^2}{32\pi^3} \frac{m_\sigma^3}{F_a^2}. \quad (3.60)$$

Through this decay the universe is “reheated” again, and its reheating temperature T_σ is estimated as

$$T_\sigma = 10 \text{ GeV} \left(\frac{m_\sigma}{1 \text{ TeV}} \right)^{3/2} \left(\frac{10^{10} \text{ GeV}}{F_a} \right). \quad (3.61)$$

Notice that the saxion decay takes place far before the beginning of the Big-Bang Nucleosynthesis, and hence is cosmologically harmless.

The saxion decay increases the entropy of the universe by the rate

$$\begin{aligned} \Delta \equiv \frac{s_{\text{after}}}{s_{\text{before}}} &= \frac{\rho_\sigma}{s_{\text{before}}} \times \frac{s_{\text{after}}}{\rho_\sigma} \\ &= \frac{1}{8} T_R \left(\frac{X_0}{M_G} \right)^2 \times \frac{4}{3T_\sigma} \\ &= 2 \times 10^6 \left(\frac{T_R}{10^8 \text{ GeV}} \right) \left(\frac{1 \text{ TeV}}{m_\sigma} \right)^{3/2} \left(\frac{F_a}{10^{10} \text{ GeV}} \right) \left(\frac{X_0}{M_G} \right)^2. \end{aligned} \quad (3.62)$$

Because of this entropy production by the saxion decay, the primordial baryon asymmetry Eq. (3.57) is also diluted by the rate Δ . Then the present baryon asymmetry is given by

$$\frac{n_B}{s} = 0.2 \times 10^{-10} \left(\frac{10^{-4} \text{ eV}}{m_{\nu 1}} \right) \left(\frac{10^{10} \text{ GeV}}{F_a} \right)^2 \left(\frac{m_\sigma}{1 \text{ TeV}} \right)^{3/2} \left(\frac{M_G}{X_0} \right) \delta_{\text{ph}}. \quad (3.63)$$

Notice that the present baryon asymmetry is independent of the reheating temperature T_R , while it depends on the Peccei-Quinn scale F_a . We see that the desired baryon asymmetry is obtained even with the lightest neutrino mass of $m_{\nu 1} \simeq 10^{-4} \text{ eV}$ for $F_a \simeq 10^{10} \text{ GeV}$.

Finally, we comment on the cosmological consequences of this model.

¹⁸ This is realized when $m_X^2 \simeq m_{\overline{X}}^2$.

(i) The entropy production by the saxion in Eq. (3.62) ensures that we are free from the cosmological gravitino problems, since the number density of gravitinos produced at the reheating process is diluted by the rate Δ .

(ii) This model does not suffer from the cosmological problem of the “axino”, which is a fermionic superpartner of the axion. The axinos are produced in the reheating process by the thermal scatterings and may lead to a cosmological difficulty [119]. However, in this model, the interaction of axino at the reheating epoch is suppressed by $X_0 \simeq M_G$, not by F_a , and hence the production of axino is less effective. Furthermore, the entropy production by the saxion decay dilutes the axino abundance. Thus, the axino becomes completely cosmologically harmless in this model.

(iii) The reheating temperature of the saxion decay Eq. (3.61) is high enough to thermalize the lightest SUSY particles (LSPs) if the mass of the LSP is less than 200 GeV, and hence the stable LSP can be a dominant component of the dark matter.

3.4 Effects of the gauged $U(1)_{B-L}$ symmetry

In Sec. 3.2, we found a very interesting aspect of the leptogenesis via LH_u flat direction, the “reheating temperature independence of the cosmological baryon asymmetry.” The resultant baryon asymmetry is almost determined by the mass of the lightest neutrino (and the gravitino mass), and hence we can predict the mass of the lightest neutrino $m_{\nu 1}$ as $m_{\nu 1} \simeq 10^{-9}$ eV from the observed baryon asymmetry.

In this section, we discuss the effects of gauging the $U(1)_{B-L}$ symmetry on the leptogenesis via the LH_u flat direction. As mentioned in Sec. 1.1, the absence of $U(1)_{B-L}$ gauge anomaly automatically requires three families of right-handed neutrinos, and the breaking of $B - L$ symmetry naturally gives rise to the heavy Majorana masses of these right-handed neutrinos, which explains the tiny neutrino masses via the seesaw mechanism.

Below the $B - L$ breaking scale, the behavior of the scalar potential of the ϕ field does not change much from the one discussed in Sec. 3.2. Therefore, if the $B - L$ breaking scale v is larger than the initial value of ϕ field obtained in Sec. 3.2.2,

$$v \gtrsim \sqrt{M_{\text{eff}} H_{\text{inf}}}, \quad (3.64)$$

the amplitude of the ϕ field is always smaller than the $B - L$ breaking scale, and the dynamics of the ϕ field is basically the same as the one discussed in Sec. 3.2. Hence, the resultant baryon asymmetry reduces to the one obtained in Sec. 3.2.4 [see Eq. (3.41) and Fig. 3.4]. We will call this case “ F -term stopping case.”

On the other hand, if $v \lesssim \sqrt{M_{\text{eff}} H_{\text{inf}}}$, the ϕ field develops its initial value as large as the $B - L$ breaking scale. At this scale, the potential of the flat direction ϕ can be lifted by the effect of the $U(1)_{B-L}$ D -term [106]. (We denote this case by “ D -term stopping case.”) As we shall show in this section, the dynamics of the ϕ field is drastically changed in this D -term stopping case. Actually, the obtained baryon asymmetry is much larger than the case without a $U(1)_{B-L}$ symmetry, and hence than the F -term stopping case.

It is interesting that the resultant baryon asymmetry in the D -term stopping case linearly depends on the reheating temperatures of inflation, but it is completely independent of the gravitino mass. This is totally an opposite situation to the result obtained in Sec. 3.2, in which the final baryon asymmetry is almost independent of the reheating temperatures of inflation and linearly depends on the gravitino mass [see Eq. (3.41) and Fig. 3.4]. This “gravitino-mass independence of the cosmological baryon asymmetry” provides us a great advantage in gauge-mediated SUSY breaking scenarios [45].

3.4.1 The model

In order to discuss the effect of the $U(1)_{B-L}$ gauge symmetry, we should investigate the $B - L$ breaking sector, which gives the masses of the right-handed neutrinos. To demonstrate our point, we use the following superpotential:

$$W = hNLH_u + \frac{1}{2}\xi SNN + \lambda X (S\bar{S} - v^2) + \mu H_u H_d, \quad (3.65)$$

where, h , ξ , λ are coupling constants and we assume $\lambda = \mathcal{O}(1)$. X , S and $\bar{S}$ are singlets under the MSSM gauge groups and they carry the $U(1)_{B-L}$ charges of 0, -2 and $+2$, respectively. v is the the breaking scale of the $U(1)_{B-L}$ symmetry. Here and hereafter, we take the couplings h , ξ , λ and μ , v to be real, by field redefinitions. We will omit the family indices, since the “effectively flat” direction relevant for the present baryon asymmetry is again given by the flattest flat direction, $L = L_1$.

The shape of the superpotential in Eq. (3.65) is the same as that of the SUSY hybrid inflation discussed in Sec. 2.3.3.a.¹⁹ Actually, If the X field in Eq. (3.65) develops a large expectation value and the energy density of the universe is dominated by the vacuum energy $V \simeq \lambda^2 v^4$, hybrid inflation takes place. Here, however, we assume that the $B - L$ breaking sector in Eq. (3.65) is not directly related to the inflation, and discuss the effects of this $U(1)_{B-L}$ on the dynamics of the LH_u flat direction.

As we have shown in Sec. 3.2.2, the Kähler potential must have non-minimal couplings of the ϕ field to the inflaton χ , since otherwise the ϕ field gets a large positive mass term of the order of the Hubble parameter and it is driven exponentially towards the origin during inflation [104]. We assume that there are also non-minimal couplings of other fields to the inflaton in the Kähler potential:

$$\begin{aligned} K &= \sum_Y Y^\dagger Y + \chi^\dagger \chi \\ &+ \sum_Y \left(\frac{c_Y}{M_G} \chi Y^\dagger Y + \text{H.c.} \right) + \sum_Y \frac{b_Y}{M_G^2} \chi^\dagger \chi Y^\dagger Y + \dots, \end{aligned} \quad (3.66)$$

where c_Y and b_Y are complex and real couplings of order unity, respectively, and Y denotes L , H_u , H_d , X , S , $\bar{S}$, and N . Then, the full scalar potential relevant to the flat direction

¹⁹The superpotential in Eq. (3.65) is also similar to the one in the model in Sec. 3.3.2. However, the model adopted here is different from that, since the $U(1)$ symmetry here is gauged, while in Sec. 3.3.2 the Peccei-Quinn $U(1)$ symmetry is global.

field $L = H_u = \phi/\sqrt{2}$ is given by ²⁰

$$\begin{aligned}
V = & \frac{1}{2}g^2 \left(-2|S|^2 + 2|\bar{S}|^2 + |N|^2 - \frac{1}{2}|\phi|^2 \right)^2 \\
& + \frac{1}{2}\mu^2|\phi|^2 + \left| \frac{1}{2}h\phi^2 + \xi SN \right|^2 + h^2|N\phi|^2 \\
& + \left| \frac{1}{2}\xi N^2 + \lambda X \bar{S} \right|^2 + \lambda^2 |S\bar{S} - v^2|^2 + \lambda^2 |XS|^2 \\
& + 3H^2 \left(\sum_Y (1 - b'_Y) |Y|^2 \right) - \sqrt{3}H \left(\sum_Y c_Y Y W_Y + \text{H.c.} \right) + V_{\text{SB}}, \quad (3.67)
\end{aligned}$$

where $b'_Y \equiv b_Y - |c_Y|^2$ and $W_Y \equiv \partial W / \partial Y$, and we redefined the phase of c_Y . $g = \mathcal{O}(1)$ is the gauge coupling constant of the $U(1)_{B-L}$. V_{SB} represents SUSY breaking terms in the true vacuum. All fields denote the scalar components of the corresponding superfields. Here, we omit the potential coming from thermal effects, which will be discussed in Sec. 3.4.4.

The scalar potential in Eq. (3.67) is so complicated that it seems very difficult to solve the dynamics of the relevant fields. Fortunately, however, we only need to know the shape around the minimum of this potential, if the curvatures around the bottom of the potential are as large as the Hubble parameter during inflation. This is because all the scalar fields, which have masses as large as the Hubble parameter, settle down at the bottom of the potential during inflation and trace this potential minimum in the subsequent evolution. Therefore, we first find out the relations to minimize the potential in Eq. (3.67). Though this is also a very hard task because of the complexity of the potential, one can find out the minimum of the potential at least when $|\phi| \lesssim v$ and $H \lesssim v$ are satisfied. From Eq. (3.67), we find the approximate minimum of F -terms for $|\phi|$, $H \lesssim v$,

$$\bar{S} \simeq \frac{v^2}{S}, \quad X \simeq 0, \quad N \simeq -\frac{h\phi^2}{2\xi S}. \quad (3.68)$$

Notice that the curvatures associated with these three relations are of the order of the $B - L$ breaking scale v , and hence these relations hold as far as $H \lesssim v$. By using these relations, we can reduce the scalar potential in Eq. (3.67) to the following one:

$$\begin{aligned}
V \simeq & \frac{1}{2}g^2 \left(-2|S|^2 + 2\frac{v^4}{|S|^2} + \frac{h^2}{4\xi^2} \frac{|\phi|^4}{|S|^2} - \frac{1}{2}|\phi|^2 \right)^2 + \frac{h^4}{4\xi^2} \frac{|\phi|^6}{|S|^2} + \frac{h^4}{64\xi^2} \frac{|\phi|^8}{|S|^4} \\
& - H^2|\phi|^2 + 3(1 - b'_S)H^2|S|^2 + 3(1 - b'_S)H^2\frac{v^4}{|S|^2} + 3(1 - b'_N)H^2\frac{h^2}{4\xi^2} \frac{|\phi|^4}{|S|^2}
\end{aligned}$$

²⁰Here, we take $H_d \simeq 0$ assuming a positive Hubble-order mass term for the H_d field. This is necessary in order to avoid a contamination of H_d to the LH_u flat direction. We have checked both analytically and numerically that the contamination of the H_d field becomes relevant only after the $H \lesssim \mu$, and hence the following discussion is not affected much by the H_d contamination.

$$+ \frac{\sqrt{3}}{2} H \left(c'_\phi \frac{h^2 \phi^4}{\xi S} + \text{H.c.} \right) + \frac{1}{2} \mu^2 |\phi|^2 + V_{\text{SB}}, \quad (3.69)$$

where $c'_\phi \equiv c_\phi - (1/4)c_S$. Here and hereafter, we assume $3(1 - b'_\phi) \simeq -1$ for simplicity. (This corresponds to $c_H \simeq 1$ in the previous analysis. See Eq. (3.13).)

One may wonder why the “flat” direction field ϕ can develop a large expectation value in spite of the presence of the $U(1)_{B-L}$ D -term. This is because the S field shifts and absorbs the D -term potential:

$$|S|^2 \simeq -\frac{1}{8} |\phi|^2 + \sqrt{v^4 + \left(\frac{1}{64} + \frac{h^2}{8\xi^2} \right) |\phi|^4}. \quad (3.70)$$

The curvature associated with this relation is also of the order of the $B - L$ breaking scale. Then we get the following effective potential:

$$\begin{aligned} V \simeq & V_{\text{SB}} + \frac{1}{2} \mu^2 |\phi|^2 + \frac{\sqrt{3}}{2} H \left(c'_\phi \frac{h^2 \phi^4}{\xi S} + \text{H.c.} \right) + \frac{h^4}{4\xi^2} \frac{|\phi|^6}{|S|^2} + \mathcal{O} \left(\frac{|\phi|^8}{|S|^4} \right) \\ & - H^2 |\phi|^2 + 3(1 - b'_S) H^2 |S|^2 + 3(1 - b'_S) H^2 \frac{v^4}{|S|^2} + 3(1 - b'_N) H^2 \frac{h^2}{4\xi^2} \frac{|\phi|^4}{|S|^2}. \end{aligned} \quad (3.71)$$

In the following discussion, we assume $H \lesssim v$ and that the relations in Eq. (3.68) and in Eq. (3.70) are satisfied during the leptogenesis works. Although the approximation we have made here is crude, we will justify by a numerical calculation that the effective potential in Eq. (3.71) is valid at least for the following discussion.

As we will see later, the third term in Eq. (3.71),

$$V_{\text{phase}} = \frac{\sqrt{3}}{2} H \left(c'_\phi \frac{h^2 \phi^4}{\xi S} + \text{H.c.} \right), \quad (3.72)$$

which depends on the phase of ϕ , plays a crucial role in the D -term stopping case. We will call this term “Hubble A -term.” (This corresponds to the Hubble A -term in Eq. (3.13) in the previous analysis.)

3.4.2 Initial amplitude and initial phase

In this subsection, we discuss the evolution of the ϕ field during the inflation.

3.4.2.a initial amplitude

First, we discuss the amplitude of the ϕ field. Let us first investigate the potential in the range $|\phi| \lesssim v$. In this range, the S field in Eq. (3.70) is expanded into the following form:

$$|S|^2 \simeq v^2 - \frac{1}{8}|\phi|^2 + \mathcal{O}\left(\frac{|\phi|^4}{v^2}\right), \quad \text{for } |\phi| \lesssim v. \quad (3.73)$$

By substituting Eq. (3.73) into Eq. (3.71), we get the effective potential for $\phi \lesssim v$ in the following form:

$$\begin{aligned} V \simeq & V_{\text{SB}} + \frac{1}{2}\mu^2|\phi|^2 + \frac{\sqrt{3}}{2}\frac{H}{M_{\text{eff}}}\left(c'_\phi\phi^4 + \text{H.c.}\right) + \frac{|\phi|^6}{4M_{\text{eff}}^2} \\ & - H^2|\phi|^2 + \frac{3}{8}(b'_S - b'_{\bar{S}})H^2|\phi|^2 + \dots, \end{aligned} \quad (3.74)$$

where the ellipsis denotes higher order terms in $|\phi|^2/v^2$. Here, $M_{\text{eff}} \equiv \xi v/h^2$ corresponds to the mass scale which was denoted by the same symbol in Sec. 3.2. (The mass of the right-handed neutrino in the present model is given by ξv .) Recall that it is related to the neutrino mass as $m_{\nu 1} = \langle H_u \rangle^2 / M_{\text{eff}}$.

The scalar potential in Eq. (3.74) is nothing but the potential obtained in Eq. (3.12) in the previous analysis, with some modifications.²¹ However, there is an important difference in the Hubble-induced SUSY breaking mass terms. Since there appears an extra Hubble mass term, we need the following condition in order that the ϕ field develops a large expectation value during the inflation:

$$b'_S - b'_{\bar{S}} \gtrsim \frac{8}{3}. \quad (3.75)$$

If this is not satisfied, the ϕ field is driven toward the origin during the inflation, and the leptogenesis cannot work. In the following discussion, we assume that the above condition is satisfied. Then the ϕ field has a negative Hubble mass term at least in the range of $|\phi| \lesssim v$.

Now let us discuss the scale where the flat direction is lifted. If the balance point between the negative Hubble mass term and the F -term potential $|\phi|^6/(4M_{\text{eff}}^2)$ in Eq. (3.74) is below the $B - L$ breaking scale v , the ϕ field is stopped at this balance point,

$$|\phi| \simeq \sqrt{M_{\text{eff}} H_{\text{inf}}} < v. \quad (3.76)$$

This is the condition for the “ F -term stopping case.” In this case, the evolution of the ϕ field and the amount of the generated baryon asymmetry result in the same conclusions as discussed in Sec. 3.2.

²¹As we mentioned, the potential terms coming from thermal effects will be discussed in Sec. 3.4.4.

On the other hand, if the condition

$$\sqrt{M_{\text{eff}} H_{\text{inf}}} \gtrsim v, \quad (3.77)$$

is satisfied, the ϕ field can develop its expectation value as large as v . In this case, we can stop the ϕ field below the balance point $\sqrt{M_{\text{eff}} H_{\text{inf}}}$ by using the D -term potential. This is very crucial to enhance the final baryon asymmetry in the D -term stopping case. To see this, let us discuss what happens if the ϕ field develops its value as large as the $B - L$ breaking scale v . At this scale (i.e., $|\phi| \simeq v$), the expansion of the S field given in Eq. (3.73) becomes invalid and above this scale we must use another expansion of the S field as follows:

$$|S|^2 \simeq 4 \frac{v^4}{|\phi|^2} + \frac{h^2}{2\xi^2} |\phi|^2 + \left[\mathcal{O}\left(\frac{v^4}{|\phi|^4}\right) + \mathcal{O}\left(\frac{h^2}{\xi^2}\right) \right]^2 |\phi|^2 \quad \text{for } |\phi| \gtrsim v. \quad (3.78)$$

One might wonder whether the above expansion is reliable, since Eq. (3.70) is based on Eq. (3.68), which may not be valid for $|\phi| \gg v$. Here, we first derive the conditions to fix the ϕ field at the scale v during the inflation, assuming this expansion is effectively valid at least for $|\phi| \sim v$. We will justify later the validity of the obtained conditions by numerical calculations.

By substituting the expansion Eq. (3.78) into Eq. (3.71), we obtain the following Hubble-induced mass term:²²

$$\delta V \simeq \left(-1 + \frac{3}{4}(1 - b'_{\bar{S}}) \right) H^2 |\phi|^2 + \mathcal{O}\left(\frac{v^4}{|\phi|^2} H^2\right) \quad \text{for } |\phi| \gtrsim v. \quad (3.79)$$

Therefore, the ϕ field can have a positive Hubble mass term for $|\phi| \gtrsim v$, if

$$b'_{\bar{S}} \lesssim -\frac{1}{3}. \quad (3.80)$$

Therefore, if this condition is satisfied, the ϕ field cannot develop its expectation value above the scale v , and hence it is fixed at the $B - L$ breaking scale v during the inflation.

To summarize, the ϕ field, and hence S and $\bar{S}$ fields as well, are stopped at the $B - L$ breaking scale v during the inflation, if the following conditions are satisfied,

$$b'_S - b'_{\bar{S}} \lesssim \frac{8}{3}, \quad \sqrt{M_{\text{eff}} H_{\text{inf}}} \gtrsim v, \quad b'_{\bar{S}} \lesssim -\frac{1}{3}. \quad (3.81)$$

These are the conditions for the D -term stopping case.

²² Here, we assume $h^2/\xi^2 \ll 1$. If this is not the case, we must include the Hubble mass term coming from the coupling of the right-handed Majorana neutrino to the inflaton, the last term in Eq. (3.71).




Figure 3.6: The amplitudes of the ϕ field fixed during inflation which are determined by numerical calculations. In this calculation, we have used the full scalar potential in Eq. (3.67) to follow the evolution of the relevant fields. Here, we have assumed that $H_{\text{inf}}/v = 0.1$, $3(1 - b'_\phi) = -1$, $h = 10^{-4}$, and $g = \xi = \lambda = 1$, and have randomly generated other coupling constants b_Y , $|c_Y|$ in the range $-2.5 \leq b_Y \leq 2.5$ and $0 \leq |c_Y| \leq 2.5$, respectively. Various symbols denote the $|\phi|$ to v ratio at the end of inflation. We see that the ϕ field is, in fact, stopped at the $B - L$ breaking scale v if the conditions in Eq. (3.81) are satisfied.

In order to check the reliability of the conditions in Eq. (3.81), (especially that of the last condition $b'_{\bar{S}} \lesssim -1/3$), we have numerically solved the coupled equations of motions

$$\ddot{Y} + 3H\dot{Y} + \frac{\partial V}{\partial Y^*} = 0, \quad (3.82)$$

for the relevant fields $Y = \phi, N, S, \bar{S}$ and X , using the full scalar potential in Eq. (3.67). We show the result in Fig. 3.6, where the amplitude of the ϕ field at the end of the inflation is plotted in $b'_S - b'_{\bar{S}}$ plane. Here, we have taken $M_{\text{eff}} = \xi v/h^2 = 10^8 v$ and $H_{\text{inf}} = 0.1v$, and hence the second condition $\sqrt{M_{\text{eff}} H_{\text{inf}}} \gtrsim v$ is satisfied.

It is found from Fig. 3.6 that the conditions in Eq. (3.81) well explain the result of this numerical calculation. Actually, for $b'_S - b'_{\bar{S}} \gtrsim 8/3$ (positive Hubble mass term for $|\phi| \lesssim v$), the amplitude is damped as $|\phi| \ll v$, while for $b'_{\bar{S}} \gtrsim -1/3$ (negative Hubble mass term for $|\phi| \gtrsim v$), the ϕ field overshoots the $B - L$ breaking scale v .

Thus, the ϕ field can stop at the $B - L$ breaking scale v if the conditions $b'_S - b'_{\bar{S}} \lesssim 8/3$ and $b'_{\bar{S}} \lesssim -1/3$ are satisfied. In fact, we see from Fig. 3.6 that the amplitude of the ϕ field at the end of the inflation lies in the range

$$|\phi| \simeq (1 - 3) \times v, \quad (3.83)$$

in most of the parameter space where the conditions in Eq. (3.81) are satisfied.

3.4.2.b initial phase

Now let us turn to discuss the phase of the ϕ field at the end of the inflation, which is crucial to estimate the resultant lepton asymmetry.

First, in the F -term stopping case, the amplitude of the ϕ field is fixed at $\sqrt{M_{\text{eff}} H_{\text{inf}}}$. An important point here is that the curvature around the valley of the Hubble A -term potential in Eq. (3.72) (which we denote by the symbol m_{phase}^2) is of the order of the Hubble parameter in this case. This can be seen from the following relation:

$$\begin{aligned} m_{\text{phase}}^2 &\simeq \left| \frac{\partial^2}{\partial \phi^2} V_{\text{phase}} \right| \\ &\simeq \frac{H_{\text{inf}}}{M_{\text{eff}}} |\phi|^2 \simeq H_{\text{inf}}^2. \end{aligned} \quad (3.84)$$

Therefore, as discussed in Sec. 3.2, the phase of the ϕ field settles down at the bottom of the valley of the A -term potential during the inflation, and hence the Hubble A -term cannot supply a torque to rotate the ϕ field. In this case, the relevant torque for the ϕ field only comes from the ordinary A -term potential proportional to the gravitino mass.

As stressed in Sec. 3.2.4, this leads to a suppression of the baryon asymmetry in gauge-mediation models with a small gravitino mass.

However, if the conditions in Eq. (3.81) are satisfied, namely, in the D -term stopping case, the curvature of the potential along the phase direction is smaller than the Hubble parameter during the inflation:

$$\begin{aligned} m_{\text{phase}}^2 &\simeq \left| \frac{\partial^2}{\partial \phi^2} V_{\text{phase}} \right| \\ &\simeq \frac{H_{\text{inf}}}{M_{\text{eff}}} |\phi|^2 \simeq H_{\text{inf}}^2 \left(\frac{|\phi|}{\sqrt{M_{\text{eff}} H_{\text{inf}}}} \right)^2 < H_{\text{inf}}^2. \end{aligned} \quad (3.85)$$

In this case, therefore, there is no reason to expect that the ϕ field sits down at the bottom of the valley of the A -term potential in Eq. (3.72) during the inflation. In other words, unless there is an accidental fine tuning of the initial phase of the ϕ field, it is generally displaced from the bottom of the valley of this A -term potential when the inflation ends.

Therefore, the A -term potential kicks the ϕ field along the phase direction when the Hubble parameter becomes comparable to the curvature along the phase direction m_{phase}^2 . As we will show in the next subsection, this phase rotation caused by the Hubble A -term is very crucial to enhance the final baryon asymmetry.

3.4.3 Lepton asymmetry

Now let us discuss the dynamics of the ϕ field and calculate the resultant lepton asymmetry in the present model. In the F -term stopping case, the amplitude of the ϕ field is always smaller than the $B - L$ breaking scale v [see Eq. (3.76)] and the evolution of the ϕ field is determined by the effective potential in Eq. (3.74) plus thermal effects, which we have investigated in Sec. 3.2. Thus, the resultant baryon asymmetry is given by Eq. (3.41) (or Fig. 3.4).

We turn to the D -term stopping case. As discussed in the previous subsection, the amplitude of the ϕ field at the end of the inflation is given by

$$|\phi| \simeq v, \quad (3.86)$$

and the phase of the ϕ field is generally displaced from the bottom of the valley of the Hubble A -term potential.

After the end of the inflation, the field value of the ϕ field takes an almost constant value for a while. Then the Hubble A -term in Eq. (3.72) kicks the ϕ field along the phase direction when the Hubble parameter H becomes comparable with the curvature:

$$H^2 \simeq m_{\text{phase}}^2 \left(\simeq \frac{H}{M_{\text{eff}}} v^2 \right). \quad (3.87)$$

Thus, the Hubble parameter at this time H_{ph} is given by

$$H_{\text{ph}} \simeq \frac{v^2}{M_{\text{eff}}} . \quad (3.88)$$

At this time, the phase of the ϕ field, $\arg(\phi)$, starts to oscillate around the bottom of the valley of the Hubble A -term potential. At the same time, the amplitude of the ϕ field, $|\phi|$, starts to decrease according to the following equation:

$$|\phi| \simeq \sqrt{M_{\text{eff}} H} , \quad (3.89)$$

which is the balance point between the negative Hubble mass term and the $|\phi|^6/(4M_{\text{eff}}^2)$ potential.

Since the phase of the ϕ field starts its oscillation at $H = H_{\text{ph}}$, it has already had a large acceleration along the *phase* direction before it starts coherent oscillation along the *radius* direction around the origin when $H = H_{\text{osc}}$. This H_{osc} is determined by thermal effects, soft SUSY breaking mass term and μ -term in the same way as in Sec. 3.2. Here, we have assumed that $H_{\text{ph}} > H_{\text{osc}}$, which we will discuss in Sec. 3.4.4.

In Fig. 3.7 and Fig. 3.8, we show the evolution of the amplitude and the phase of the ϕ field obtained by numerical calculations, respectively. In these figures, t is the cosmic time in the matter dominated universe $t = 2/(3H)$. We see from these figures that the evolution of the ϕ field is well explained by above arguments.

We are now at the point to estimate the baryon asymmetry. From the equation of motion (3.21), the evolution of the lepton asymmetry $n_L = (i/2)(\dot{\phi}^* \phi - \phi^* \dot{\phi})$ is given by

$$\dot{n}_L + 3Hn_L = 2\sqrt{3} \frac{H}{M_{\text{eff}}} \text{Im} \left(c'_\phi \phi^4 \right) , \quad (3.90)$$

where we have used $|S| \simeq v$ for $H \lesssim H_{\text{ph}}$. This can be easily integrated and we obtain the lepton number at time t as

$$\left[R^3 n_L \right] (t) \simeq \frac{2\sqrt{3}}{M_{\text{eff}}} \int^t dt' R^3 H \text{Im} \left(c'_\phi \phi^4 \right) . \quad (3.91)$$

Before the ϕ 's oscillation, i.e., for $H_{\text{osc}} \lesssim H \lesssim H_{\text{ph}}$, the amplitude of the ϕ field decreases as $|\phi| \simeq \sqrt{M_{\text{eff}} H}$ and the sign of $\text{Im}(c'_\phi \phi^4)$ in Eq. (3.91) changes with $1/H$ time scale. (See Fig. 3.7 and Fig. 3.8.) Therefore, we see from Eq. (3.91) that the total lepton number oscillates with $1/H$ time scale with almost constant amplitude, since $R^3 H |\phi^4| \propto H \propto t^{-1}$ in this period.

Then, the total lepton number is fixed when the ϕ field starts its coherent oscillation around the origin at $H = H_{\text{osc}}$, since the amplitude of the ϕ field decreases as fast as




Figure 3.7: The evolution of the ϕ field estimated by a numerical calculation. We assume that $M_{\text{eff}} = 3 \times 10^{23}$ GeV (i.e., $m_{\nu 1} = 10^{-10}$ eV), $v = 10^{14}$ GeV, $H_{\text{inf}} = 10^{12}$ GeV, $m_\phi = 10^3$ GeV. We see that the ϕ field starts to move at $H = H_{\text{ph}} \simeq 3 \times 10^4$ GeV according to the equation; $|\phi| \simeq \sqrt{M_{\text{eff}} H}$. Here, $t = 2/(3H)$ is the cosmic time. The ϕ field starts to oscillate around the origin at $H_{\text{osc}} \simeq m_\phi$ since we neglect the thermal effects in this calculation.




Figure 3.8: The evolution of the phase of the ϕ field estimated by a numerical calculation. The parameters used in this figure are the same as in Fig. 3.7. Here, we also defined that c'_ϕ is real, and hence the valleys coming from the Hubble A -term lie along $\arg(\phi) = \pi/4, 3\pi/4, 5\pi/4, 7\pi/4$. We see that the phase of the ϕ field starts to oscillate around the bottom of the valley at $H \simeq H_{\text{ph}}$.

$|\phi| \propto H$ and the integrand of Eq. (3.91) decreases as $\sim t^{-3}$ after the oscillation. The lepton number density at time $t_{\text{osc}} = 2/(3H_{\text{osc}})$ is then given by

$$\begin{aligned} n_L(t_{\text{osc}}) &\simeq \frac{2\sqrt{3}}{M_{\text{eff}}} H_{\text{osc}} \text{Im} \left[c'_\phi \phi^4(t_{\text{osc}}) \right] t_{\text{osc}} \\ &\simeq \frac{4}{\sqrt{3}} |c'_\phi| M_{\text{eff}} H_{\text{osc}}^2 \delta_{\text{ph}}, \end{aligned} \quad (3.92)$$

where $\delta_{\text{ph}} \simeq \sin(4 \arg \phi + \arg c'_\phi)$ is the effective CP -phase. By using Eq. (3.39), the final lepton asymmetry is given by

$$\frac{n_L}{s} \simeq \frac{M_{\text{eff}} T_R}{\sqrt{3} M_G^2} |c'_\phi| \delta_{\text{ph}}. \quad (3.93)$$

Here, we have used the fact that the ϕ field starts its oscillation before the reheating process completes, i.e., $H_{\text{osc}} > \Gamma_\chi$, which was verified in Sec. 3.2.4. Notice that the resultant lepton asymmetry is independent of the time when the ϕ field starts the oscillation around the origin, H_{osc} .²³ This is because the total lepton number itself oscillates with a *constant* amplitude for $H_{\text{osc}} \lesssim H \lesssim H_{\text{ph}}$, as discussed just above. Thus, the order of the lepton asymmetry is already fixed just after $H \simeq H_{\text{ph}}$, before the oscillation time H_{osc} .

By using the relation between the lepton and baryon asymmetry in the presence of the sphaleron effect, we obtain

$$\begin{aligned} \frac{n_B}{s} &\simeq 0.35 \times \frac{n_L}{s} \\ &\simeq 1 \times 10^{-10} \left(\frac{T_R}{10^8 \text{ GeV}} \right) \left(\frac{10^{-6} \text{ eV}}{m_{\nu 1}} \right)^{-1} |c'_\phi| \delta_{\text{ph}}. \end{aligned} \quad (3.94)$$

Here, we give several comments on the obtained baryon asymmetry. First, by comparing Eq. (3.93) and Eq. (3.40), we can see that the lepton asymmetry in the D -term stopping case is enhanced compared with the F -term stopping case (and the case without $U(1)_{B-L}$), since it is no longer suppressed by the oscillation time H_{osc} .

Next, as stressed before, one sees that the resultant baryon asymmetry is independent of the gravitino mass $m_{3/2}$. Therefore, we can obtain an enough baryon asymmetry even with a very light gravitino. This is the crucial point in this D -term stopping case. Here, we have to take care of the gravitino problem. For example, if the mass of the lightest neutrino is as small as $m_{\nu 1} \simeq 10^{-10}$ eV, the present baryon asymmetry is explained with a low reheating temperature $T_R \simeq 10^4$ GeV [see Eq. (3.94)]. In such a low reheating temperature, we are free from the overproduction of gravitinos even in gauge-mediated SUSY breaking scenarios with the small gravitino mass $m_{3/2} = \mathcal{O}(100 \text{ keV})$ [20].

²³We have also confirmed by numerical calculations that the final lepton asymmetry n_L/s is independent of the oscillation time H_{osc} .




Figure 3.9: The evolution of the baryon asymmetry n_B/s estimated by a numerical calculation. Parameters used in this figure are the same as in Fig. 3.7 and Fig. 3.8. Here, we also take $T_R = 10^4$ GeV. The thick line denotes the analytic estimation in Eq. (3.94). We can see that the asymmetry begins to oscillate with constant amplitude at $H \simeq H_{\text{ph}}$ and it is fixed when the ϕ field starts the oscillations around the origin. The analytical estimation of the baryon asymmetry agrees well with the obtained value by the numerical calculation.

On the other hand, if the mass of the gravitino is large enough, say, $m_{3/2} \simeq 1$ TeV, we can avoid the cosmological gravitino problem even if the reheating temperature is rather high as $T_R \simeq 10^9$ GeV (see Sec. 1.5). In such a case, we see from Eq. (3.94) that the mass of the lightest neutrino $m_{\nu 1} \simeq 10^{-4}$ eV is small enough to explain the present baryon asymmetry.

Finally, we show the evolution of the baryon asymmetry n_B/s obtained by a numerical calculation in Fig. 3.9. From this figure we see that the asymmetry begins to oscillate with almost constant amplitude at $H = H_{\text{ph}}$ and it is fixed at the time when the ϕ field starts to oscillate around the origin, $H = H_{\text{osc}}$. We also see that the amount of generated baryon asymmetry agrees well with our analytic estimation in Eq. (3.94). We confirm from this figure that the resultant asymmetry is well explained by the arguments described in this subsection.

3.4.4 Conditions to avoid an early oscillation

So far, we have neglected the thermal effects and assumed that there is no early oscillation, i.e., $H_{\text{ph}} > H_{\text{osc}}$. If the ϕ field starts its oscillation before $H = H_{\text{ph}}$, i.e., if $H_{\text{osc}} < H_{\text{ph}}$, the resultant baryon asymmetry is strongly suppressed. (In this case, the resultant baryon asymmetry is inversely proportional to H_{osc}^2 .) Thus, we verify the assumption $H_{\text{ph}} > H_{\text{osc}}$ here.

First, in order to avoid the early oscillation caused by the soft SUSY breaking mass m_ϕ and the μ -term, we need

$$m_\phi, \mu < H_{\text{ph}} \simeq \frac{v^2}{M_{\text{eff}}}. \quad (3.95)$$

Next, the thermal mass term $c_k f_k^2 T^2$ discussed in Sec. 3.2.3 may cause the early oscillation. This can be avoided if the following condition is satisfied:

$$H^2 > \sum_{f_k |\phi| < T} c_k f_k^2 T^2, \quad \text{for } H > H_{\text{ph}}. \quad (3.96)$$

Finally, we need the following third condition to avoid the early oscillation caused by the thermal log term $T^4 \ln(|\phi|^2)$, which was also explained in Sec. 3.2.3:

$$H^2 > \frac{a_g \alpha_S^2 T^4}{v^2}, \quad \text{for } H > H_{\text{ph}}. \quad (3.97)$$

Here, we have used $|\phi| \simeq v$ for $H > H_{\text{ph}}$.

From the second (3.96) and third (3.97) conditions, the reheating temperature is bounded from above as ²⁴

$$T_R < \min \left[T_{Ri}, \frac{v^2}{a_g^{1/2} \alpha_S M_G^{1/2} M_{\text{eff}}^{1/2}} \right], \quad (3.98)$$

where

$$T_{Ri} = \max \left(\frac{f_i v^{3/2}}{c_i^{1/4} M_G^{1/2}}, \frac{v^3}{c_i f_i^2 M_G^{1/2} M_{\text{eff}}^{3/2}} \right). \quad (3.99)$$

We show the allowed regions which are free from the early oscillation in Fig. 3.10, taking account the conditions in Eqs. (3.95) and (3.98). From this figure, we see that the early oscillations can be easily avoided, especially when the $B - L$ breaking scale satisfies $v \gtrsim 10^{14}$ GeV.

²⁴At first sight, Eq. (3.98) seems to be obtained just by rewriting the relation $H_{\text{osc}} < H_{\text{ph}}$ by using the formula of H_{osc} in Eq. (3.34). However, it is slightly different, since $|\phi|$ takes a constant value v for $H \gtrsim H_{\text{ph}}$ in the present case, while $|\phi| \simeq \sqrt{M_{\text{eff}} H}$ in the previous case.




Figure 3.10: The plot of the parameter regions which are free from the early oscillation in $m_{\nu 1} - T_R$ plane. The regions below the four solid lines are free from the early oscillation. These four lines correspond to the breaking scale of the $U(1)_{B-L}$, $v = 10^{16}, 10^{15}, 10^{14}, 10^{13}$ GeV from left to right, respectively. The shaded region corresponds to the present baryon asymmetry, $n_B/s \simeq (0.4 - 1) \times 10^{-10}$. We have taken $\delta_{\text{ph}} = 1$ and $|c'_\phi| = 1$ in this figure. We see that if we take the breaking scale of the $U(1)_{B-L}$ as $v \gtrsim 10^{14}$ GeV, the early oscillation of the ϕ field $H_{\text{osc}} > H_{\text{ph}}$ can be avoided in most region of parameter space.

3.5 The neutrino-less double-beta decay

In Sec. 3.2, we have investigated the leptogenesis via LH_u flat direction in detail and have found that the produced baryon asymmetry is determined almost independently of the reheating temperature T_R , because of the early oscillation of the flat direction field ϕ caused by thermal effects. As shown there, an ultralight neutrino $m_{\nu 1} \simeq 10^{-9}$ eV is required to explain the empirical baryon asymmetry $n_B/s \simeq 10^{-10}$, except for the case where the model in Sec. 3.3.2 is adopted.²⁵ By introducing a gauged $U(1)_{B-L}$ symmetry, the early oscillation of the flat direction due to the thermal effect can be avoided (Sec. 3.4), but small neutrino mass $m_{\nu 1} \ll 10^{-4}$ eV is still necessary if we require low enough reheating temperature $T_R \lesssim 10^9$ GeV to avoid the cosmological gravitino problem in a wide range of gravitino mass (see Fig. 3.10). In this section we show that such a tiny mass of the lightest neutrino mass $m_{\nu 1} \ll 10^{-4}$ eV have an interesting implication on a low-energy experiment; the neutrino-less double beta ($0\nu\beta\beta$) decay, which will be a strong evidence for the lepton number violation if observed.

The crucial parameter to determine the $0\nu\beta\beta$ decay rate is the ν_e - ν_e component of the neutrino mass matrix, which is given by

$$m_{ee} \equiv \sum_i U_{ei}^2 m_{\nu i}, \quad (3.100)$$

where the mixing matrix $U_{\alpha i}$ is defined in Sec. 2.1. If the mass of the lightest neutrino is actually so small as $m_{\nu 1} \ll 10^{-4}$ eV, the contribution from $m_{\nu 1}$ to m_{ee} can be neglected as long as we consider an accessible $0\nu\beta\beta$ decay rate, which corresponds to $|m_{ee}| \gtrsim 10^{-3}$ eV. Thus, the parameter $|m_{ee}|$ is written in terms of masses and mixings of two other neutrinos as

$$|m_{ee}| = |U_{e2}^2 m_{\nu 2} + U_{e3}^2 m_{\nu 3}|, \quad (3.101)$$

which immediately leads to the following upper and lower bounds:

$$\begin{aligned} |m_{ee}|_{\max} &= |U_{e2}|^2 m_{\nu 2} + |U_{e3}|^2 m_{\nu 3}, \\ |m_{ee}|_{\min} &= \left| |U_{e2}|^2 m_{\nu 2} - |U_{e3}|^2 m_{\nu 3} \right|. \end{aligned} \quad (3.102)$$

In general, the mass pattern of the neutrinos can be classified into two cases: the normal hierarchy and the inverted hierarchy. Let us consider these two cases in turn.

²⁵See also the last comment in Sec. 3.2.4.

normal mass hierarchy

In the case of normal mass hierarchy, the parameters of atmospheric and solar neutrino oscillations are given by

$$\delta m_{\text{atm}}^2 = m_{\nu 3}^2 - m_{\nu 2}^2, \quad \delta m_{\text{sol}}^2 = m_{\nu 2}^2, \quad \tan^2 \theta_{\text{sol}} \equiv |U_{e2}/U_{e1}|^2, \quad (3.103)$$

and the mixing angle constrained by CHOOZ experiment [116] corresponds to $\sin \theta_{\text{chooz}} = |U_{e3}|$, which is severely bounded as $\sin \theta_{\text{chooz}} < 0.15$. In terms of these oscillation parameters, Eqs. (3.102) become

$$\begin{aligned} |m_{ee}|_{\text{max}} &= c_{\text{chooz}}^2 s_{\text{sol}}^2 \sqrt{\delta m_{\text{sol}}^2} + s_{\text{chooz}}^2 \sqrt{\delta m_{\text{atm}}^2 + \delta m_{\text{sol}}^2}, \\ |m_{ee}|_{\text{min}} &= \left| c_{\text{chooz}}^2 s_{\text{sol}}^2 \sqrt{\delta m_{\text{sol}}^2} - s_{\text{chooz}}^2 \sqrt{\delta m_{\text{atm}}^2 + \delta m_{\text{sol}}^2} \right|, \end{aligned} \quad (3.104)$$

where $c_X^2 \equiv \cos^2 \theta_X$ and $s_X^2 \equiv \sin^2 \theta_X$. Here, we can safely neglect the contribution of δm_{sol}^2 in the second term $s_{\text{chooz}}^2 \sqrt{\delta m_{\text{atm}}^2 + \delta m_{\text{sol}}^2}$, since it changes $|m_{ee}|$ at most $\Delta|m_{ee}| < s_{\text{chooz}}^2 \delta m_{\text{sol}}^2 / (2\sqrt{\delta m_{\text{atm}}^2}) < 10^{-4}$ eV.

Then, the ratio of the mass parameter $|m_{ee}|$ to the mass scale of atmospheric neutrino oscillation $\sqrt{\delta m_{\text{atm}}^2}$ is given by

$$\begin{aligned} \frac{|m_{ee}|_{\text{max}}}{\sqrt{\delta m_{\text{atm}}^2}} &= c_{\text{chooz}}^2 s_{\text{sol}}^2 \sqrt{\frac{\delta m_{\text{sol}}^2}{\delta m_{\text{atm}}^2}} + s_{\text{chooz}}^2, \\ \frac{|m_{ee}|_{\text{min}}}{\sqrt{\delta m_{\text{atm}}^2}} &= \left| c_{\text{chooz}}^2 s_{\text{sol}}^2 \sqrt{\frac{\delta m_{\text{sol}}^2}{\delta m_{\text{atm}}^2}} - s_{\text{chooz}}^2 \right|. \end{aligned} \quad (3.105)$$

Fig. 3.11 shows the upper and lower bounds on $|m_{ee}|$ in units of $\sqrt{\delta m_{\text{atm}}^2}$, for $\sin \theta_{\text{chooz}} = 0.15, 0.10, 0.05$ and 0 . As can be seen in Fig. 3.11, the mass parameter $|m_{ee}|$ is predicted in a narrow range. We stress that this is because of the absence of the contribution from the lightest neutrino, i.e., $m_{\nu 1} \ll 10^{-4}$ eV. It is quite interesting to see that the predicted $|m_{ee}|$ can be as large as $|m_{ee}| \simeq 0.001\text{--}0.01$ eV, which could be accessible at future $0\nu\beta\beta$ decay experiments [120]. Furthermore, it is also found from Fig. 3.11 that, if the $\sin \theta_{\text{chooz}}$ becomes more constrained by future experiments, $|m_{ee}|$ is predicted in a much narrower range.




Figure 3.11: The predicted value of $|m_{ee}|$ in units of $\sqrt{\delta m^2_{\text{atm}}}$, in the case of normal mass hierarchy. The lines represent the upper and lower values of $|m_{ee}|$ for $\sin \theta_{\text{chooz}} = 0.15, 0.10, 0.05$ and 0 . The vertical line corresponds to the set of best fit values, $\sin^2 \theta_{\text{sol}} \simeq 0.3$, $\delta m^2_{\text{sol}} \simeq 7 \times 10^{-5} \text{ eV}^2$ [115] and $\delta m^2_{\text{atm}} = 2.5 \times 10^{-3} \text{ eV}^2$ [10].




Figure 3.12: The predicted value of $|m_{ee}|$ in units of $\sqrt{\delta m^2_{\text{atm}}}$, in the case of inverted mass hierarchy. The lines represent the upper and lower values of $|m_{ee}|$. The vertical line corresponds to the best fit value, $\sin^2 \theta_{\text{sol}} \simeq 0.3$ [115].

inverted hierarchy

Now let us turn to discuss the case of inverted mass hierarchy, where the parameters of atmospheric and solar neutrino oscillations are given by²⁶

$$\delta m_{\text{atm}}^2 = m_{\nu 3}^2, \quad \delta m_{\text{sol}}^2 = m_{\nu 3}^2 - m_{\nu 2}^2, \quad \tan^2 \theta_{\text{sol}} \equiv |U_{e3}/U_{e2}|^2. \quad (3.106)$$

The mixing angle constrained by CHOOZ experiment [116] now corresponds to $\sin \theta_{\text{chooz}} = |U_{e1}|$. Then from Eqs. (3.102) we obtain

$$\begin{aligned} \frac{|m_{ee}|_{\text{max}}}{\sqrt{\delta m_{\text{atm}}^2}} &= c_{\text{chooz}}^2 \left[c_{\text{sol}}^2 \sqrt{1 - \frac{\delta m_{\text{sol}}^2}{\delta m_{\text{atm}}^2}} + s_{\text{sol}}^2 \right], \\ \frac{|m_{ee}|_{\text{min}}}{\sqrt{\delta m_{\text{atm}}^2}} &= c_{\text{chooz}}^2 \left| c_{\text{sol}}^2 \sqrt{1 - \frac{\delta m_{\text{sol}}^2}{\delta m_{\text{atm}}^2}} - s_{\text{sol}}^2 \right|. \end{aligned} \quad (3.107)$$

In Fig. 3.12, we show the upper and lower bounds on $|m_{ee}|$ in units of $\sqrt{\delta m_{\text{atm}}^2}$, for given $\tan^2 \theta_{\text{sol}}$. Here, we have used $\sin \theta_{\text{chooz}} = 0$ for simplicity, but $|m_{ee}|$ changes (reduces) only 2% even for $\sin \theta_{\text{chooz}} = 0.15$. As for the ratio $r \equiv \delta m_{\text{sol}}^2 / \delta m_{\text{atm}}^2$, we have adopted $r = 0.008\text{--}0.15$ [10, 115]. As seen from the figure, the $|m_{ee}|$ is restricted in a very narrow range such as $|m_{ee}| \simeq (0.01\text{--}0.05) \text{ eV} \times (\sqrt{\delta m_{\text{atm}}^2} / 0.05 \text{ eV})$ for $\tan^2 \theta_{\text{sol}} < 0.7$, which is indeed in the reach of the future $0\nu\beta\beta$ decay experiments [120].

The prediction of $|m_{ee}|$ discussed in this section is a generic consequence of a hierarchical neutrino masses $m_{\nu 1} \ll m_{\nu 2}, m_{\nu 3}$. Thus the $0\nu\beta\beta$ decay experiments, combined with neutrino oscillation experiments, can provide a consistency test for the tiny mass of the lightest neutrino $m_{\nu 1} \ll 10^{-4} \text{ eV}$ required in the leptogenesis via LH_u flat direction.

²⁶The convention adopted here is different from the one in Ref. [103], where $m_{\nu i}$ are defined as $m_{\nu 3} < m_{\nu 1} < m_{\nu 2}$ for inverted hierarchy.

3.A D -flat condition for the LH_u flat direction

In this appendix, we discuss the D -flat condition along the LH_u direction. Let us take the number of generations of the lepton doublets L_i to be N in general. Then, there are $(2 + 2N)$ complex scalar fields in the L - H_u sector.

$$\begin{aligned}\widetilde{L}_i &= \begin{pmatrix} \widetilde{\nu}_{Li} \\ \widetilde{e}_{Li} \end{pmatrix} \quad i = 1 \cdots N, \\ H_u &= \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix}.\end{aligned}\tag{3.108}$$

First, the freedom to make $SU(2)_L$ gauge transformation allows us to rotate away a possible vacuum expectation value of one of the weak isospin components of one of the scalar fields. Thus, we can take $H_u^+ = 0$ without loss of generality. Then, vanishing D -term potential requires the following relations:

$$-|\widetilde{\nu}_{Li}|^2 - |\widetilde{e}_{Li}|^2 + |H_u^0|^2 = 0 \quad \text{for } U(1)_Y,\tag{3.109}$$

$$\begin{aligned}\widetilde{\nu}_{Li}^* \widetilde{e}_{Li} + \widetilde{e}_{Li}^* \widetilde{\nu}_{Li} &= 0 \\ \widetilde{\nu}_{Li}^* \widetilde{e}_{Li} - \widetilde{e}_{Li}^* \widetilde{\nu}_{Li} &= 0 \quad \text{for } SU(2)_L, \\ |\widetilde{\nu}_{Li}|^2 - |\widetilde{e}_{Li}|^2 - |H_u^0|^2 &= 0\end{aligned}\tag{3.110}$$

where the summations over i are suppressed for simplicity. From these equations, the D -flat condition is given by

$$\sum_{i=1}^N |\widetilde{\nu}_{Li}|^2 = |H_u^0|^2.\tag{3.111}$$

It is still possible to rotate away one of the phases by using the $U(1)_Y$ gauge transformation (or third component of the $SU(2)_L$ gauge transformations²⁷). Therefore, this flat “direction” is parameterized by $[(N+1) \text{ complex fields}] - [\text{one real constraint Eq. (3.111)}] - [\text{one phase}] = 2N$ real fields. In the absence of effective superpotential Eq. (3.2), this “direction” would be completely flat in the SUSY limit. Furthermore, it would be isotropic for N directions, $\widetilde{\nu}_{L1}, \dots, \widetilde{\nu}_{LN}$. However, due to the scalar potential induced by the operator Eq. (3.2), the potential is lifted and modified. Then, the flattest direction among the above field space is parameterized by one scalar field ϕ , which corresponds to the first family $i = 1$.

²⁷Notice that one $U(1)$ group remains unbroken by the condensation of the flat direction field.

Chapter 4

Conclusions and discussion

In this thesis, we have investigated various leptogenesis mechanisms in the framework of SUSY. We have also discussed some aspects of those scenarios, such as constraints from cosmological gravitino problems and possible experimental signatures. In particular, we have made essential improvements compared with earlier works in the following points:

- In the leptogenesis scenario where the lepton asymmetry is created by the decay of right-handed sneutrino $\tilde{N}$ having dominated the early universe (Sec. 2.4), the dilution of the gravitino due to the entropy production by the $\tilde{N}$'s decay has been discussed. As a result, it has been shown that the cosmological gravitino problem is drastically ameliorated.
- In the leptogenesis via the LH_u flat direction (Chapter 3), all the relevant thermal effects has been taken into account, and detailed analyses on the dynamics of the flat direction field has been done both by analytic and numerical calculations.

In this chapter, we summarize the aspects of various leptogenesis mechanisms investigated in this thesis. The produced baryon asymmetry in each mechanism is well summarized in Table 4.1.

First of all, we see that in all the mechanisms the neutrino mass plays a very crucial role to determine the resultant baryon asymmetry. In the mechanisms where lepton asymmetry is produced by the asymmetric decay of the right-handed (s)neutrino (Chapter 2), the asymmetry parameter ϵ_1 is proportional to the mass of the heaviest neutrino $m_{\nu 3}$ for fixed effective CP -violating phase δ_{eff} . We see that the information from the atmospheric neutrino oscillation [10], which leads to $m_{\nu 3} \simeq 0.05$ eV as long as masses of the light neutrinos are not degenerate, is very important to calculate the final baryon asymmetry. On the other hand, when the lepton asymmetry is produced via the LH_u flat direction (Chapter 3), the mass of the lightest neutrino $m_{\nu 1}$ is the crucial parameter.

	produced baryon asymmetry
thermal ⁽¹⁾	$\frac{n_B}{s} \simeq 0.35 \times \kappa \frac{\epsilon_1}{240}$ $\simeq 0.3 \times 10^{-10} \left(\frac{\kappa}{0.1} \right) \left(\frac{M_1}{10^9 \text{ GeV}} \right) \cdot \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}$
inflaton decay ^{(2) (a)}	$\frac{n_B}{s} \simeq 0.35 \times \frac{3}{2} \epsilon_1 B_r \frac{T_R}{m_\chi}$ $\simeq 0.5 \times 10^{-10} B_r \left(\frac{T_R}{10^6 \text{ GeV}} \right) \left(\frac{2 M_1}{m_\chi} \right) \cdot \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}$
$\widetilde{N}$ dominant ⁽³⁾	$\frac{n_B}{s} \simeq 0.35 \times \frac{3}{4} \epsilon_1 \frac{T_{N_1}}{M_1}$ $\simeq 0.5 \times 10^{-10} \left(\frac{T_{N_1}}{10^6 \text{ GeV}} \right) \cdot \left(\frac{m_{\nu 3}}{0.05 \text{ eV}} \right) \delta_{\text{eff}}$
LH_u flat direction ^{(4) (b)}	$\frac{n_B}{s} \simeq 0.35 \times \frac{M_{\text{eff}} T_R}{12 M_G^2} \left(\frac{m_{3/2} a_m }{H_{\text{osc}}} \right) \delta_{\text{ph}}$ $\rightarrow m_{\nu 1} \simeq (0.1\text{--}3) \times 10^{-9} \text{ eV}^{(c)} \text{ for } T_R \gtrsim 10^5 \text{ GeV}$ (See Fig. 3.4.)
+ gauged $U(1)_{B-L}$ D -term stopping ⁽⁵⁾	$\frac{n_B}{s} \simeq 0.35 \times \frac{M_{\text{eff}} T_R}{\sqrt{3} M_G^2} c'_\phi \delta_{\text{ph}}$ $\simeq 1 \times 10^{-10} \left(\frac{T_R}{10^8 \text{ GeV}} \right) \left(\frac{10^{-6} \text{ eV}}{m_{\nu 1}} \right)^{-1} c'_\phi \delta_{\text{ph}}$

Table 4.1: Summary of produced baryon asymmetries in various leptogenesis mechanisms. Necessary conditions are: (1) $T_R > M_1$. (2) $M_1 > T_R$ and $m_\chi > 2M_1$. (3) $M_1 > T_{N_1}$ and $T_R \gg T_{N_1}$ (for $|\widetilde{N}_{1i}| \simeq M_G$). (4) negative Hubble mass for ϕ . (5) See Eq. (3.81) and Fig. 3.10. Some comments are in order: (a) Perturbative decay is assumed. For preheating, see Sec. 2.3.6. (b) $H_{\text{inf}} > H_{\text{osc}}$ is assumed. (See the last comment in Sec. 3.2.4.) (c) This is not the case if effective mass for the right-handed neutrinos are very large during leptogenesis (Sec. 3.3.2).

It is interesting to see that the dependences on the neutrino masses are opposite in those two scenarios. As for the ϵ_1 parameter of the right-handed neutrino decay, larger Yukawa coupling is favored to make ϵ_1 larger. When we normalize the Yukawa couplings by neutrino masses (Sec. 2.1), ϵ_1 turns out to be proportional to $m_{\nu 3}$ for fixed δ_{eff} . On the other hand, for leptogenesis via LH_u flat direction, flatter direction is favored to enhance the final lepton asymmetry. This is because, the flatter the potential is, the larger the amplitude of ϕ becomes, and the larger the final lepton asymmetry is produced. Thus in this case the relevant neutrino mass is the lightest one, $m_{\nu 1}$.

Now let us comment on each mechanism in turn.

- **thermal production of the right-handed (s)neutrino**

The simplest one is the thermal production of the right-handed (s)neutrino (Sec. 2.2). The attractive point in this scenario is nothing but the fact that the production of the right-handed (s)neutrino requires no assumption besides high enough reheating temperature $T_R > M_1$. The production occurs just by ordinary thermal scatterings.

On the other hand, since this scenario requires high enough reheating temperature, cosmological gravitino problems should be considered seriously. One solution is the stable (LSP) gravitino with mass $m_{3/2} \sim 100$ GeV. In this case, one has to take care of the decays of the next-to-lightest SUSY particle (NLSP) after the big-bang nucleosynthesis [20, 48], which gives severe constraints on the lifetime and the abundance of the NLSP at the decay. Since both of the lifetime and the abundance of the NLSP is determined by the properties of the NLSP (and the gravitino mass), they might be going to be well calculated if SUSY particles are discovered and their properties are studied in future collider experiments. This is quite an interesting possibility.

If the gravitino is not the LSP, cosmological gravitino problem is relatively severe. Therefore, it will be interesting and might be necessary to determine a definite lower bound on the reheating temperature T_R at which the leptogenesis from thermally produced right-handed (s)neutrino works marginally. This requires to see what happens if the reheating temperature is just around the mass of the right-handed neutrino, $T_R \sim M_1$.

- **leptogenesis in inflaton decay**

The advantage of this mechanism is that the production of the right-handed (s)neutrino is possible for $T_R < M_1$. Thus the cosmological gravitino problem can be avoided in a wider range of the gravitino mass. Actually, we have shown that the enough lepton asymmetry can be produced even with a reheating temperature $T_R \simeq 10^6$ GeV. On the other hand,

the produced baryon asymmetry crucially depends on the physics of the inflation. In particular, we have taken the branching ratio $B_r(\chi \rightarrow N_1 N_1)$ to be $B_r \simeq 1$. In this sense the first one of the hybrid inflation models we have discussed in Sec. 2.3.3.a, which has Yukawa couplings between the inflaton and right-handed neutrinos, is attractive since it automatically gives rise to $B_r \simeq 1$.

We should mention that the current and future satellite experiments on anisotropies of the cosmic microwave background radiation [90] will give precious information about the inflation physics, especially the spectrum index n_s . Thus we will be able to distinguish which inflation model is favored by using future data.

By the way, from Table 4.1, it is found that the formula of the baryon asymmetry from inflaton decay reproduces that of the $\widetilde{N}$ -dominant universe if we take $B_r = 1$, $T_R = T_{N_1}$ and $2M_1 = m_\chi$. This corresponds to the case where the right-handed sneutrino plays a role of the inflaton itself [24, 93].

• $\widetilde{N}$ -dominant universe

The resultant baryon asymmetry in this scenario is expressed in a very simple way, as given in Table 4.1. It is determined only by the decay temperature T_{N_1} for fixed $m_{\nu 3}$ and δ_{eff} . This is because the $\widetilde{N}_1$'s decay produces dominant component of the radiation as well as the lepton asymmetry at once.

The most attractive feature of this scenario is the dilution of the gravitino. In fact, as we have shown, cosmological gravitino problems can be avoided even for $T_R \gg 10^{11}$ GeV, which is very special among other baryogenesis scenarios.

It is also quite an interesting possibility that the cosmic density perturbation coming from the $\widetilde{N}_1$'s fluctuation might be detected at future experiments [99].

• LH_u flat direction

We have performed a detailed analysis on this scenario, including all the relevant thermal effects. In spite of the complicated potential and dynamics of the flat direction field ϕ , it was shown that the result is very simple and attractive. The baryon asymmetry is determined almost only by the mass of the lightest neutrino $m_{\nu 1}$. Remarkably, we can estimate (or predict) the $m_{\nu 1}$ from the empirical baryon asymmetry $n_B/s \simeq (0.4-1) \times 10^{-10}$ as

$$m_{\nu 1} \simeq (0.1 - 3) \times 10^{-9} \text{ eV}, \quad (4.1)$$

almost independently of the reheating temperature for $T_R \gtrsim 10^5$ GeV. It is amazing that the baryogenesis which happened in the very early universe is directly related to the neutrino mass.

- **LH_u flat direction + gauged $U(1)_{B-L}$, D -term stopping case**

When we introduce an gauged $U(1)_{B-L}$ symmetry, the dynamics of the LH_u flat direction field ϕ can be modified, if certain conditions are satisfied (D -term stopping case). Then the resultant baryon asymmetry is enhanced compared with the case with F -term stopping case, or the case without a gauged $U(1)_{B-L}$ symmetry. Although the “reheating temperature independence” disappears in this case (and $m_{\nu 1}$ is no longer predicted), this scenario has an advantage in the case of gauge-mediated SUSY breaking (GMSB) models [45], compared to the previous case. The point is that the phase rotational motion of the ϕ field is provided not by the usual SUSY-breaking A -term which is proportional to the gravitino mass, but by a Hubble-induced A -term. Thus this mechanism can work even with a very light gravitino as in the GMSB models.

- **The neutrinoless double beta decay**

We have also shown that a very light mass of the lightest neutrino $m_{\nu 1} \ll 10^{-4}$ eV, which is suggested by the above leptogenesis mechanisms via LH_u flat direction, leads to a high predictability of the rate of the neutrino-less double beta ($0\nu\beta\beta$) decay. Although it is highly difficult to determine the mass of the lightest neutrino directly, future experiments on $0\nu\beta\beta$ decay and neutrino oscillations will provide us precious information by which we can test the consistency of the very small lightest neutrino mass $m_{\nu 1} \ll 10^{-4}$ eV, as shown in Fig. 3.11 and 3.12.

Acknowledgement

I am very grateful to my advisor T. Yanagida for various instructive suggestions, stimulating discussions and collaborations, hearty encouragements and continuous support.

I would like to thank T. Asaka, M. Fujii, M. Kawasaki and H. Murayama, who are collaborators in various parts of this thesis, for fruitful discussions and collaborations.

I would also like to thank T. Watari for helpful discussions, and K. Ichikawa for helping me maintain the computer.

Finally, I thank all the members of particle physics theory group at University of Tokyo for their hospitality.

Appendix A

Notations

We use natural units $c = h/(2\pi) = 1$, where c is the speed of light (in vacuum) and h is the Planck constant. We also take the Boltzmann constant k_B to be unity, which leads to $1 \text{ eV} = 11605K$.

$M_G \equiv (8\pi G)^{-1/2} = 2.4 \times 10^{18} \text{ GeV}$ denotes the reduced Planck scale, where G is the Newton constant.

Throughout this thesis, except for generic discussions in Sec. 3.1, 3.2, 3.5 and 3.A, we assume the existence of the right-handed neutrinos, which have superpotentials:

$$W = \frac{1}{2}M_i N_i N_i + h_{i\alpha} N_i L_\alpha H_u, \quad (\text{A.1})$$

where N_i ($i = 1, 2, 3$), L_α ($\alpha = e, \mu, \tau$) and H_u denote the supermultiplets of the heavy right-handed neutrinos, lepton doublets and the Higgs doublet which couples to up-type quarks, respectively.

We often use a basis where the mass matrix of the light neutrinos becomes diagonal, where the Yukawa couplings are given by $\hat{h}_{ij}$. (The left subscript i denotes the family of the heavy right-handed neutrinos N_i , while the right subscript j denotes the family of the light neutrinos ν_j in mass eigenstates. See Sec. 2.1 and Eq. (2.12).)

As for the parameters related to the cosmology, see next Appendix B.

Appendix B

Some notes of standard cosmology

In this appendix, we briefly review some ingredients of the standard cosmology, in order to introduce our notations and relevant equations. For more details, see, for example, Ref. [28].

B.1 Friedmann-Robertson-Walker universe

The standard cosmology is based on the following homogeneous and isotropic metric (Robertson-Walker metric):

$$ds^2 = dt^2 - R(t)^2 \left\{ \frac{dr^2}{1 - kr^2} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right\}, \quad (\text{B.1})$$

where $R(t)$ is the “scale factor” of the expanding universe, and t is the cosmic time. (t, r, θ, ϕ) is called the comoving coordinates. From the above metric and the perfect-fluid form of the energy-momentum tensor $T_{\mu\nu} = \text{diag}(\rho, p, p, p)$ ($\rho(t)$ and $p(t)$ denote the energy density and the pressure, respectively), the Einstein equations lead to the following two equations:

$$\left(\frac{\dot{R}}{R} \right)^2 + \frac{k}{R^2} = \frac{8\pi G}{3} \rho, \quad (\text{B.2})$$

$$\frac{d}{dt} (\rho R^3) = -p \frac{d}{dt} (R^3), \quad (\text{B.3})$$

where the overdot denotes the derivative with cosmic time t , and G is the Newton constant.

The second equation (B.3), which corresponds to the conservation of the energy and momentum $T^{\mu\nu}_{;\nu} = 0$, describes the dilution of the energy density due to the expansion:

$$\text{radiation} \quad (p_{\text{rad}} = \tfrac{1}{3}\rho_{\text{rad}}) \quad \rho_{\text{rad}} \propto R^{-4},$$

$$\begin{aligned}
\text{matter} \quad & (p_{\text{mat}} = 0) \quad \rho_{\text{mat}} \propto R^{-3}, \\
\text{vacuum} \quad & (p_{\text{vac}} = -\rho_{\text{vac}}) \quad \rho_{\text{vac}} = \text{const}.
\end{aligned} \tag{B.4}$$

Notice that the above relations hold for each component of the energy density ρ_X , as long as the energy-momentum conservation $T_X^{\mu\nu}{}_{;\nu} = 0$ is satisfied separately for each X .

The other equation (B.2), called Friedmann equation, describes the expansion of the universe. Notice that the curvature term k/R^2 decreases with time more slowly than the right-hand side of Eq. (B.2) as long as ρ is dominated by matter (ρ_{mat}) or radiation (ρ_{rad}) [see Eq. (B.4)]. Therefore, it can be safely neglected as far as we consider the early universe.

The Hubble parameter is defined as

$$H \equiv \frac{\dot{R}}{R}, \tag{B.5}$$

which means how fast the universe is expanding at a given stage of its expansion. In terms of this Hubble parameter H , reduced Planck mass $M_G \equiv (8\pi G)^{-1/2} = 2.4 \times 10^{18}$ GeV, critical energy density $\rho_{\text{crit}} \equiv 3M_G^2 H^2$ and density parameter $\Omega \equiv \rho/\rho_{\text{crit}}$, Eq. (B.2) is rewritten as follows:

$$\Omega = \frac{\rho}{3M_G^2 H^2} = 1 + \frac{k}{R^2 H^2}. \tag{B.6}$$

Neglecting the curvature term, usually we use the following form of the Friedmann equation:

$$\rho = 3M_G^2 H^2. \tag{B.7}$$

The present universe ($t = t_0$) observed is as follows [29]. The present Hubble expansion rate is $H_0 = 100 h [\text{km sec}^{-1} \text{Mpc}^{-1}]$, where $h \simeq 0.7$, and hence critical energy density is $\rho_{\text{crit}}(t_0) = 3M_G^2 H_0^2 = 1.9 h^2 \times 10^{-29} [\text{g cm}^{-3}]$. In terms of density parameter $\Omega_X \equiv \rho_X/\rho_{\text{crit}}$, total energy density is given by $\Omega(t_0) \simeq 1$, which includes radiation $\Omega_{\text{rad}}(t_0) = (2.47 + 0.56 N_\nu^{\text{eff}}) h^{-2} \times 10^{-5}$ (N_ν^{eff} is the number of the generations of neutrinos which are still relativistic today), matter $\Omega_{\text{mat}}(t_0) \simeq 0.3$ (including baryon $\Omega_B(t_0) \simeq 0.02 h^{-2}$) and “dark energy” $\Omega_{\text{vac}}(t_0) \simeq 0.7$.¹ For more details and the errors included in these values, see Ref. [29]. The nature of the cold dark matter (dominant component of the Ω_{mat}) as well as that of the “dark energy” are big puzzles in cosmology, which is beyond the scope of this appendix. (As for the cold dark matter, however, supersymmetry provides a good candidate, the lightest SUSY particle (LSP). See also Sec. 2.4.3.)

¹The notation “ Ω_{vac} ” might not be correct, since the “dark energy” density could be a dynamical value (like “quintessence”), not a cosmological constant.

From the above ratio of the present energy densities, it is found that the early universe was dominated by radiation (for $R \lesssim 10^{-4}R(t_0)$), and subsequently dominated by matter. Eqs. (B.4) and (B.7) lead to the following familiar relations for each epoch.²

$$\begin{aligned} \text{radiation-dominated universe} \quad R &\propto t^{1/2}, \quad H = \frac{1}{2t}, \\ \text{matter-dominated universe} \quad R &\propto t^{2/3}, \quad H = \frac{2}{3t}. \end{aligned} \quad (\text{B.8})$$

B.2 Thermodynamics

Here, we briefly review some basic thermodynamics in the radiation-dominated early universe, in which many particles are relativistic and in thermal equilibrium. First of all, the equilibrium density of particles of type i with momenta in a range d^3p centered on $\mathbf{p}$ is given by

$$g_i \frac{d^3p}{2\pi^3} f_i(\mathbf{p}), \quad (\text{B.9})$$

where g_i is the number of degrees of freedom and $f_i(\mathbf{p})$ is the Fermi-Dirac or Bose-Einstein distribution function:

$$f_i(\mathbf{p}) = \frac{1}{\exp\left(\frac{E_i - \mu_i}{T}\right) \pm 1}. \quad (\text{B.10})$$

Here, E_i is the energy $E_i \equiv \sqrt{\mathbf{p}^2 + m_i^2}$, μ_i is the chemical potential of the particle i , and the plus (minus) sign is for fermions (bosons). The number density n_i , energy density ρ_i and pressure p_i of particle i are then given by the following equations:

$$n_i = \frac{g_i}{2\pi^3} \int f_i(\mathbf{p}) d^3p, \quad (\text{B.11})$$

$$\rho_i = \frac{g_i}{2\pi^3} \int E_i f_i(\mathbf{p}) d^3p, \quad (\text{B.12})$$

$$p_i = \frac{g_i}{2\pi^3} \int \frac{\mathbf{p}^2}{3E_i} f_i(\mathbf{p}) d^3p. \quad (\text{B.13})$$

In Table B.1, we show these quantities for the relativistic ($T \gg m_i$) and non-relativistic ($T \ll m_i$) limits. Here, we have assumed $|\mu_i| \ll T$ and no Bose-Einstein condensation ($|\mu_i| < m_i$).

²If we include the change of $g_*(T)$ with time, which is defined below, the relation in Eq. (B.8) is slightly modified.

$T \gg m_i$		$T \ll m_i$
fermion	boson	
$n_i = \frac{3}{4}g_i \left(\frac{\zeta(3)}{\pi^2}\right) T^3$	$n_i = g_i \left(\frac{\zeta(3)}{\pi^2}\right) T^3$	$n_i = g_i \left(\frac{m_i T}{2\pi}\right)^{3/2} \exp\left(-\frac{m_i}{T}\right)$
$\rho_i = \frac{7}{8}g_i \left(\frac{\pi^2}{30}\right) T^4$	$\rho_i = g_i \left(\frac{\pi^2}{30}\right) T^4$	$\rho_i = m_i n_i$
$p_i = \frac{1}{3}\rho_i$	$p_i = \frac{1}{3}\rho_i$	$p_i = T n_i (\ll \rho_i)$

Table B.1: The number density n_i , energy density ρ_i and pressure p_i of the particle i , which is thermal equilibrium, in the limits of $T \gg m_i$ and $T \ll m_i$. We have assumed $|\mu_i| \ll T$ and $|\mu_i| < m_i$.

Because the energy density of a non-relativistic particle is exponentially suppressed compared with the relativistic one, the total energy density of the radiation ρ_{rad} is given by the following simple form:

$$\rho_{\text{rad}} = \frac{\pi^2}{30} g_*(T) T^4, \quad (\text{B.14})$$

where

$$g_*(T) \equiv \sum_{\substack{m_i \ll T \\ i = \text{boson}}} g_i + \frac{7}{8} \sum_{\substack{m_j \ll T \\ j = \text{fermion}}} g_j. \quad (\text{B.15})$$

If there are particles which have different temperatures from that of the photon T , another factor $(T_i/T)^4$ should be multiplied in the above expression. (For example, at $T \ll \text{MeV}$, neutrinos have temperature $T_\nu = (4/11)^{1/3} T$ for $m_\nu \ll T_\nu$.)

Notice that all leptogenesis mechanisms discussed in this thesis work at temperatures far above the electroweak scale $T \gg 1 \text{ TeV}$, where all the MSSM particles are expected to be in thermal equilibrium. In this case, we obtain

$$g_* = 228.75 \quad \text{for MSSM}. \quad (\text{B.16})$$

In the expanding universe, it is convenient to introduce the entropy density s , which is defined by

$$\begin{aligned} s &\equiv \frac{\rho + p}{T} \\ &= \frac{4}{3T} \rho = \frac{2\pi^2}{45} g_*(T) T^3. \end{aligned} \quad (\text{B.17})$$

(Again, in the presence of particles with different temperatures, a factor of $(T_i/T)^3$ is multiplied in Eq. (B.17). In this case, the g_* in Eq. (B.17) becomes slightly different from the g_* in Eq. (B.14).) Notice that the entropy per comoving volume sR^3 is conserved as far as no entropy production takes place. Thus it is quite convenient to take the ratio n_X/s when we discuss some number density n_X . For example, if some X -number is conserved, the ratio of the X -number density to the entropy density takes a constant value

$$\frac{n_X}{s} = \text{const} , \quad (\text{B.18})$$

as long as there is no entropy production, since both n_X and s scales as R^{-3} as the universe expands. As another example, if the X -particle is in thermal equilibrium and relativistic ($T \gg m_X$), the ratio is given by

$$\frac{n_X^{\text{eq}}}{s} = \frac{45\zeta(3)}{2\pi^4} \frac{g_X}{g_*(T)} \quad \left(\times \frac{3}{4} \quad \text{for fermion} \right) , \quad (\text{B.19})$$

where the temperature (or time) dependence only comes from $g_*(T)$.

Before closing this section, we calculate the relations between the particle number asymmetry $n_i^{(+)} - n_i^{(-)}$ and the particle's chemical potential μ_i , which can be obtained by integrating Eq. (B.11). In order to calculate the asymmetry, it is necessary to calculate higher order terms than those in Table B.1. The results are given by

$$\begin{aligned} n_i^{(+)} - n_i^{(-)} &= \frac{1}{6} g_i T^3 \left[\left(\frac{\mu_i}{T} \right) + \dots \right] & \text{for fermion} , \\ n_i^{(+)} - n_i^{(-)} &= \frac{1}{3} g_i T^3 \left[\left(\frac{\mu_i}{T} \right) + \dots \right] & \text{for boson} , \end{aligned} \quad (\text{B.20})$$

where ellipses denote higher order terms in the expansions of m_i/T and μ_i/T . Here, we have assumed no Bose-Einstein condensation $|\mu_i| < m_i$ for boson, and relativistic limit $m_i \ll T$. Eq. (B.20) is used in Sec. 1.4, in deriving the relation between the baryon and lepton asymmetries in the presence of sphaleron effect. In this case, the chemical potentials are of order $|\mu| \sim 10^{-10} T$, and hence the above assumptions are justified.

Bibliography

- [1] A. D. Sakharov, Pisma Zh. Eksp. Teor. Fiz. **5** (1967) 32 [JETP Lett. **5** (1967) 24].
- [2] M. Yoshimura, Phys. Rev. Lett. **41** (1978) 281 [Erratum-ibid. **42** (1978) 746];
A. Y. Ignatev, N. V. Krasnikov, V. A. Kuzmin and A. N. Tavkhelidze, Phys. Lett. B **76** (1978) 436.
- [3] D. Toussaint, S. B. Treiman, F. Wilczek and A. Zee, Phys. Rev. D **19** (1979) 1036;
S. Weinberg, Phys. Rev. Lett. **42** (1979) 850;
M. Yoshimura, Phys. Lett. B **88** (1979) 294.
See also S. Dimopoulos and L. Susskind, Phys. Rev. D **18** (1978) 4500.
- [4] H. Georgi and S. L. Glashow, Phys. Rev. Lett. **32** (1974) 438;
J. C. Pati and A. Salam, Phys. Rev. D **10** (1974) 275.
- [5] G. 't Hooft, Phys. Rev. Lett. **37** (1976) 8; Phys. Rev. D **14** (1976) 3432 [Erratum-ibid. D **18** (1976) 2199].
- [6] V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. B **155** (1985) 36.
- [7] A. G. Cohen, D. B. Kaplan and A. E. Nelson, Ann. Rev. Nucl. Part. Sci. **43** (1993) 27 [arXiv:hep-ph/9302210], and references therein.
- [8] For recent reviews and references, see, for example,
J. M. Cline, arXiv:hep-ph/9902328,
M. Quiros, Nucl. Phys. Proc. Suppl. **101** (2001) 401 [arXiv:hep-ph/0101230].
- [9] M. Fukugita and T. Yanagida, Phys. Lett. **B174** (1986) 45.
- [10] Y. Fukuda *et al.* [Super-Kamiokande Collaboration],
Phys. Lett. B **433** (1998) 9 [arXiv:hep-ex/9803006];
Phys. Lett. B **436** (1998) 33 [arXiv:hep-ex/9805006];
Phys. Rev. Lett. **81** (1998) 1562 [arXiv:hep-ex/9807003].

See also a recent data,

T. Nakaya [SUPER-KAMIOKANDE Collaboration], arXiv:hep-ex/0209036.

- [11] Y. Fukuda *et al.* [Super-Kamiokande Collaboration],
Phys. Rev. Lett. **81** (1998) 1158 [Erratum-ibid. **81** (1998) 4279] [arXiv:hep-ex/9805021];
Phys. Rev. Lett. **82** (1999) 2430 [arXiv:hep-ex/9812011];
Phys. Rev. Lett. **82** (1999) 1810 [arXiv:hep-ex/9812009];
Q. R. Ahmad *et al.* [SNO Collaboration], Phys. Rev. Lett. **87** (2001) 071301 [arXiv:nucl-ex/0106015];
B. T. Cleveland *et al.*, Astrophys. J. **496** (1998) 505;
V. N. Gavrin [SAGE Collaboration], Nucl. Phys. Proc. Suppl. **91** (2001) 36;
E. Bellotti, Nucl. Phys. Proc. Suppl. **91** (2001) 44.
- [12] K. Eguchi *et al.*, [KamLAND Collaboration], arXiv:hep-ex/0212021.
- [13] T. Yanagida, in Proceedings of the “*Workshop on the Unified Theory and the Baryon Number in the Universe*”, Tsukuba, Japan, 1979, edited by O. Sawada and A. Sugamoto, KEK Report No. KEK-79-18, p. 95; Prog. Theor. Phys. **64** (1980) 1103;
M. Gell-Mann, P. Ramond and R. Slansky, in “*Supergravity*”, edited by D.Z. Freedman and P. van Nieuwenhuizen (North-Holland, Amsterdam, 1979).
- [14] J. Wess and B. Zumino, Nucl. Phys. B **70** (1974) 39.
For a review and references of the supersymmetric extension of the standard model, see, for example, S. P. Martin, arXiv:hep-ph/9709356.
- [15] J. R. Ellis, S. Kelley and D. V. Nanopoulos, Phys. Lett. B **260** (1991) 131;
U. Amaldi, W. de Boer and H. Furstenuau, Phys. Lett. B **260** (1991) 447;
P. Langacker and M. x. Luo, Phys. Rev. D **44** (1991) 817;
C. Giunti, C. W. Kim and U. W. Lee, Mod. Phys. Lett. A **6** (1991) 1745.
- [16] N. Sakai, Z. Phys. C **11** (1981) 153;
S. Dimopoulos and H. Georgi, Nucl. Phys. B **193** (1981) 150.
- [17] K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, Prog. Theor. Phys. **68** (1982) 927 [Erratum-ibid. **70** (1982) 330].
L. E. Ibanez and G. G. Ross, Phys. Lett. B **110** (1982) 215.
- [18] For a review and references, see, G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept. **267** (1996) 195 [arXiv:hep-ph/9506380].

- [19] M. Y. Khlopov and A. D. Linde, Phys. Lett. **B138** (1984) 265;
 J. Ellis, J. E. Kim and D. V. Nanopoulos, Phys. Lett. **B145** (1984) 181;
 M. Kawasaki and T. Moroi, Prog. Theor. Phys. **93** (1995) 879 [arXiv:hep-ph/9403364].
 see also a recent analysis,
 M. Kawasaki, K. Kohri and T. Moroi, Phys. Rev. D **63** (2001) 103502 [arXiv:hep-ph/0012279].
- [20] T. Moroi, H. Murayama and M. Yamaguchi, Phys. Lett. B **303** (1993) 289;
 A. de Gouvea, T. Moroi and H. Murayama, Phys. Rev. D **56** (1997) 1281 [arXiv:hep-ph/9701244].
- [21] M. H. Reno and D. Seckel, Phys. Rev. D **37** (1988) 3441;
 S. Dimopoulos, R. Esmailzadeh, L. J. Hall and G. D. Starkman, Astrophys. J. **330** (1988) 545; Nucl. Phys. B **311** (1989) 699.
 see also a recent analysis,
 K. Kohri, Phys. Rev. D **64** (2001) 043515 [arXiv:astro-ph/0103411].
- [22] For recent reviews and references, see, for example,
 M. Plumacher, Nucl. Phys. B **530** (1998) 207 [arXiv:hep-ph/9704231];
 W. Buchmuller and M. Plumacher, Phys. Rept. **320** (1999) 329 [arXiv:hep-ph/9904310];
 W. Buchmuller and M. Plumacher, Int. J. Mod. Phys. A **15** (2000) 5047 [arXiv:hep-ph/0007176].
- [23] K. Kumekawa, T. Moroi and T. Yanagida, Prog. Theor. Phys. **92** (1994) 437 [arXiv:hep-ph/9405337];
 G. Lazarides, Springer Tracts Mod. Phys. **163** (2000) 227 [arXiv:hep-ph/9904428] and references therein;
 G. F. Giudice, M. Peloso, A. Riotto and I. Tkachev, JHEP **9908** (1999) 014 [arXiv:hep-ph/9905242];
 T. Asaka, K. Hamaguchi, M. Kawasaki and T. Yanagida, Phys. Lett. B **464** (1999) 12 [arXiv:hep-ph/9906366], Phys. Rev. D **61** (2000) 083512 [arXiv:hep-ph/9907559];
 M. Kawasaki, M. Yamaguchi and T. Yanagida, Phys. Rev. D **63** (2001) 103514 [arXiv:hep-ph/0011104].
- [24] H. Murayama, H. Suzuki, T. Yanagida and J. Yokoyama, Phys. Rev. Lett. **70** (1993) 1912.

- [25] H. Murayama and T. Yanagida, Phys. Lett. B **322** (1994) 349 [arXiv:hep-ph/9310297].
- [26] K. Hamaguchi, H. Murayama and T. Yanagida, Phys. Rev. D **65** (2002) 043512 [arXiv:hep-ph/0109030].
- [27] I. Affleck and M. Dine, Nucl. Phys. B **249** (1985) 361.
- [28] See, for example, E. Kolb and M. Turner, *The Early Universe* (Addison-Wisley, 1990).
- [29] D. E. Groom *et al.* [Particle Data Group Collaboration], “Review of particle physics,” Eur. Phys. J. C **15** (2000) 1.
- [30] N. S. Manton, Phys. Rev. D **28** (1983) 2019.
- [31] F. R. Klinkhamer and N. S. Manton, Phys. Rev. D **30** (1984) 2212.
- [32] This solution was also found in the following papers in different contexts:
R. F. Dashen, B. Hasslacher and A. Neveu, Phys. Rev. D **10** (1974) 4138;
J. Boguta, Phys. Rev. Lett. **50** (1983) 148;
J. Burzlaff, Nucl. Phys. B **233** (1984) 262;
P. Forgacs and Z. Horvath, Phys. Lett. B **138** (1984) 397.
- [33] See also A. D. Linde, Phys. Lett. B **70** (1977) 306, where a possible unsuppression of the baryon number violation via the tunneling at $T \sim T_C$ was briefly commented, and S. Dimopoulos and L. Susskind, in Ref. [3], where it was pointed out that the wash-out of the baryon asymmetry due to the tunneling between the distinct $(B+L)$ vacua could be efficient at temperature higher than the electroweak scale.
- [34] P. Arnold and L. D. McLerran, Phys. Rev. D **36** (1987) 581.
- [35] H. V. Klapdor-Kleingrothaus, S. Kolb and V. A. Kuzmin, Phys. Rev. D **62** (2000) 035014 [arXiv:hep-ph/9909546], and references therein.
- [36] P. Arnold, D. Son and L. G. Yaffe, Phys. Rev. D **55** (1997) 6264 [arXiv:hep-ph/9609481];
D. Bodeker, Phys. Lett. B **426** (1998) 351 [arXiv:hep-ph/9801430];
D. Bodeker, G. D. Moore and K. Rummukainen, Phys. Rev. D **61** (2000) 056003 [arXiv:hep-ph/9907545];
G. D. Moore, arXiv:hep-ph/0009161, and references therein.

- [37] S. Y. Khlebnikov and M. E. Shaposhnikov, Nucl. Phys. **B308** (1988) 885;
J. A. Harvey and M. S. Turner, Phys. Rev. **D 42** (1990) 3344.
- [38] S. Y. Khlebnikov and M. E. Shaposhnikov, Phys. Lett. B **387** (1996) 817 [arXiv:hep-ph/9607386];
M. Laine and M. E. Shaposhnikov, Phys. Rev. D **61** (2000) 117302 [arXiv:hep-ph/9911473], and references therein.
- [39] T. Moroi, arXiv:hep-ph/9503210.
- [40] S. Weinberg, Phys. Rev. Lett. **48** (1982) 1303.
- [41] M. Bolz, A. Brandenburg and W. Buchmuller, Nucl. Phys. B **606** (2001) 518 [arXiv:hep-ph/0012052].
- [42] M. Kawasaki, K. Kohri and T. Moroi, in Ref. [19].
- [43] L. Randall and R. Sundrum, Nucl. Phys. B **557** (1999) 79 [arXiv:hep-th/9810155];
G. F. Giudice, M. A. Luty, H. Murayama and R. Rattazzi, JHEP **9812** (1998) 027 [arXiv:hep-ph/9810442];
J. A. Bagger, T. Moroi and E. Poppitz, JHEP **0004** (2000) 009 [arXiv:hep-th/9911029].
- [44] K. Kohri, in Ref. [21].
- [45] M. Dine, A. E. Nelson and Y. Shirman, Phys. Rev. D **51** (1995) 1362 [arXiv:hep-ph/9408384].
M. Dine, A. E. Nelson, Y. Nir and Y. Shirman, Phys. Rev. D **53** (1996) 2658 [arXiv:hep-ph/9507378].
For a review, see, for example, G. F. Giudice and R. Rattazzi, Phys. Rept. **322** (1999) 419 [arXiv:hep-ph/9801271].
- [46] Y. Nomura and T. Yanagida, Phys. Lett. B **487** (2000) 140 [arXiv:hep-ph/0005211].
Y. Nomura and K. Suzuki, arXiv:hep-ph/0110040.
- [47] H. Pagels and J. R. Primack, Phys. Rev. Lett. **48** (1982) 223.
- [48] M. Bolz, W. Buchmüller and M. Plümacher, Phys. Lett. B **443** (1998) 209 [arXiv:hep-ph/9809381];
T. Gherghetta, G. F. Giudice and A. Riotto, Phys. Lett. B **446** (1999) 28 [arXiv:hep-ph/9808401];
T. Asaka, K. Hamaguchi and K. Suzuki, Phys. Lett. B **490** (2000) 136 [arXiv:hep-ph/0005136].

- [49] M. Kawasaki and T. Moroi, Phys. Lett. B **346** (1995) 27 [arXiv:hep-ph/9408321].
- [50] T. Asaka and T. Yanagida, Phys. Lett. B **494** (2000) 297 [arXiv:hep-ph/0006211].
- [51] M. Fujii and T. Yanagida, arXiv:hep-ph/0208191.
- [52] H. Fujisaki, K. Kumekawa, M. Yamaguchi and M. Yoshimura, Phys. Rev. D **54** (1996) 2494 [arXiv:hep-ph/9511381].
- [53] R. Kallosh, L. Kofman, A. D. Linde and A. Van Proeyen, Phys. Rev. D **61** (2000) 103503 [arXiv:hep-th/9907124];
G. F. Giudice, I. Tkachev and A. Riotto, JHEP **9908** (1999) 009 [arXiv:hep-ph/9907510].
- [54] H. P. Nilles, M. Peloso and L. Sorbo, Phys. Rev. Lett. **87** (2001) 051302 [arXiv:hep-ph/0102264].
- [55] M. Flanz, E. A. Paschos and U. Sarkar, Phys. Lett. B **345** (1995) 248 [Erratum-ibid. B **384** (1995) 487] [arXiv:hep-ph/9411366];
L. Covi, E. Roulet and F. Vissani, Phys. Lett. B **384** (1996) 169 [arXiv:hep-ph/9605319];
W. Buchmuller and M. Plumacher, Phys. Lett. B **431** (1998) 354 [arXiv:hep-ph/9710460].
- [56] L. Covi, E. Roulet and F. Vissani, in Ref. [55]
- [57] B. A. Campbell, S. Davidson and K. A. Olive, Nucl. Phys. B **399** (1993) 111 [arXiv:hep-ph/9302223].
- [58] M. Plumacher, in Ref. [22].
- [59] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870.
- [60] W. Buchmuller and T. Yanagida, Phys. Lett. B **445** (1999) 399 [arXiv:hep-ph/9810308].
- [61] For a discussion and references, see A. Pilaftsis, Int. J. Mod. Phys. A **14** (1999) 1811 [arXiv:hep-ph/9812256].
- [62] M. A. Luty, Phys. Rev. D **45** (1992) 455.
- [63] W. Buchmuller, P. Di Bari and M. Plumacher, Nucl. Phys. B **643** (2002) 367 [arXiv:hep-ph/0205349]; Phys. Lett. B **547** (2002) 128 [arXiv:hep-ph/0209301]. See also P. Di Bari, arXiv:hep-ph/0211175.

- [64] A. D. Dolgov and Y. B. Zeldovich, *Rev. Mod. Phys.* **53** (1981) 1.
- [65] E. W. Kolb and S. Wolfram, *Nucl. Phys. B* **172** (1980) 224 [Erratum-ibid. B **195** (1980) 542].
- [66] W. Buchmuller, [arXiv:hep-ph/0101102](#).
- [67] M. Fujii, K. Hamaguchi and T. Yanagida, *Phys. Rev. D* **65** (2002) 115012 [[arXiv:hep-ph/0202210](#)].
- [68] C. D. Froggatt and H. B. Nielsen, *Nucl. Phys. B* **147** (1979) 277.
- [69] W. Buchmuller and T. Yanagida, *Phys. Lett. B* **302** (1993) 240.
- [70] S. Davidson and A. Ibarra, *Phys. Lett. B* **535** (2002) 25 [[arXiv:hep-ph/0202239](#)].
- [71] T. Asaka, K. Hamaguchi, M. Kawasaki and T. Yanagida, in Ref. [23].
- [72] M. Kawasaki, M. Yamaguchi and T. Yanagida, *Phys. Rev. Lett.* **85** (2000) 3572 [[arXiv:hep-ph/0004243](#)].
- [73] M. Kawasaki, M. Yamaguchi and T. Yanagida, in Ref. [23].
- [74] G. F. Giudice, M. Peloso, A. Riotto and I. Tkachev, in Ref. [23].
- [75] M. Fukugita and T. Yanagida, *Phys. Rev. D* **42** (1990) 1285.
- [76] J. Sato and T. Yanagida, *Nucl. Phys. Proc. Suppl.* **77** (1999) 293 [[arXiv:hep-ph/9809307](#)];
P. Ramond, *Nucl. Phys. Proc. Suppl.* **77** (1999) 3 [[arXiv:hep-ph/9809401](#)].
- [77] J. Sato and T. Yanagida, *Phys. Lett. B* **493** (2000) 356 [[arXiv:hep-ph/0009205](#)].
- [78] G. Lazarides, in Ref. [23].
- [79] G. R. Dvali, Q. Shafi and R. Schaefer, *Phys. Rev. Lett.* **73** (1994) 1886 [[arXiv:hep-ph/9406319](#)].
- [80] E. J. Copeland, A. R. Liddle, D. H. Lyth, E. D. Stewart and D. Wands, *Phys. Rev. D* **49** (1994) 6410 [[arXiv:astro-ph/9401011](#)].
- [81] C. Panagiotakopoulos, *Phys. Rev. D* **55** (1997) 7335 [[arXiv:hep-ph/9702433](#)];
A. D. Linde and A. Riotto, *Phys. Rev. D* **56** (1997) 1841 [[arXiv:hep-ph/9703209](#)].
- [82] C. L. Bennett *et al.*, *Astrophys. J.* **464** (1996) L1 [[arXiv:astro-ph/9601067](#)].

- [83] A. D. Linde, Phys. Lett. B **108** (1982) 389;
A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48** (1982) 1220.
- [84] K. I. Izawa and T. Yanagida, Phys. Lett. B **393** (1997) 331 [arXiv:hep-ph/9608359].
- [85] See, for example, A.D. Linde, *Particle Physics and Inflationary Cosmology*, (Harwood, Chur, Switzerland, 1990).
- [86] K. I. Izawa, M. Kawasaki and T. Yanagida, Phys. Lett. B **411** (1997) 249 [arXiv:hep-ph/9707201];
T. Asaka, M. Kawasaki and M. Yamaguchi, Phys. Rev. D **61** (2000) 027303 [arXiv:hep-ph/9906365].
- [87] A. D. Linde, Phys. Lett. B **327** (1994) 208 [arXiv:astro-ph/9402031];
A. Vilenkin, Phys. Rev. Lett. **72** (1994) 3137 [arXiv:hep-th/9402085].
- [88] K. I. Izawa, M. Kawasaki and T. Yanagida, Prog. Theor. Phys. **101** (1999) 1129 [arXiv:hep-ph/9810537].
- [89] N. Sakai, H. A. Shinkai, T. Tachizawa and K. i. Maeda, Phys. Rev. D **53** (1996) 655 [Erratum-ibid. D **54** (1996) 2981] [arXiv:gr-qc/9506068];
I. Cho and A. Vilenkin, Phys. Rev. D **56** (1997) 7621 [arXiv:gr-qc/9708005];
A. A. de Laix, M. Trodden and T. Vachaspati, Phys. Rev. D **57** (1998) 7186 [arXiv:gr-qc/9801016].
- [90] <http://map.gsfc.nasa.gov/>
<http://astro.estec.esa.nl/SA-general/Projects/Planck/>
- [91] M. Yoshimura, Prog. Theor. Phys. **94** (1995) 873 [arXiv:hep-th/9506176];
H. Fujisaki, K. Kumekawa, M. Yamaguchi and M. Yoshimura, Phys. Rev. D **53** (1996) 6805 [arXiv:hep-ph/9508378];
L. Kofman, A. D. Linde and A. A. Starobinsky, Phys. Rev. D **56** (1997) 3258 [arXiv:hep-ph/9704452], and references therein.
- [92] J. Baacke, K. Heitmann and C. Patzold, Phys. Rev. D **58** (1998) 125013 [arXiv:hep-ph/9806205];
P. B. Greene and L. Kofman, Phys. Lett. B **448** (1999) 6 [arXiv:hep-ph/9807339].
- [93] H. Murayama, H. Suzuki, T. Yanagida and J. Yokoyama, Phys. Rev. D **50** (1994) 2356 [arXiv:hep-ph/9311326].
- [94] T. Moroi, H. Murayama and M. Yamaguchi, in Ref. [20].

- [95] M. Fujii, K. Hamaguchi and T. Yanagida, Phys. Rev. D **63** (2001) 123513 [arXiv:hep-ph/0102187].
- [96] A. Hebecker, J. March-Russell and T. Yanagida, arXiv:hep-ph/0208249.
- [97] R. Allahverdi, B. A. Campbell and J. Ellis, Nucl. Phys. **B579** (2000) 355 [arXiv:hep-ph/0001122].
- [98] A. Anisimov and M. Dine, Nucl. Phys. B **619** (2001) 729 [arXiv:hep-ph/0008058].
See also, A. Anisimov, arXiv:hep-ph/0111233.
- [99] T. Moroi and H. Murayama, arXiv:hep-ph/0211019.
- [100] T. Moroi and T. Takahashi, Phys. Lett. B **522** (2001) 215 [Erratum-ibid. B **539** (2002) 303] [arXiv:hep-ph/0110096]; Phys. Rev. D **66** (2002) 063501 [arXiv:hep-ph/0206026], and references therein.
- [101] T. Asaka, M. Fujii, K. Hamaguchi and T. Yanagida, Phys. Rev. D **62** (2000) 123514 [arXiv:hep-ph/0008041].
- [102] M. Fujii, K. Hamaguchi and T. Yanagida, Phys. Rev. D **65** (2002) 043511 [arXiv:hep-ph/0109154].
- [103] M. Fujii, K. Hamaguchi and T. Yanagida, Phys. Lett. B **538** (2002) 107 [arXiv:hep-ph/0203189].
- [104] M. Dine, L. Randall and S. Thomas, Phys. Rev. Lett. **75** (1995) 398 [arXiv:hep-ph/9503303]. Nucl. Phys. B **458** (1996) 291 [arXiv:hep-ph/9507453].
- [105] T. Moroi and H. Murayama, JHEP **0007** (2000) 009 [arXiv:hep-ph/9908223].
- [106] M. Fujii, K. Hamaguchi and T. Yanagida, Phys. Rev. D **64** (2001) 123526 [arXiv:hep-ph/0104186].
- [107] M. Fujii and K. Hamaguchi, Phys. Lett. B **525** (2002) 143 [arXiv:hep-ph/0110072]; Phys. Rev. D **66** (2002) 083501 [arXiv:hep-ph/0205044].
- [108] B. A. Campbell, S. Davidson and K. A. Olive, Phys. Lett. B **303** (1993) 63 [arXiv:hep-ph/9302222].
- [109] M. Senami and K. Yamamoto, Phys. Lett. B **524** (2002) 332 [arXiv:hep-ph/0105054].

- [110] M. Senami and K. Yamamoto, Phys. Rev. D **66**, 035006 (2002) [arXiv:hep-ph/0205041]; arXiv:hep-ph/0210073.
- [111] M. K. Gaillard, H. Murayama and K. A. Olive, Phys. Lett. B **355** (1995) 71 [arXiv:hep-ph/9504307].
- [112] J. Grundberg, T. H. Hansson and U. Lindstrom, arXiv:hep-th/9510045.
- [113] A. de Gouvêa, T. Moroi and H. Murayama, in Ref. [20].
- [114] T. Asaka, Phys. Lett. B **521** (2001) 329 [arXiv:hep-ph/0110073].
- [115] V. Barger and D. Marfatia, arXiv:hep-ph/0212126;
G. L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo and A. M. Rotunno, arXiv:hep-ph/0212127;
M. Maltoni, T. Schwetz and J. W. Valle, arXiv:hep-ph/0212129;
A. Bandyopadhyay, S. Choubey, R. Gandhi, S. Goswami and D. P. Roy, arXiv:hep-ph/0212146;
J. N. Bahcall, M. C. Gonzalez-Garcia and C. Pena-Garay, arXiv:hep-ph/0212147;
H. Nunokawa, W. J. C. Teves, R. Zukanovich Funchal, arXiv:hep-ph/0212202;
P. Aliani, V. Antonelli, M. Picariello, E. Torrente-Lujan arXiv:hep-ph/0212212.
- [116] M. Apollonio *et al.*, Phys. Lett. **B466** (1999) 415 [arXiv:hep-ex/9907037].
- [117] R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38** (1977) 1440; Phys. Rev. D **16** (1977) 1791.
- [118] See, for a review, J. E. Kim, Phys. Rept. **150** (1987) 1.
- [119] K. Rajagopal, M. S. Turner and F. Wilczek, Nucl. Phys. B **358** (1991) 447.
- [120] See, for the present status of the double beta decay experiments, S. R. Elliott and P. Vogel, arXiv:hep-ph/0202264, and references therein.